

Streptococcus pneumoniae and *Streptococcus pyogenes* Resistant to Macrolides but Sensitive to Clindamycin: a Common Resistance Pattern Mediated by an Efflux System

JOYCE SUTCLIFFE,* AMELIA TAIT-KAMRADT, AND LILLIAN WONDRACK

Department of Infectious Diseases, Central Research Division, Pfizer, Inc., Groton, Connecticut 06340

Received 28 March 1996/Returned for modification 3 May 1996/Accepted 29 May 1996

Macrolide-resistant *Streptococcus pyogenes* isolates from Finland, Australia, and the United Kingdom and, more recently, *Streptococcus pneumoniae* and *S. pyogenes* strains from the United States were shown to have an unusual resistance pattern to macrolides, lincosamides, and streptogramin B antibiotics. This pattern, referred to as M resistance, consists of susceptibility to clindamycin and streptogramin B antibiotics but resistance to 14- and 15-membered macrolides. An evaluation of the macrolide-lincosamide-streptogramin B resistance phenotypes among our streptococcal strains collected from 1993 to 1995 suggested that this unusual resistance pattern is not rare. Eighty-five percent ($n = 66$) of the *S. pneumoniae* and 75% ($n = 28$) of the *S. pyogenes* strains in our collection had an M phenotype. The mechanism of M resistance was not mediated by target modification, as isolated ribosomes from a pneumococcal strain bearing the M phenotype were fully sensitive to erythromycin. Further, the presence of an *erm* methylase was excluded with primers specific for an *erm* consensus sequence. However, results of studies that determined the uptake and incorporation of radio-labeled erythromycin into cells were consistent with the presence of a macrolide efflux determinant. The putative efflux determinant in streptococci seems to be distinct from the multicomponent macrolide efflux system in coagulase-negative staphylococci. The recognition of the prevalence of the M phenotype in streptococci has implications for sensitivity testing and may have an impact on the choice of antibiotic therapy in clinical practice.

There are presently three recognized mechanisms of resistance to macrolide antibiotics: target modification, inactivation, and efflux (36–38, 72). Target modification occurs at the level of the ribosomes via an *erm* gene encoding a 23S rRNA methylase (36, 38, 72). There are currently at least eight classes of *erm* genes distinguishable by hybridization criteria (36, 38, 72). *Erm* methylases add either one or two methyl residues to a highly conserved adenine residue in domain V, the peptidyl transferase center, of 23S rRNA (33, 72). This modification renders the strain resistant to most macrolides, lincosamides, and streptogramin B compounds; phenotypically, this resistance pattern is known as MLS_B resistance (21, 33). Expression of MLS_B resistance can be inducible or constitutive and is unrelated to the class of an *erm* determinant (10, 18, 36, 71, 73). Regulation of the methylase depends upon the sequence of the regulatory domain upstream of the structural gene and is accounted for by a translational attenuation mechanism (10, 18, 36, 71, 73).

Resistance to MLS antibiotics due to inactivation has been described for a number of clinically important organisms, including *Staphylococcus aureus* (12, 16, 35, 39, 40, 75) *Staphylococcus haemolyticus* (12, 35), and *Escherichia coli* (1, 3, 4, 9, 31, 51–54). To our knowledge, no form of macrolide inactivation has been described for streptococci.

Staphylococci appear to have an efflux system (19, 28, 37, 42, 43, 46, 59–61, 75) with specificity for 14- and 15-membered macrolides and type B streptogramin molecules but not for lincosamide antibiotics (the MS phenotype). The efflux system appears to be multicomponent, involving *msrA* (60, 61), *smp*

(59) and/or *stp* (59) genes to constitute a fully operational efflux pump. *msrA* and *stp* genes encode ATP-binding proteins homologous to a superfamily of related proteins that contain two conserved ATP-binding sites that are involved in transport (27, 41, 49), while *smp* genes encode a hydrophobic transmembrane protein, the putative pump. Presently, it is unclear what role each protein plays in the functional efflux system; however, it is clear that *MsrA* must be present to confer the macrolide and streptogramin B resistance (MS) phenotype (59). Efflux has not been described for streptococci.

Macrolide resistance via an energy-dependent *mtr* efflux system in gonococci was recently reported (26, 55). This efflux system, similar to the *mexAB-oprK* efflux system of *Pseudomonas aeruginosa* and the *acrAB* and *acrEF* efflux pumps of *E. coli*, is not specific for macrolides (49). Rather, these pumps appear to be important for preventing entry of structurally diverse, but undesirable agents, thereby enhancing the survival of the organism in its ecological niche.

Clinical *Streptococcus pyogenes* strains from Finland (63, 64), Australia (67), the United Kingdom (58, 62, 77), and North America (15) have been described with an M phenotype, that is, resistance to macrolides but susceptibility to lincosamide and streptogramin B antibiotics. A study of pneumococci recovered from middle ear and sinus cultures of children from Houston, Texas, noted isolates with an M phenotype (48). Evaluation of our clinical streptococcal strains revealed that the majority of *S. pyogenes* and *Streptococcus pneumoniae* strains had an M phenotype. Our studies showed that the determinant responsible for the M phenotype appears to be distinct from known *erm* genes and is not mediated through target modification. Rather, we found that strains bearing the M phenotype contain an efflux system for erythromycin apparently distinct from the efflux system described for erythromycin-resistant staphylococci.

* Corresponding author. Mailing address: Department of Infectious Diseases, Central Research Division, Pfizer, Inc., Eastern Point Rd., Groton, CT 06340. Phone: (860) 441-4693. Fax: (860) 441-6159.

TABLE 1. Macrolide resistance in *S. pneumoniae*

Yr	Source	No. of strains	MLS _B phenotype ^a	Efflux demonstrated ^b	PCR or hybridization analysis ^c			No. of Pen ^f strains (I/H) ^d
					<i>erm</i> class	<i>msr</i>	<i>smp</i>	
1993	Kentucky	2	M	NT ^e	None	No	No	0/1
	Texas	8	M	NT	None	NT	NT	5/3
	France	2	cMLS _B	No	<i>B</i>	No	No	1/1
1994	Alabama	1	M	NT	None	No	NT	1/0
	Arizona	3	M	NT	None	No	No	1/2
	California	6	M	NT	None	No	No	3/1
	California	3	cMLS _B	NT	<i>B</i>	NT	NT	2/1
	Maryland	4	M	Yes	None	NT	NT	0/2
	Missouri	3	M	NT	None	No	No	1/0
	Ohio	1	M	NT	None	NT	NT	0/0
	Virginia	1	M	NT	None	No	NT	1/0
	Texas	5	M	Yes	None	No	No	1/3
	France	1	cMLS _B	No	<i>B</i>	NT	NT	0/1
1995	Alabama	1	M	NT	None	No	NT	0/0
	California	1	M	NT	None	No	NT	0/1
	Kentucky	12	M	NT	None	No	No	2/7
	Massachusetts	1	M	NT	None	No	NT	0/0
	Michigan	1	M	NT	None	No	NT	1/0
	Nebraska	4	M	NT	None	No	No	3/0
	Nebraska	1	cMLS _B	NT	<i>B</i>	No	No	1/0
	Oregon	1	cMLS _B	NT	<i>B</i>	No	NT	1/0
	South Africa	2	M	NT	None	No	No	0/1
	South Africa	2	cMLS _B	NT	<i>B</i>	No	No	0/2

^a Results are from a disk assay. cMLS_B, constitutive resistance to erythromycin, clindamycin, and streptogramin B; M, erythromycin resistance but clindamycin and streptogramin B susceptibility.

^b Efflux was demonstrated (yes) or found not to be present (no) by radiolabeled uptake studies.

^c None, no PCR product was obtained with *erm*-specific or *erm*-degenerate primer sets; *B*, *ermB* was present; No, the indicated gene was not present by PCR (*msr*) or hybridization (*smp*) analysis.

^d Number of strains with intermediate (I) or high-level (H) resistance to penicillin as described by the report of the National Committee for Clinical Laboratory Standards (47).

^e NT, not tested.

MATERIALS AND METHODS

Chemicals. Erythromycin and clindamycin disks were obtained from Difco Laboratories (Detroit, Mich.). Disks containing 25 µg of a streptogramin B analog, CP-37,277 (13), were freshly made by adding 25 µl of CP-37,277 (stock solution of 1 mg/ml in ethanol) to sterile disks. Brain heart infusion and Todd-Hewitt broth were obtained from Difco Laboratories; brain heart infusion and blood agar plates were purchased from Remel Microbiology Products (Lenexa, Kans.). PCR buffers and DNA polymerase were supplied by Perkin-Elmer (Danbury, Conn.), and all DNA primers were supplied by GenoSys Biotechnologies (The Woodlands, Tex.). Lysozyme and lysostaphin were purchased from Sigma Chemical Co. (St. Louis, Mo.), and proteinase K (biotechnology grade) was from Amresco (Solon, Ohio). Azithromycin (17) was prepared in the laboratories of Pfizer Central Research, Groton, Conn. Erythromycin, carbonyl cyanide *m*-

chlorophenylhydrazone (CCCP), sodium arsenate, and all chemicals required for ribosome isolation and in vitro protein translation except tRNA and radiolabeled lysine were purchased from Sigma Chemical Co. Dithiothreitol and tRNA were purchased from Boehringer Mannheim (Indianapolis, Ind.). ReadySafe cocktail was purchased from Beckman (Fullerton, Calif.).

Radiochemicals. [*N*-methyl-¹⁴C]erythromycin (55 mCi/mmol) was purchased from DuPont, NEN Research Products (Boston, Mass.). [¹⁴C]lysine (312 mCi/mmol) was purchased from Amersham (Arlington Heights, Ill.). MicroScint 20 was purchased from Packard (Meriden, Conn.).

Bacterial strains. *S. pneumoniae* and *S. pyogenes* strains, collected between 1993 and 1995 from the United States and other areas of the world, are described in Tables 1 and 2. Control strains not included in the tables are as follows. *Staphylococcus aureus* RN4220, a restriction-deficient strain that serves as the

TABLE 2. Macrolide resistance in *S. pyogenes*

Yr	Source	No. of strains	MLS _B phenotype ^a	Efflux demonstrated ^b	PCR or hybridization analysis ^c		
					<i>erm</i> class	<i>msr</i>	<i>smp</i>
1994	Ireland	9	M	Yes	None	No	No
	Ireland	2	cMLS _B	No	None	No	No
	Ireland	2	iMLS _B	NT ^d	Variante	NT	NT
	Ireland	1	iMLS _B	No	<i>B</i>	NT	NT
	Indiana	7	M	NT	None	No	No
	Indiana	1	iMLS _B	NT	Variante	No	No
1995	Sweden	5	M	NT	None	No	No
	Sweden	1	iMLS _B	NT	Variante	NT	NT

^a See Table 1, footnote a. For the iMLS_B phenotype, MLS_B resistance was fully expressed only after induction of the *erm* operon.

^b See Table 1, footnote b.

^c See Table 1, footnote c. Variante, a PCR product was detected only with degenerate primers.

^d NT, not tested.

source of S150 extracts for *in vitro* translation (32), was obtained from Saleem Kahn (University of Pittsburgh School of Medicine); *Staphylococcus epidermidis* S1187 and *E. coli* XL-1 Blue, containing the 1.9-kb *Hind*III fragment of *msrA* (60) were obtained through the courtesy of Anne Eady (University of Leeds); *S. pyogenes* 02C1061 (strain AC1 [pAC1]; serotype M22:T12; *ermB*⁺ [14]) and 02C1062 (strain AC1 cured of plasmid pAC1 [14]) were obtained from Don Clewell (University of Michigan); *S. pneumoniae* R6, an unencapsulated derivative of the Rockefeller University strain R36A (7), was obtained through the courtesy of Regine Hakenbeck (Max-Planck Institute, Berlin, Germany); *S. pneumoniae* 02J1047 (strain 13433 [8]), an *ermB* strain, was obtained from A. Fremaux (Centre Hospitalier Intercommunal, Creteil, France); and *S. pneumoniae* 02J1016 (25), an MLS_B-susceptible serotype 3 strain, was obtained from J.-J. Pocardalo (Hôpital Claude Bernard, Paris, France).

Strain characterization. All strains were screened for MLS_B resistance by an agar disk diffusion assay. Isolated colonies were selected by a BBL Prompt system (Becton Dickinson, Cockeysville, Md.). This standardized cell suspension served as an inoculum for streaking the surface of a blood agar plate. Clindamycin (2 µg), erythromycin (15 µg), and CP-37,277 (25 µg) disks were placed on the plates approximately 12 mm apart. The plates were incubated overnight at 37°C in an atmosphere containing 5% carbon dioxide. Blunting of the zones around clindamycin or the streptogramin B analog (CP-37,277) indicated inducible resistance to these compounds (36, 38), whereas constitutive resistance was indicated by no zones to clindamycin and CP-37,277 and either a small zone or no zone to erythromycin.

Inactivation of radiolabeled erythromycin in concentrated cell suspensions was assessed as described previously (75, 76).

Macrolide efflux by actively growing cells. Freshly grown colonies from blood agar plates were used to inoculate Todd-Hewitt broth for growth overnight in 5% CO₂ at 37°C. The overnight culture was diluted to an optical density at 650 nm (OD₆₅₀) of 0.1 and grown in 5% CO₂ without shaking until an OD₆₅₀ of 0.3 was reached. At this point, 0.02 µg of erythromycin per ml was added to ensure full induction of the efflux determinant and the culture was incubated for another hour. [¹⁴C]erythromycin was added at a final concentration of 0.2 µg/ml, and uptake was assessed by removing aliquots of the culture in duplicate, filtering each sample onto prewet GF/C filters, and washing the filters twice with 0.9% NaCl containing 1 µg of erythromycin per ml. The amount of radiolabeled erythromycin that is cell associated was determined after scintillation counting of the dried filters. In a typical experiment, the OD₆₅₀ from the addition of [¹⁴C]erythromycin to the end of the experiment (45 min) increased by ~35%. Samples receiving 10 mM arsenate or 25 µM CCCP were exposed to the drug for 10 min prior to the addition of radiolabeled erythromycin; no change in OD₆₅₀ was noted during the course of the experiment.

DNA isolation, PCR, and hybridization conditions. Total genomic DNA was isolated from streptococcal strains as described in reference 6, with slight modifications. For maximal lysis of *S. pneumoniae*, the period of proteinase K-sodium dodecyl sulfate (SDS) digestion at 37°C was increased to 3 h. To provide adequate DNA yields, group A streptococci were incubated with lysozyme (6 mg/ml) at 37°C for 1 h prior to the addition of proteinase K-SDS. For staphylococcal strains (controls), cell walls were digested with lysostaphin (50 µg/ml for 1 h at 37°C) prior to the addition of proteinase K-SDS. The PCR reaction mixture was as recommended by Perkin-Elmer, with the concentration of magnesium being optimized for each primer set (4 mM for *ermA* and *msrA-msrB* primer sets and 2 mM for *ermB*, *ermC*, and consensus *erm* primer sets). DNA in the reaction mixture (20 µl) was denatured at 93°C for 3 min and then annealed at 52°C. The amplification cycles consisted of elongation at 72°C for 60 s, denaturation at 93°C for 60 s, and annealing at 52°C for 60 s. After 35 amplification cycles, the last elongation step was performed at 72°C for 5 min. Primer sets specific for the detection of *ermA*, *ermB*, *ermC*, and *msrA-msrB* were as described previously (68, 75). Primers for *smp* were based on the published sequence of *smpA* (59): 5'-AAATGTGTTAAAAAGAAATC-3' and 5'-TTTGAACATAATATTCATC-3'. These primers, when used on plasmid DNA isolated from *Staphylococcus epidermidis* S1187, produced the expected 616-bp PCR product. The *smp* probe was labeled with a digoxigenin DNA random prime DNA labelling and detection kit (Boehringer Mannheim) and hybridized to DNA on a Magnagraph nylon membrane (Micron Separations Inc., Westboro, Mass.) according to the manufacturer's instructions.

Isolation of *S. pneumoniae* ribosomes. *S. pneumoniae* strains R6 (susceptible) and 02J1175 (M phenotype; one of the strains from Kentucky from 1995) (Table 1) were grown in 5% CO₂ overnight in Todd-Hewitt broth supplemented with 0.05% sodium thioglycolate and 2% neutralized yeast extract at 37°C. The overnight culture was used as an inoculum (20%) in 9 liters of the above-described medium, and cells were harvested after 6 h of aerobic growth at 37°C. Cells were lysed in a 0.01 M Tris (pH 7.5) buffer containing 0.15 M NaCl, 0.01 M MgCl₂, 0.35 M NH₄Cl, 0.25% SDS, 1 µg of DNase I per ml, and 0.25% sodium deoxycholate as described in the report of Swendsen and Johnson (69). The ammonium sulfate precipitation procedure described by Fogel and Sypherd (22) was used to isolate the ribosomes. Isolated ribosomes from the pneumococci were reconstituted with an S150 extract from *Staphylococcus aureus* RN4220. Translation was monitored *in vitro* by a microtiter-based poly(A) system (75).

RESULTS

Phenotypic analysis of streptococcal strains. By the agar disk diffusion assay, 66 erythromycin-resistant strains of *S. pneumoniae* and 28 erythromycin-resistant *S. pyogenes* strains were evaluated (Tables 1 and 2). Blunting of the zones around clindamycin or the streptogramin B analog (CP-37,277) indicated inducible resistance to these compounds (36, 38), whereas constitutive resistance was indicated by either a small zone or no zone. As described by Seppälä et al. (64), three different phenotypes were distinguishable for the group A isolates: constitutive MLS_B resistance (cMLS_B), inducible MLS_B resistance (iMLS_B), and a novel phenotype which we designate M. The M phenotype is typified by strains remaining susceptible to clindamycin and streptogramin B antibiotics while being resistant to 14- and 15-membered macrolides. The M phenotype was also found in our recent clinical isolates of pneumococci. To our surprise, the M phenotype constituted 75% of the erythromycin-resistant *S. pyogenes* isolates and 85% of the erythromycin-resistant *S. pneumoniae* isolates.

For this study, clinical laboratories were requested to send erythromycin-resistant *S. pneumoniae* and *S. pyogenes* strains. It is interesting that the vast majority of the pneumococci are also penicillin resistant (Pen^r) (76%) and that 52% of the Pen^r isolates show high-level Pen^r (MIC ≥ 2 µg/ml). The association of macrolide and penicillin resistance in *S. pneumoniae* has been noted by others (11, 29, 48, 74). The 10 pneumococci with an MLS_B-resistant phenotype were all coresistant to penicillin, while 71.4% of the isolates bearing an M phenotype were coresistant to penicillin. None of the *S. pyogenes* isolates were resistant to penicillin *in vitro*.

M strains do not have an *erm* methylase. Primer sets designed to specifically detect *ermA*, *ermB*, or *ermC* methylase revealed the presence of the archetypal *ermB* in all of the MLS_B-resistant strains of *S. pneumoniae* (Table 1). However, only one of seven MLS_B-resistant strains of *S. pyogenes* gave a specific PCR product with the *ermB*-specific primers. With a set of degenerate primers (5) designed to detect *ermA*, *ermB*, *ermC*, *ermG*, and related variants, four of the seven strains were found to have an *erm*-type sequence (listed as variants in Table 2 because they were not detected with the archetypal *ermB*-specific primers). Interestingly, the two remaining MLS_B-resistant strains of *S. pyogenes* did not give a specific PCR product even with the degenerate primers. The most likely explanation for erythromycin resistance in the strains labeled as having a variant *erm* is that the resident *erm* has a sequence slightly different from that region recognized by the *ermB*-specific primers. The two *S. pyogenes* strains that do not give a PCR product with the degenerate *erm* primer set may have an alteration in the peptidyl transferase center or some other unknown mechanism that confers coresistance to MLS_B antibiotics. In an effort to address the former possible explanation, the copy number of 23S rRNA genes in *S. pyogenes* was investigated; six copies were found in both MLS_B-susceptible and MLS_B-resistant strains (data not shown). Others have found that organisms containing four or less 23S rRNA copies can be phenotypically MLS_B resistant when only one of the rRNA genes contains a base change at the methylatable adenine (A-2058 [72]) in the peptidyl transferase center (45, 57, 72). In *E. coli*, with seven rRNA genes, a base change at A-2058 in only one of the gene copies is phenotypically recessive (72). Given that *S. pyogenes*' copy number was close to seven, it seemed unlikely that MLS_B resistance in these strains could be accounted for by an alteration in the rRNA sequence. However, future studies with these strains are necessary to determine if the determinant(s) is novel.




FIG. 1. Sensitivity of *S. pneumoniae* ribosomes to erythromycin and azithromycin. An *in vitro* translation system was reconstituted with the S150 extract from *Staphylococcus aureus* RN4220 and purified ribosomes from macrolide-susceptible *S. pneumoniae* R6 (■) or 02J1175, an *S. pneumoniae* strain with an M phenotype (●).

Streptococcal strains bearing an M phenotype failed to yield a PCR product with any of the *erm* primer sets, including the consensus *erm* primers. To determine if these strains bear a target modification specific for macrolides, ribosomes were isolated from strain 02J1175, an *S. pneumoniae* strain with an M phenotype. Figure 1 compares the sensitivity of ribosomes isolated from 02J1175 to that of a well-known laboratory pneumococcal strain, R6, in a reconstituted translation assay. Ribosomes isolated from either strain are fully sensitive to the inhibition of either erythromycin or azithromycin (Fig. 1). Thus, it appears that the determinant specifying erythromycin resistance in strain 02J1175 does not alter the binding of macrolides to the ribosome. (Attempts to isolate functional ribosomes from MLS_B-resistant *S. pneumoniae* strains were unsuccessful, but ribosomes isolated from MLS_B-resistant *S. aureus* cells have 50% inhibitory concentrations of >100 µg/ml [75].)

Streptococci with the M phenotype efflux erythromycin. Inactivation of erythromycin by clinical gram-positive strains has been described only for *Staphylococcus aureus* (75). Although degradation was considered a mechanism that could account for the M phenotype, we found no evidence for inactivation of erythromycin by any of the *S. pyogenes* or *S. pneumoniae* strains in our culture collection (reference 76 and data not shown).

The M phenotype in streptococci is reminiscent of the MS phenotype (resistance to macrolide and streptogramin B antibiotics but susceptible to lincosamide antibiotics) described for staphylococci (19, 28, 34, 42, 43, 46, 59–61, 75). To determine if resistance to macrolides is mediated by an efflux mechanism, exponentially growing cells of *S. pyogenes* and *S. pneumoniae* were evaluated for their uptake and incorporation of radiolabeled erythromycin in the absence or in the presence of CCCP or arsenate, agents that disrupt proton motive force. Figures 2 and 3 provide examples of erythromycin uptake by susceptible strains (02C1062 in Fig. 2 and R6 and 02J1016 in Fig. 3), isolates with an M phenotype (02C1063 and 02C1064 in Fig. 2 and 02J1086 and 02J1088 in Fig. 3), and control strains that are resistant to macrolides via an *ermB* methylase (strains 02C1061 and 02J1047 in Fig. 2 and 3, respectively).

The initial rate of incorporation of erythromycin by susceptible streptococci is rapid, after which either a plateau phase (02J1062 in Fig. 2 and R6 in Fig. 3) or a slower rate of uptake (02J1016 in Fig. 3) occurs. Pretreatment with CCCP or arsenate did not greatly alter the accumulation in 02C1062 (Fig. 2) or 02J1016 (Fig. 3), while a more substantial amount of erythromycin is evident in R6 following pretreatment with arsenate (Fig. 3). The increased amount of uptake was noted in other susceptible strains of pneumococci (data not shown); a similar phenomenon has been eloquently explained by Nikaido and Thanassi (50). The cytosolic total concentration of erythromycin is lower than that of the external medium because of the pH gradient (a major component of proton motive force). The addition of CCCP or arsenate collapses the pH gradient, producing an influx of erythromycin as a result of equilibration of free drug (50).

Strains with an M phenotype do not maximally accumulate erythromycin (even after 30 to 45 min) unless they are pretreated with 25 µM CCCP (Fig. 2) or 10 mM arsenate (Fig. 3). The additional accumulation in these strains appears to result from the inhibition of a proton motive force-sensitive efflux pump. This interpretation is made in conjunction with the knowledge that these strains are erythromycin resistant. MICs observed for *S. pneumoniae* or *S. pyogenes* strains with an M phenotype are in the 4 to 16-µg/ml range, distinguishable from MLS_B-susceptible strains (MICs ≤ 0.25 µg/ml).

There is no substantial retention of radiolabeled erythromycin in MLS_B-resistant strains; this is expected because ErmB-modified ribosomes can no longer bind erythromycin. Further, pretreatment of strains containing ErmB with CCCP or arsenate does not alter the small amount of erythromycin accumulated.

Macrolide efflux in M strains appears to be distinct. To determine if the efflux system in streptococci was distinct from the multicomponent efflux pump described for coagulase-negative staphylococci, strains with the M phenotype were probed with *smpA* from *Staphylococcus epidermidis* S1187 (59, 60) and evaluated by PCR using primers specific to the C-terminal ATP-binding region of *msrA* (75). None of the strains ap-




FIG. 2. Efflux studies with *S. pyogenes* strains. Uptake and incorporation of radiolabeled erythromycin in the presence (●) or absence (■) of CCCP are shown.

peared to have either of the staphylococcal pump components (Tables 1 and 2). Additional studies using an *msrA* probe (data not shown) substantiated the results obtained with the *msrA*-specific PCR primers. Thus, it appears that the efflux determinant(s) in *S. pyogenes* and *S. pneumoniae* is distinguishable from the efflux system in staphylococci bearing the MS phenotype.

DISCUSSION

We found that recent clinical isolates of erythromycin-resistant *S. pneumoniae* and *S. pyogenes* are predominantly resistant to 14- and 15-membered macrolides, a resistance phenotype known as M. This is contrary to earlier reports in which the mechanism of erythromycin resistance in streptococci was confined to MLS_B resistance (29, 30, 36, 66, 72). Although other studies have described streptococcal strains that appear to bear the M phenotype (15, 48, 58, 62–64, 67, 77), this is the first report that provides a mechanism for the novel resistance phenotype.

Using PCR analysis, we were able to show that none of the *S. pyogenes* and *S. pneumoniae* strains with an M phenotype contain an *erm* determinant. Further, ribosomes isolated from a pneumococcal strain with an M phenotype proved to be fully sensitive to erythromycin and azithromycin. Several strains were characterized for their ability to take up radiolabeled erythromycin. The M strains appear to have an efflux pump, as discerned by the reduced uptake of [¹⁴C]erythromycin in the

absence of CCCP or arsenate. The putative efflux pump appears to be distinct from the *msr-stp-smp* pump described for staphylococci (37, 59–61, 73). To our knowledge, this is the first report describing a macrolide efflux pump in streptococci.

The M phenotype is not found in a single geographical location (Tables 1 and 2) (15, 24, 48, 58, 62–64, 67, 77). Even recent *S. pneumoniae* isolates from South Africa have the M phenotype, whereas 10 isolates from 1978 were all found to contain an *ermB* methylase (data not shown and reference 29). Thus far, we have found a 100% correlation between the M phenotype and erythromycin resistance mediated by a non-*erm* determinant. Our results are consistent with a putative efflux mechanism for strains with an M phenotype.

Currently, the primary set of antibiotics to be tested by clinical laboratories as recommended by the National Committee for Clinical Laboratory Standards for streptococci (47) does not include lincosamides (i.e., clindamycin) or streptogramin B, primarily because resistance to erythromycin in the past has signaled cross-resistance to these antibiotics. However, results from this epidemiological survey and others (23, 48) may well have an impact on these recommendations, since a streptococcal strain with an M phenotype remains susceptible to clindamycin and streptogramin B. The value in testing these antibiotics as part of the primary evaluation could be especially meaningful in cases in which *S. pneumoniae* has been recovered from patients with life-threatening infections (e.g., meningitis and bacteremia [44]). As regards bacteremia, it is noteworthy that reports of multidrug-resistant pneumococci of




FIG. 3. Efflux studies with *S. pneumoniae* strains. Uptake and incorporation of radiolabeled erythromycin in the presence (●) or absence (■) of arsenate are shown.

diverse serotypes are appearing worldwide (2, 11, 29, 44, 70, 74). It is important that the clinician be outfitted with a full armamentarium for treating these dangerous infections. Further, Nelson et al. (48) and McCracken (44) have suggested that a clinical trial in the United States to evaluate clindamycin as a treatment for middle ear and sinus infections caused by *S. pneumoniae* strains resistant to other oral antibiotics may be rational; a trial to evaluate clindamycin for use in treatment of penicillin-tolerant, macrolide-resistant *S. pyogenes* might also prove useful. Recently, clindamycin therapy was reported to be successful in the treatment of a child with recurrent pneumococcal sepsis caused by an intermediately penicillin-resistant strain (65). A recent evaluation of clindamycin to treat experimental meningitis caused by penicillin- and cephalosporin-resistant *S. pneumoniae* was also reported (56). Because of the poor clinical response to *Haemophilus influenzae*-mediated otitis media (20), it is unlikely that empirical clindamycin use will be advised for patients with otitis media, however.

Evaluation of the novelty of the putative macrolide efflux determinant(s) in *S. pyogenes* and *S. pneumoniae* awaits the cloning, sequencing, and functional analysis of the requisite genes. The determination of the substrate specificity and the appropriate classification of the efflux superfamily (major facilitator, resistance-nodulation-division, staphylococcal multidrug resistance, or ATP-binding cassette [41, 49]) will also be interesting.

ACKNOWLEDGMENTS

We thank Marge Anderson and Jim Retsema for managing the Pfizer strain collection and Juan Walterspiel and Eric Utt for reviewing the manuscript. *S. pneumoniae* strains from Nebraska and Kentucky were collected by Stephen Chartrand, Creighton University School of Medicine, and Stanley Block, Kentucky Pediatric Research, Inc., respectively, and were obtained from Ellen Smith Moland of the Antimicrobials and Biotechnology Department, Creighton University School of Medicine.

REFERENCES

1. Andreumont, A., G. Gerbaud, and P. Courvalin. 1986. Plasmid-mediated high-level resistance to erythromycin in *Escherichia coli*. *Antimicrob. Agents Chemother.* **29**:515-518.
2. Appelbaum, P. C. 1992. Antimicrobial resistance in *Streptococcus pneumoniae*: an overview. *Clin. Infect. Dis.* **15**:77-83.
3. Arthur, M., A. Andreumont, and P. Courvalin. 1987. Distribution of erythromycin esterase and rRNA methylase genes in members of the family *Enterobacteriaceae* highly resistant to erythromycin. *Antimicrob. Agents Chemother.* **31**:404-409.
4. Arthur, M., D. Autissier, and P. Courvalin. 1986. Analysis of the nucleotide sequence of the *ereB* gene encoding the erythromycin esterase type II. *Nucleic Acids Res.* **14**:4987-4999.
5. Arthur, M., C. Molinas, C. Mabilat, and P. Courvalin. 1990. Detection of erythromycin resistance by the polymerase chain reaction using primers in conserved regions of *erm* rRNA methylase genes. *Antimicrob. Agents Chemother.* **34**:2024-2026.
6. Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl (ed.). 1987. *Current protocols in molecular biology*, vol. 1, p. 2.4.1-2.4.4. John Wiley & Sons, New York.
7. Avery, O. T., C. M. MacLeod, and M. McCarty. 1944. Studies on the chem-

- ical nature of the substance inducing transformation of pneumococcal types. *J. Exp. Med.* **79**:137–158.
8. **Azoulay-Dupuis, E., E. Vallee, B. Veber, and J. J. Pocidalo.** 1992. Efficacy of RP 59500, a new-semi-synthetic streptogramin, against penicillin- and multi-resistant strains of *Streptococcus pneumoniae* (sp), p. 329, abstr. 1311. In Program and Abstracts of the 32nd Interscience Conference on Antimicrobial Agents and Chemotherapy. American Society for Microbiology, Washington, D.C.
 9. **Barthélémy, P., D. Autissier, G. Gerbaud, and P. Courvalin.** 1984. Enzymatic hydrolysis of erythromycin by a strain of *Escherichia coli*: a new mechanism of resistance. *J. Antibiot.* **37**:1692–1696.
 10. **Bechhofer, D. H.** 1990. Triple post-transcriptional control. *Mol. Microbiol.* **4**:1419–1423.
 11. **Breiman, R. F., J. C. Butler, F. C. Tenover, J. A. Elliott, and R. R. Facklam.** 1994. Emergence of drug-resistant pneumococcal infections in the United States. *JAMA* **271**:1831–1835.
 12. **Brisson-Noël, A., P. Delrieu, D. Samain, and P. Courvalin.** 1988. Inactivation of lincosamide antibiotics in *Staphylococcus*. *J. Biol. Chem.* **263**: 15880–15887.
 13. **Celmer, W. D., W. P. Cullen, C. E. Moppett, J. B. Routien, R. Shibakawa, and J. Tome.** July 1977. U.S. patent 4,038,383.
 14. **Clewell, D. B., and A. E. Franke.** 1974. Characterization of a plasmid determining resistance to erythromycin, lincomycin, and vernamycin B_α in a strain of *Streptococcus pyogenes*. *Antimicrob. Agents Chemother.* **5**:534–537.
 15. **Coonan, K. M., and E. L. Kaplan.** 1994. In vitro susceptibility of recent North American Group A streptococcal isolates to eleven oral antibiotics. *Pediatr. Infect. Dis. J.* **13**:630–635.
 16. **Devriese, L. A.** 1980. Two new types of resistance to lincomycin in pathogenic staphylococci from animals. *Ann. Microbiol. (Paris)* **131**:261–266.
 17. **Djokic, S., G. Kobrehel, G. Lazarevski, N. Lopotar, Z. Tamburasev, G. Kamenar, A. Nagl, and I. Vickovic.** 1986. Erythromycin series. Part II. Ring expansion of erythromycin A oxime by the Beckmann rearrangement. *J. Chem. Soc. Perkin Trans. I* **1986**:1881–1890.
 18. **Dubnau, D.** 1984. Translational attenuation: the regulation of bacterial resistance to the macrolide-lincosamide-streptogramin B antibiotics. *Crit. Rev. Biochem.* **16**:103–132.
 19. **Eady, E. A., J. I. Ross, J. L. Tipper, C. E. Walters, J. H. Cove, and W. C. Noble.** 1993. Distribution of genes encoding erythromycin ribosomal methylases and an erythromycin efflux pump in epidemiologically distinct groups of staphylococci. *J. Antimicrob. Chemother.* **31**:211–217.
 20. **Feigin, R. D., R. E. Keeney, J. Nusrata, P. G. Shackelford, and R. D. Lins.** 1973. Efficacy of clindamycin therapy for otitis media. *Arch. Otolaryngol.* **98**:27–31.
 21. **Fernandez-Munoz, R., R. E. Monro, R. Torres-Pinedo, and D. Vasquez.** 1971. Substrate- and antibiotic-binding sites at the peptidyl-transferase centre of *Escherichia coli* ribosomes. Studies on the chloramphenicol, lincomycin and erythromycin sites. *Eur. J. Biochem.* **23**:185–193.
 22. **Fogel, S., and P. S. Sypher.** 1968. Extraction and isolation of individual ribosomal proteins from *Escherichia coli*. *J. Bacteriol.* **96**:358–364.
 23. **Friedland, I. R., S. Shelton, M. Paris, et al.** 1993. Dilemmas in diagnosis and management of cephalosporin-resistant *Streptococcus pneumoniae* meningitis. *Pediatr. Infect. Dis. J.* **12**:196–200.
 24. **Gerber, M. A.** 1995. Antibiotic resistance in Group A streptococci. *Pediatr. Clin. North Am.* **42**:539–551.
 25. **Girard, A. E., E. Girard, T. D. Gootz, J. A. Faiella, and C. R. Cimochofski.** 1995. In vivo efficacy of trovafloxacin (CP-99,219), a new quinolone with extended activities against gram-positive pathogens, *Streptococcus pneumoniae*, and *Bacteroides fragilis*. *Antimicrob. Agents Chemother.* **39**:2210–2216.
 26. **Hagman, K. E., W. P. Pan, B. G. Spratt, J. T. Balthazar, R. C. Judd, and W. M. Shafer.** 1995. Resistance of *Neisseria gonorrhoeae* to antimicrobial agents is modulated by the *mrRCDE* efflux system. *Microbiology* **141**:611–622.
 27. **Higgins, C. F., S. C. Hyde, M. M. Mimmack, U. Gileadi, D. R. Gill, and M. P. Gallagher.** 1990. Binding protein-dependent transport systems. *J. Bioenerg. Biomembr.* **22**:571–592.
 28. **Janosi, L., Y. Nakajima, and H. Hashimoto.** 1990. Characterization of plasmids that confer inducible resistance to 14-membered macrolides and streptogramin type B antibiotics in *Staphylococcus aureus*. *Microbiol. Immunol.* **34**:723–735.
 29. **Klugman, K. P., and H. J. Koornhof.** 1988. Drug resistance patterns and serogroups or serotypes of pneumococcal isolates from cerebrospinal fluid or blood, 1979–1986. *J. Infect. Dis.* **158**:956–964.
 30. **Klugman, K. P., H. J. Koornhof, and V. Kuhnle.** 1986. Clinical and nasopharyngeal isolates of unusual multiply resistant pneumococci. *Am. J. Dis. Child.* **140**:1186–1190.
 31. **Kono, M., K. O'Hara, and T. Ebisu.** 1992. Purification and characterization of macrolide 2'-phosphotransferase type II from a strain of *Escherichia coli* highly resistant to macrolide antibiotics. *FEMS Microbiol. Lett.* **97**:89–94.
 32. **Kreiswirth, B. M., S. Löfdahl, M. J. Betley, M. O'Reilly, P. M. Schlievert, M. S. Bergdoll, and R. P. Novick.** 1983. The toxic shock syndrome exotoxin structural gene is not detectably transmitted by a prophage. *Nature (London)* **305**:709–712.
 33. **Lai, C. J., and B. Weisblum.** 1971. Altered methylation of ribosomal RNA in an erythromycin-resistant strain of *Staphylococcus aureus*. *Proc. Natl. Acad. Sci. USA* **68**:856–860.
 34. **Lampson, B. C., W. von David, and J. T. Parisi.** 1986. Novel mechanism for plasmid-mediated erythromycin resistance by pNE24 from *Staphylococcus epidermidis*. *Antimicrob. Agents Chemother.* **30**:653–658.
 35. **Leclercq, R., A. Brisson-Noël, J. Duval, and P. Courvalin.** 1987. Phenotypic expression and genetic heterogeneity of lincosamide inactivation in *Staphylococcus* spp. *Antimicrob. Agents Chemother.* **31**:1887–1891.
 36. **Leclercq, R., and P. Courvalin.** 1991. Bacterial resistance to macrolide, lincosamide, and streptogramin antibiotics by target modification. *Antimicrob. Agents Chemother.* **35**:1267–1272.
 37. **Leclercq, R., and P. Courvalin.** 1991. Intrinsic and unusual resistance to macrolide, lincosamide, and streptogramin antibiotics in bacteria. *Antimicrob. Agents Chemother.* **35**:1273–1276.
 38. **Leclercq, R., and P. Courvalin.** 1993. Mechanisms of resistance to macrolides and functionally related antibiotics, p. 125–141. In A. J. Bryskier, J. P. Butzler, H. C. Neu, and P. M. Tulkens (ed.), *Macrolides—chemistry, pharmacology, and clinical uses*. Arnette, Blackwell, Paris.
 39. **Le Goffic, F., M. L. Capmau, J. Abbe, C. Cerceau, A. Dublanquet, and J. Duval.** 1977. Plasmid-mediated pristinamycin resistance: PH1A, a pristinamycin 1A hydrolase. *Ann. Inst. Pasteur (Paris)* **128**:471–474.
 40. **Le Goffic, F., M. L. Capmau, M. L. Bonnet, C. Cerceau, C. J. Soussy, A. Dublanquet, and J. Duval.** 1977. Plasmid-mediated pristinamycin resistance: PACIIA, a new enzyme which modifies pristinamycin IIA. *J. Antibiot.* **30**: 665–669.
 41. **Lewis, K.** 1994. Multidrug resistance pumps in bacteria: variations on a theme. *Trends Biochem. Sci.* **19**:119–123.
 42. **Matsuoka, M., K. Endou, S. Saitoh, M. Katoh, and Y. Nakajima.** 1995. A mechanism of resistance to partial macrolide and streptogramin B antibiotics in *Staphylococcus aureus* clinically isolated in Hungary. *Biol. Pharm. Bull.* **18**:1482–1486.
 43. **Matsuoka, M., L. Janosi, K. Endou, S. Saitoh, H. Hashimoto, and Y. Nakajima.** 1993. An increase of 63kDa-protein present in the cell membranes of *Staphylococcus aureus* that bears a plasmid mediating inducible resistance to partial macrolide and streptogramin B antibiotics. *Biol. Pharm. Bull.* **16**:1288–1290.
 44. **McCracken, G. H., Jr.** 1995. Emergence of resistant *Streptococcus pneumoniae*: a problem in pediatrics. *Pediatr. Infect. Dis. J.* **14**:424–428.
 45. **Meier, A., P. Kirschner, B. Springer, V. A. Steingrube, B. A. Brown, R. J. Wallace, Jr., and E. C. Böttger.** 1994. Identification of mutations in 23S rRNA gene of clarithromycin-resistant *Mycobacterium intracellulare*. *Antimicrob. Agents Chemother.* **38**:381–384.
 46. **Milton, E. D., C. L. Hewitt, and C. R. Harwood.** 1992. Cloning and sequencing of a plasmid-mediated erythromycin resistance determinant from *Staphylococcus xylosum*. *FEMS Microbiol. Lett.* **97**:141–147.
 47. **National Committee for Clinical Laboratory Standards.** 1995. Suggested groupings of U.S. FDA-approved antimicrobial agents that should be considered for routine testing and reporting on fastidious organisms by clinical microbiology laboratories, table 1A. In J. H. Jorgensen, W. A. Craig, G. V. Doern, M. J. Ferraro, S. M. Finegold, J. Jung-Tomc, S. L. Hansen, J. Hindler, D. A. Preston, L. B. Reller, J. M. Swenson, F. C. Tenover, M. A. Wikler, and W. R. Wilson (ed.), *Performance standards for antimicrobial susceptibility testing*, vol. 15, no. 14. National Committee for Clinical Laboratory Standards, Wayne, Pa.
 48. **Nelson, C. T., E. O. Mason, Jr., and S. L. Kaplan.** 1994. Activity of oral antibiotics in middle ear and sinus infections caused by penicillin-resistant *Streptococcus pneumoniae*: implications for treatment. *Pediatr. Infect. Dis. J.* **13**:585–589.
 49. **Nikaido, H.** 1994. Prevention of drug access to bacterial targets: permeability barriers and active efflux. *Science* **264**:382–388.
 50. **Nikaido, H., and D. G. Thanassi.** 1993. Penetration of lipophilic agents with multiple protonation sites into bacterial cells: tetracyclines and fluoroquinolones as examples. *Antimicrob. Agents Chemother.* **37**:1393–1399.
 51. **Noguchi, N., A. Emura, H. Matsuyama, K. O'Hara, M. Sasatsu, and M. Kono.** 1995. Nucleotide sequence and characterization of erythromycin resistance determinant that encodes macrolide 2'-phosphotransferase I in *Escherichia coli*. *Antimicrob. Agents Chemother.* **39**:2359–2363.
 52. **O'Hara, K.** 1994. Application of nuclear magnetic resonance spectrometry to measure the activity of bacterial macrolide esterase. *Microbios* **79**:231–239.
 53. **O'Hara, K., T. Kanda, K. Ohmiya, T. Ebisu, and M. Kono.** 1989. Purification and characterization of macrolide 2'-phosphotransferase from a strain of *Escherichia coli* that is highly resistant to erythromycin. *Antimicrob. Agents Chemother.* **33**:1354–1357.
 54. **Ounissi, H., and P. Courvalin.** 1985. Nucleotide sequence of the gene *ereA* encoding the erythromycin esterase in *Escherichia coli*. *Gene* **35**:271–278.
 55. **Pan, W., and B. G. Spratt.** 1994. Regulation of the permeability of the gonococcal envelope by the *mtr* system. *Mol. Microbiol.* **11**:769–775.
 56. **Paris, M. M., S. Shelton, M. Trujillo, S. M. Hickey, and G. H. McCracken, Jr.** 1996. Clindamycin therapy of experimental meningitis caused by penicil-

- lin- and cephalosporin-resistant *Streptococcus pneumoniae*. *Antimicrob. Agents Chemother.* **40**:122–126.
57. **Pernodet, J. L., F. Boccard, M.-T. Alegre, M. H. Blondelet-Rouault, and M. Guérineau.** 1988. Resistance to macrolides, lincosamides and streptogramin type B antibiotics due to a mutation in an rRNA operon of *Streptomyces ambofaciens*. *EMBO J.* **7**:277–282.
 58. **Phillips, G., D. Parratt, G. V. Orange, I. Harper, H. McEwan, and N. Young.** 1990. Erythromycin-resistant *Streptococcus pyogenes*. *J. Antimicrob. Chemother.* **25**:723–724.
 59. **Ross, J. I., E. A. Eady, J. H. Cove, and S. Baumberg.** 1995. Identification of a chromosomally encoded ABC-transport system with which the staphylococcal erythromycin exporter MsrA may interact. *Gene* **153**:93–98.
 60. **Ross, J. I., E. A. Eady, J. H. Cove, W. J. Cunliffe, S. Baumberg, and J. C. Wootton.** 1990. Inducible erythromycin resistance in staphylococci is encoded by a member of the ATP-binding transport super-gene family. *Mol. Microbiol.* **4**:1207–1214.
 61. **Ross, J. I., A. M. Farrell, E. A. Eady, J. H. Cove, and W. J. Cunliffe.** 1989. Characterization and molecular cloning of the novel macrolide-streptogramin B resistance determinant from *Staphylococcus epidermidis*. *J. Antimicrob. Chemother.* **24**:851–862.
 62. **Scott, R. J. D., J. Naidoo, N. F. Lightfoot, and R. C. George.** 1989. A community outbreak of group A beta haemolytic streptococci with transferable resistance to erythromycin. *Epidemiol. Infect.* **102**:95–91.
 63. **Seppälä, H., A. Nissinen, H. Järvinen, S. Huovinen, T. Henriksson, E. Herva, S. E. Holm, M. Jahkola, M.-L. Katila, T. Klaukka, S. Kontiainen, O. Liimatainen, S. Oinonen, L. Passi-Metsomaa, and P. Huovinen.** 1992. Resistance to erythromycin in Group A streptococci. *N. Engl. J. Med.* **326**:292–297.
 64. **Seppälä, H., A. Nissinen, Q. Yu, and P. Huovinen.** 1993. Three different phenotypes of erythromycin-resistant *Streptococcus pyogenes* in Finland. *J. Antimicrob. Chemother.* **32**:885–891.
 65. **Shandler, L., J. Gonzales, and G. D. Overturf.** 1996. Recurrence of pneumococcal sepsis caused by an intermediately penicillin-resistant organism treated with loracarbef. *Pediatr. Infect. Dis. J.* **15**:379–380.
 66. **Spika, J. S., R. R. Facklam, B. D. Plikaytis, and M. J. Oxtoby.** 1991. Antimicrobial resistance of *Streptococcus pneumoniae* in the United States, 1979–1987. The pneumococcal surveillance working group. *J. Infect. Dis.* **163**:1273–1278.
 67. **Stingemore, N., G. R. J. Francis, M. Toohey, and D. B. McGeachie.** 1989. The emergence of erythromycin resistance in *Streptococcus pyogenes* in Fremantle, Western Australia. *Med. J. Aust.* **150**:626–631.
 68. **Sutcliffe, J., T. Grebe, A. Tait-Kamradt, and L. Wondrack.** Detection of erythromycin-resistant determinants using the polymerase chain reaction. Submitted for publication.
 69. **Swendsen, C. L., and W. Johnson.** 1976. Humoral immunity to *Streptococcus pneumoniae* induced by a pneumococcal ribosomal protein fraction. *Infect. Immun.* **14**:345–354.
 70. **Thornsberry, C., S. D. Brown, Y. C. Yee, S. K. Bouchillon, J. K. Marler, and T. Rich.** 1993. Increasing penicillin resistance in *Streptococcus pneumoniae* in the U.S. Effect on susceptibility to oral cephalosporins. *Infections Med.* **10**:15–24.
 71. **Weisblum, B.** 1985. Inducible resistance to macrolides, lincosamides, and streptogramin type B antibiotics: the resistance phenotype, its biological diversity, and structural elements that regulate expression—a review. *J. Antimicrob. Chemother.* **16**(Suppl. A):63–90.
 72. **Weisblum, B.** 1995. Erythromycin resistance by ribosome modification. *Antimicrob. Agents Chemother.* **39**:577–585.
 73. **Weisblum, B.** 1995. Insights into erythromycin action from studies of its activity as inducer of resistance. *Antimicrob. Agents Chemother.* **39**:797–805.
 74. **Welby, P. E., D. S. Keller, J. L. Cromien, P. Tebas, and G. A. Storch.** 1994. Resistance to penicillin and non- β -lactam antibiotics of *Streptococcus pneumoniae* at a children's hospital. *Pediatr. Infect. Dis. J.* **13**:281–287.
 75. **Wondrack, L., M. Massa, B. V. Yang, and J. Sutcliffe.** 1996. A clinical strain of *Staphylococcus aureus* inactivates and effluxes macrolides. *Antimicrob. Agents Chemother.* **40**:992–998.
 76. **Wondrack, L., and J. Sutcliffe.** 1995. *Streptococcus* strains with a M phenotype efflux erythromycin, a 14-membered macrolide, p. 158, abstr. A-85. In Abstracts of the 95th General Meeting of the American Society for Microbiology, Washington, D.C.
 77. **Youngs, E. R.** 1984. Erythromycin resistant *Streptococcus pyogenes* in Merseyside. *J. Infect.* **8**:86–87.