

Mechanism of Fluconazole Resistance in *Candida krusei*

ALISON S. OROZCO,¹ LINDSEY M. HIGGINBOTHAM,¹ CHRISTOPHER A. HITCHCOCK,²
TANYA PARKINSON,² DEREK FALCONER,² ASHRAF S. IBRAHIM,¹
MAHMOUD A. GHANNOUM,^{1,3†} AND SCOTT G. FILLER^{1,3*}

St. John's Cardiovascular Research Center, Division of Infectious Diseases, Harbor-UCLA Research and Education Institute, Torrance, California 90502¹; Pfizer Central Research, Sandwich, Kent, United Kingdom²; and the UCLA School of Medicine, Los Angeles, California 90024³

Received 2 April 1998/Returned for modification 30 May 1998/Accepted 9 July 1998

The mechanisms of fluconazole resistance in three clinical isolates of *Candida krusei* were investigated. Analysis of sterols of organisms grown in the absence and presence of fluconazole demonstrated that the predominant sterol of *C. krusei* is ergosterol and that fluconazole inhibits 14 α -demethylase in this organism. The 14 α -demethylase activity in cell extracts of *C. krusei* was 16- to 46-fold more resistant to inhibition by fluconazole than was 14 α -demethylase activity in cell extracts of two fluconazole-susceptible strains of *Candida albicans*. Comparing the carbon monoxide difference spectra of microsomes from *C. krusei* with those of microsomes from *C. albicans* indicated that the total cytochrome P-450 content of *C. krusei* is similar to that of *C. albicans*. The Soret absorption maximum in these spectra was located at 448 nm for *C. krusei* and at 450 nm for *C. albicans*. Finally, the fluconazole accumulation of two of the *C. krusei* isolates was similar to if not greater than that of *C. albicans*. Thus, there are significant qualitative differences between the 14 α -demethylase of *C. albicans* and *C. krusei*. In addition, fluconazole resistance in these strains of *C. krusei* appears to be mediated predominantly by a reduced susceptibility of 14 α -demethylase to inhibition by this drug.

As the use of azole antifungal agents has risen dramatically, there has been an increase in the number of reports of azole-resistant isolates of *Candida*, especially in patients with advanced AIDS. In addition to the development of resistance by *Candida albicans* and *Candida glabrata*, it has been reported that the frequent use of fluconazole can select for the emergence of *Candida krusei* as a commonly isolated opportunistic pathogen in some medical centers (1, 33). This finding is clinically significant because *C. krusei* can cause serious infections in susceptible patients (8, 19). Furthermore, this organism is usually intrinsically resistant to fluconazole, both in vitro (3) and in vivo (4).

Three general mechanisms of azole resistance have been described for species of *Candida*. The first is an alteration in the target enzyme, 14 α -demethylase. Inhibition of this enzyme by azoles causes an accumulation of C₁₄ methylated sterols which likely disrupt membrane structure (9). In some resistant organisms, there is overexpression of the 14 α -demethylase gene and/or the enzyme is less susceptible to azole inhibition (15, 24, 32). The second mechanism is decreased drug accumulation, mediated by either diminished uptake or increased efflux of the drug (22, 26). The third mechanism of resistance is the presence of a deficiency in C₅₍₆₎ desaturase. Organisms deficient in this enzyme produce 14-methylfecosterol and remain viable when 14 α -demethylase activity is inhibited (5, 14).

To determine if fluconazole resistance in *C. krusei* is mediated by one or more of these mechanisms, we analyzed the effects of this antifungal agent on sterol synthesis by three strains of *C. krusei*. In addition, the fluconazole uptake and cytochrome P-450 content of these organisms were measured. Our results indicate that the predominant mechanism of flu-

conazole resistance in these organisms is a 14 α -demethylase with reduced susceptibility to the inhibitory effects of fluconazole.

MATERIALS AND METHODS

Fungal strains and susceptibility testing. Three clinical isolates of *C. krusei*, strains 91-1158, 91-1159, and 91-1161, were generously provided by Michael Rinaldi (San Antonio, Tex.). A fourth isolate of *C. krusei*, ATCC 6258, was obtained from the American Type Culture Collection (Rockville, Md.). Two clinical isolates of *C. albicans*, Y01.345 and SC5314, were supplied by Christopher Hitchcock and William Fonzi (Georgetown University School of Medicine, Washington, D.C.), respectively. The susceptibilities of the organisms to fluconazole and itraconazole were determined at 24 and 48 h by the National Committee for Clinical Laboratory Standards M27-A broth microdilution method at an inoculum of 10³ organisms per well (20). The medium was RPMI 1640 (Irvine Scientific, Santa Ana, Calif.) buffered to pH 7.0 with 0.165 M morpholinepropanesulfonic acid (MOPS). The MICs were defined as the concentrations of drug that reduced growth by 80% compared to that of organisms grown in the absence of drug.

Sterol analysis. For sterol extraction, each strain of *C. krusei* was grown for 24 h at 37°C on a rotary shaker. The medium was Sabouraud dextrose broth (Difco, Detroit, Mich.), with and without fluconazole. The concentration of fluconazole was 16 μ g/ml, the highest concentration at which these organisms would grow in this medium. The organisms were harvested by centrifugation and washed twice in 0.85% saline, and their total sterols were extracted by ethanolic KOH, as described previously (6, 7). The resultant sterols were further purified by thin-layer chromatography on PK6F silica gel 60-Å plates (Whatman, Clifton, N.J.) with a solvent system of petroleum ether-diethyl ether (3:1, vol/vol) (27). The sterols were eluted from the silica in chloroform-diethyl ether-ethanol (1:1:1). After being derivatized with hexamethyldisilazane and trimethylchlorosilane (28), the sterols were redissolved in hexane and analyzed by gas chromatography-mass spectrometry. The sterols were identified by comparison to known standards and published data (16, 17, 23).

Fluconazole accumulation. A filter-based assay was used to measure the accumulation of fluconazole by the organisms (22). These organisms were grown to exponential phase in Sabouraud dextrose broth and suspended in phosphate-buffered saline (pH 7.5) containing 5% glucose (wt/vol) at 10⁸ organisms per ml. Next, a mixture of [³H]fluconazole (specific activity, 715 GBq/mmol) and unlabeled fluconazole was added to the cells so that the final concentration of fluconazole was 100 nM (0.2 μ Ci/ml). At selected intervals, aliquots were removed and the organisms were collected by filtration. Next, the organisms were washed four times in phosphate-buffered saline containing 100 μ M unlabeled fluconazole. The amount of cell-associated radioactivity was determined by scintillation counting. All experiments were performed in triplicate.

Carbon monoxide difference spectra of microsomes. The cytochrome P-450 content of the organisms was analyzed by measuring their carbon monoxide difference spectra. Organisms were grown to exponential phase in Sabouraud

* Corresponding author. Mailing address: Division of Infectious Diseases, Harbor-UCLA Research and Education Institute, 1124 West Carson St., RB-2, Torrance, CA 90502. Phone: (310) 222-6426. Fax: (310) 782-2016. E-mail: Filler@HUMC.EDU.

† Present address: Department of Dermatology, Center for Medical Mycology, Case Western Reserve University, Cleveland, OH 44106.

TABLE 1. Antifungal susceptibilities of *C. albicans* and *C. krusei*

Organism	MIC ($\mu\text{g/ml}$)			
	Fluconazole		Itraconazole	
	24 h	48 h	24 h	48 h
<i>C. albicans</i> Y01.345	0.25	<0.25	0.125	0.125
<i>C. albicans</i> SC5314	0.5	1	0.5	0.5
<i>C. krusei</i> 91-1158	32	64	0.5	0.5
<i>C. krusei</i> 91-1159	16	32	0.5	0.5
<i>C. krusei</i> 91-1161	16	32	0.5	0.5

dextrose broth at 37°C and harvested by centrifugation. They were spheroplasted with lyticase (Sigma, St. Louis, Mo.) in sorbitol buffer (1.5 M sorbitol in 10 mM Tris buffer, pH 7.4) and resuspended in 10 mM Tris buffer, pH 7.4, containing 0.65 M sorbitol, 0.1 mM EDTA, 0.1 mM glutathione, and protease inhibitors (Boehringer Mannheim, Indianapolis, Ind.) (10). All subsequent steps were carried out at 4°C. The spheroplasts were broken into microsomes by sonication, after which debris and unbroken cells were removed by centrifugation at $1,500 \times g$ for 10 min followed by $25,000 \times g$ for 25 min. The microsomes were harvested with calcium chloride by the method of Käppeli et al. (13), washed once in 10 mM Tris buffer, pH 7.4, containing 150 mM potassium chloride, and resuspended in 100 mM Tris containing 0.65 M glycerol, 0.1 mM EDTA, and 0.1 mM glutathione. The carbon monoxide difference spectra of the microsomes were measured, and their cytochrome P-450 content was calculated by using an extinction coefficient of 91 liters/mmol/cm (21). The protein concentration of each microsome suspension was determined by the Bio-Rad protein assay (Bio-Rad, Hercules, Calif.). For each organism, the cytochrome P-450 contents of at least three different preparations of microsomes were measured.

Sterol biosynthesis by cell extracts. To determine the effects of fluconazole on the synthesis of ergosterol from [^{14}C]mevalonic acid, the organisms were grown in Sabouraud dextrose broth until late exponential phase. The organisms were broken by vortexing with glass beads in 0.1 M potassium phosphate, pH 7.5. Debris and unbroken cells were removed by centrifugation at $2,000 \times g$ for 5 min followed by $10,000 \times g$ for 10 min. To measure sterol biosynthesis, 925 μl of the resultant supernatant was added to 75 μl of cofactor buffer to achieve the following final concentrations: 1 μM NADP, 1 μM NADPH, 1 μM NAD, 7 μM glucose-6-phosphate, 5 μM ATP, 3 μM reduced glutathione, 2 μM MnCl_2 , and 0.25 μCi [^{14}C]mevalonic acid (10, 12). Selected concentrations of fluconazole were added to the reaction mixtures before the addition of the cell extracts. After incubation at 37°C for 2 h, the reaction was stopped with ethanolic KOH. The samples were saponified at 80°C for 45 min, after which the nonsaponifiable lipids were extracted with petroleum ether (bp 40 to 60°C). The samples were dried under nitrogen and redissolved in chloroform. The sterols were separated by thin-layer chromatography on silica gel LK6D (Whatman) with a solvent system of petroleum ether-diethyl ether (3:1, vol/vol) (27). The sterols were visualized by iodine staining and identified by comparison with commercially available standards which were run in parallel. Next, the sterol-containing bands were scraped from the plates and ^{14}C incorporation was determined by liquid scintillation counting. All experiments were repeated at least three times with different cell extracts.

TABLE 2. Effect of fluconazole on the sterols of *C. krusei*

Sterol	Proportion of total sterols (%)							
	Control (no drug)				Fluconazole ^a			
	91-1158	91-1159	91-1161	ATCC 6258	91-1158	91-1159	91-1161	ATCC 6258
Cholesta-7,24-dienol	1.3	1.8	1.3	1.8	0.0	0.0	0.0	0.0
Ergosterol	88.9	86.5	91.1	93.0	82.1	83.4	83.7	72.9
Ergosta-7,22-dienol	0.0	0.0	0.0	0.0	0.0	1.0	0.0	0.0
14-Methylfecosterol	0.0	0.0	0.0	0.0	2.1	1.2	1.8	1.5
Ergosta-8,24(28)-dienol	0.0	3.2	2.3	0.0	0.6	0.0	0.0	0.0
Fecosterol	2.9	0.0	0.0	1.5	0.0	0.0	0.0	0.0
Ergosta-5,7-dienol	2.6	3.7	2.3	1.4	0.5	1.5	0.6	1.1
Episterol	1.6	2.4	1.8	1.1	0.5	0.0	0.2	0.3
Lanosterol	0.4	0.4	0.3	0.5	5.9	9.0	10.9	19.5
4-Methylfecosterol	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3
4,14-Dimethyl-cholesta-8,24-dienol	2.3	2.0	1.0	0.6	5.9	0.0	0.0	0.0
Eburicol	0.0	0.0	0.0	0.0	0.4	1.8	0.5	2.0
Unidentified sterols	0.0	0.0	0.0	0.0	2.0	2.0	2.2	2.5

^a Organisms were grown with fluconazole (16 $\mu\text{g/ml}$) for 24 h.

RESULTS

Growth in fluconazole reduced the ergosterol content of *C. krusei*. Fluconazole MICs for *C. krusei* 91-1158, 91-1159, and 91-1161 were all high, whereas both strains of *C. albicans* were susceptible to this drug (Table 1). *C. albicans* Y01.345 was slightly more susceptible to itraconazole than *C. albicans* SC5314 or the three strains of *C. krusei*. Analysis of the sterols of the organisms by gas chromatography-mass spectrometry revealed that the predominant sterol in *C. krusei* was ergosterol (Table 2). Growing all three strains of *C. krusei* in the presence of fluconazole caused a decrease in their ergosterol content and a marked increase in lanosterol. Other 14-methyl sterols that were increased in organisms exposed to fluconazole were 14-methylfecosterol and eburicol. These results suggest that fluconazole inhibits 14 α -demethylase in *C. krusei*. No 14-methyl-ergosta-8,24(28)-dien-3,6-diol was detected, although this sterol has been reported to accumulate when *C. krusei* is exposed to azoles (18, 31).

The 14 α -demethylase from cell extracts of *C. krusei* showed reduced susceptibility to inhibition by fluconazole. Although fluconazole appeared to inhibit 14 α -demethylase in *C. krusei*, exposure to a relatively high concentration of drug was required to produce this inhibition. Therefore, we examined whether the enzyme itself was resistant to the inhibitory effects of fluconazole. We determined the effects of this drug on the synthesis of ergosterol from [^{14}C]mevalonic acid in cell extracts of three strains of *C. krusei* and two isolates of *C. albicans*. We found that the concentration of fluconazole required to inhibit the synthesis of ergosterol by 50% (IC₅₀) was 16- to 46-fold higher in cell extracts from *C. krusei* than in extracts from either strain of *C. albicans* (Table 3). This difference indicates that reduced fluconazole susceptibility of the 14 α -demethylase from *C. krusei* is likely the predominant mechanism of fluconazole resistance in these isolates.

Carbon monoxide difference spectra also indicated that the cytochrome P-450 of *C. krusei* is different from that of *C. albicans*. Next, we determined the carbon monoxide difference spectra of microsomes from these organisms. This procedure enabled us to evaluate further the possibility that fluconazole resistance was mediated by quantitative and/or qualitative changes in the 14 α -demethylase of the organism. We found that the cytochrome P-450 content of the three strains of *C. krusei* was similar to that of the two strains of *C. albicans* (Table 3). Since 14 α -demethylase accounts for the majority of

TABLE 3. Comparison of total cytochrome P-450 content and fluconazole susceptibility of 14 α -demethylases of *C. albicans* and *C. krusei*^a

Organism	Fluconazole IC ₅₀ (μ M) ^b	Cytochrome P-450 content (nmol/mg of protein)
<i>C. albicans</i> Y01.345	0.017 \pm 0.005	0.077 \pm 0.037
<i>C. albicans</i> SC5314	0.032 \pm 0.006	0.058 \pm 0.037
<i>C. krusei</i> 91-1158	0.751 \pm 0.384	0.052 \pm 0.021
<i>C. krusei</i> 91-1159	0.495 \pm 0.208	0.065 \pm 0.018
<i>C. krusei</i> 91-1161	0.795 \pm 0.436	0.077 \pm 0.060

^a Results are the means \pm standard deviations of at least three separate measurements.

^b Concentration of fluconazole required to inhibit the synthesis of ergosterol from [¹⁴C]mevalonic acid by 50%.

cytochrome P-450 in yeasts, these findings suggest that the mechanism of fluconazole resistance in *C. krusei* is not mediated by overproduction of the target enzyme.

In these experiments, the Soret absorption maximum of microsomes from all three strains of *C. krusei* was located at 448 nm whereas the Soret peak from microsomes of *C. albicans* Y01.345 and SC5314 was located at 450 nm. This difference in absorption maxima also suggests that there are qualitative differences in the 14 α -demethylase from the two different species of *Candida*.

Fluconazole accumulation by *C. krusei* and *C. albicans* was similar. To determine if reduced intracellular concentrations of the drug contributed to fluconazole resistance, the fluconazole accumulation of the three isolates of *C. krusei* was compared with that of the two strains of *C. albicans*. During the first 30 min, the accumulation of fluconazole by *C. krusei* was not consistently different from the accumulation of this drug by *C. albicans* (Fig. 1). However, at later time points, there was a trend towards greater accumulation of fluconazole by the strains of *C. krusei*. Of the organisms studied, *C. albicans* Y01.345 had the lowest fluconazole accumulation at 60 and 90 min. Therefore, the three isolates of *C. krusei* appear to be resistant to fluconazole by a mechanism that is independent of reduced fluconazole accumulation.

DISCUSSION

By comparing the effects of fluconazole on the synthesis of sterols by cell extracts of *C. krusei* and *C. albicans*, we deter-




FIG. 1. Accumulation of fluconazole by *C. krusei* and *C. albicans*. Results are the means of triplicate determinations. Error bars indicate standard deviations.

mined that a major mechanism of fluconazole resistance in the three strains of *C. krusei* studied appears to be a reduced susceptibility of 14 α -demethylase to fluconazole. The fluconazole IC₅₀s for sterol synthesis by cell extracts of *C. krusei* ranged from 0.495 to 0.795 μ M. These values were 16- to 46-fold greater than the corresponding IC₅₀s for extracts of *C. albicans*. In the literature, there is substantial variability in the reported fluconazole IC₅₀s for cell extracts of *C. krusei*. These values range from 0.080 to 1.38 μ M (18, 31). Possible explanations for these differences include differences in methodology or strain-to-strain variations in the mechanism of fluconazole resistance in *C. krusei*. It is noteworthy that no previous investigators have reported the direct comparison of the fluconazole IC₅₀ of *C. krusei* with that of a fluconazole-susceptible species of *Candida*. However, Venkateswarlu et al. (30) have reported that the fluconazole IC₅₀ for cell extracts of different strains of *C. albicans* ranged from 0.042 to 0.055 μ M, and we have determined previously that the fluconazole IC₅₀ for 14 α -demethylase purified from *C. albicans* is 0.074 μ M (11). These values are only slightly higher than the ones reported here.

An additional finding was that the Soret maximum of microsomes of *C. krusei* was located at 448 nm, whereas it was located at 450 nm when microsomes of *C. albicans* were analyzed. This result also suggests that the 14 α -demethylase of *C. krusei* is significantly different from that of *C. albicans*. Venkateswarlu et al. (29, 31) reported that the Soret peak of *C. krusei* microsomes was at 448 nm, which is similar to our findings. However, other investigators have found that the Soret peak of *C. albicans* microsomes is also located 448 nm, and we have determined that purified 14 α -demethylase of *C. albicans* has a Soret peak at 447 (12, 15). It is possible that the difference between these results and our present findings is the result of differences in culture conditions and/or the methods used to purify the microsomes. For example, Sanglard et al. (25) found that the Soret maximum of microsomes of *Candida tropicalis* ranged from 447 to 450 nm, depending on the conditions under which the organisms were grown.

Burgener-Kairuz et al. (2) have determined the sequence of a 1.2-kb fragment of the *C. krusei* 14 α -demethylase gene. They found that the deduced amino acid sequence of this fragment was only 80% similar to the corresponding portion of the 14 α -demethylase of *C. albicans*. This finding also supports our conclusion that the 14 α -demethylase of *C. krusei* differs significantly from that of *C. albicans*. It is not known which region(s) of the *C. krusei* 14 α -demethylase is responsible for its reduced susceptibility to fluconazole, but this question is currently being investigated.

The reduced carbon monoxide difference spectra also indicated that the cytochrome P-450 content of *C. krusei* was similar to that of *C. albicans*. Other investigators have reported that the cytochrome P-450 contents of *C. krusei* and *C. albicans* are approximately 0.03 nmol/mg of protein and 0.02 to 0.05 nmol/mg of protein, respectively (12, 29-31). These results support the conclusion that *C. krusei* is not resistant to fluconazole because of overproduction of the target enzyme.

When the different strains of *C. krusei* were grown in the presence of fluconazole, we found a decrease in the amount of ergosterol and an increase in 14-methyl sterols, mainly lanosterol. Unlike other investigators, we did not find any evidence of 14-methyl-ergosta-8,24(28)-dien-3,6-diol in the organisms exposed to fluconazole (18, 31). This sterol was not detected in cell extracts from *C. krusei* ATCC 6258, even though Venkateswarlu et al. (31) reported its presence when this strain was grown in the presence of itraconazole. A likely explanation for this difference is that even though we grew the organisms in the maximal concentration of fluconazole that did not completely

inhibit fungal growth, there was only partial inhibition of ergosterol synthesis. Others have observed that a strain of *C. krusei* grown in subinhibitory concentrations of itraconazole had a reduction in ergosterol content without the accumulation of 14-methyl-ergosta-8,24(28)-dien-3,6-diol (18). Thus, it is likely that, if it had been possible to grow the organisms in higher concentrations of fluconazole, we would have found the accumulation of this sterol. Nevertheless, because we found that fluconazole caused an accumulation of 14-methylfecosterol which was not converted to 14-methyl-ergosta-8,24(28)-dien-3,6-diol, we cannot completely rule out the possibility that C₅₍₆₎ desaturase contributes to fluconazole resistance in some strains of *C. krusei*.

An additional finding was that the fluconazole accumulations of the three strains of *C. krusei* were similar to those of *C. albicans*. These results are different from those of Marichal et al. (18), who concluded that *C. krusei* is resistant to fluconazole on the basis of diminished drug accumulation. However, these investigators found that the fluconazole accumulation by exponential-phase organisms was actually fourfold greater than that observed with the strains of *C. krusei* used in this study. It is possible that differences in either methodology or the strains of *C. krusei* account for these differences in results.

Venkateswarlu et al. (31) also reported that itraconazole resistance in *C. krusei* is mediated by reduced drug accumulation. They found that a strain of *C. krusei* that was resistant to itraconazole accumulated less drug than did an itraconazole-susceptible isolate of *C. krusei*. While the two organisms had different susceptibilities to itraconazole, both exhibited high-level fluconazole resistance. These results demonstrate that resistance to fluconazole in *C. krusei* is mediated by a different mechanism than is itraconazole resistance.

In conclusion, our data indicate that, in the strains of *C. krusei* studied, fluconazole resistance is largely the result of a decreased susceptibility of 14 α -demethylase to the inhibitory effects of fluconazole. Diminished accumulation of fluconazole did not appear to contribute to fluconazole resistance in these isolates. However, it is possible that reduced fluconazole accumulation is the predominant mechanism of fluconazole resistance in other isolates of *C. krusei*. Future work will investigate whether different strains of *C. krusei* have different mechanisms of resistance to this drug and why the 14 α -demethylase of this organism is less susceptible to inhibition by fluconazole.

ACKNOWLEDGMENTS

We thank Michael Mador and Trang Phan for technical assistance. We also appreciate the assistance of W.-N. Paul Lee and Anne Berger at the Stable Isotope Facility at Harbor-UCLA Research and Education Institute.

This work was supported by a grant from Pfizer, Inc.

REFERENCES

- Borg-von Zepelin, M., H. Eiffert, M. Kann, and R. Ruchel. 1993. Changes in the spectrum of fungal isolates: results from clinical specimens gathered in 1987/88 compared with those in 1991/92 in the University Hospital Göttingen, Germany. *Mycoses* **36**:247–253.
- Burgener-Kairuz, P., J.-P. Zuber, P. Jaunin, T. G. Buchman, J. Bille, and M. Rossier. 1994. Rapid detection and identification of *Candida albicans* and *Torulopsis (Candida) glabrata* in clinical specimens by species-specific nested PCR amplification of a cytochrome P-450 lanosterol- α -demethylase (LIA1) gene fragment. *J. Clin. Microbiol.* **32**:1902–1907.
- Dermoumi, H. 1992. In vitro susceptibility of yeast isolates from the blood to fluconazole and amphotericin B. *Chemotherapy* **38**:112–117.
- Fisher, M. A., S.-H. Shen, J. Haddad, and W. F. Tarry. 1989. Comparison of in vivo activity of fluconazole with that of amphotericin B against *Candida tropicalis*, *Candida glabrata*, and *Candida krusei*. *Antimicrob. Agents Chemother.* **33**:1443–1446.
- Geber, A., C. A. Hitchcock, J. E. Swartz, F. S. Pullen, K. E. Marsden, K. J. Kwon-Chung, and J. E. Bennett. 1995. Deletion of the *Candida glabrata* ERG3 and ERG11 genes: effect on cell viability, cell growth, sterol composition, and antifungal susceptibility. *Antimicrob. Agents Chemother.* **39**:2708–2717.
- Ghannoum, M. A., G. Janini, L. Khamis, and S. S. Radwan. 1986. Dimorphism-associated variations in the lipid composition of *Candida albicans*. *J. Gen. Microbiol.* **132**:2367–2375.
- Ghannoum, M. A., I. Swairjo, and D. R. Soll. 1990. Variation in lipid and sterol contents in *Candida albicans* white and opaque phenotypes. *J. Med. Vet. Mycol.* **28**:103–115.
- Goldman, M., J. C. Pottage, and D. C. Weaver. 1993. *Candida krusei* fungemia. Report of 4 cases and review of the literature. *Medicine* **72**:143–150.
- Hitchcock, C. A. 1993. Resistance of *Candida albicans* to azole antifungal agents. *Biochem. Soc. Trans.* **21**:1039–1047.
- Hitchcock, C. A., S. B. Brown, E. G. V. Evans, and D. J. Adams. 1989. Cytochrome P-450-dependent 14 α -demethylation of lanosterol in *Candida albicans*. *Biochem. J.* **260**:549–556.
- Hitchcock, C. A., K. Dickinson, S. B. Brown, E. G. V. Evans, and D. J. Adams. 1990. Interaction of azole antifungal antibiotics with cytochrome P-450-dependent 14 α -sterol demethylase purified from *Candida albicans*. *Biochem. J.* **266**:475–480.
- Hitchcock, C. A., K. Dickinson, S. B. Brown, E. G. V. Evans, and D. J. Adams. 1989. Purification and properties of cytochrome P-450-dependent 14 α -sterol demethylase from *Candida albicans*. *Biochem. J.* **263**:573–579.
- Käppeli, O., M. Sauer, and A. Fiechter. 1982. Convenient procedure for the isolation of highly enriched, cytochrome P-450-containing microsomal fraction from *Candida tropicalis*. *Anal. Biochem.* **126**:179–182.
- Kelly, S. L., D. C. Lamb, D. E. Kelly, N. J. Manning, J. Loeffler, H. Hebart, U. Schumacher, H. Einsele. 1997. Resistance to fluconazole and cross-resistance to amphotericin B in *Candida albicans* from AIDS patients caused by defective sterol $\Delta^{5,6}$ -desaturation. *FEBS Lett.* **400**:80–82.
- Lamb, D. C., D. E. Kelly, W. H. Schunck, A. Z. Shyadehi, M. Akhtar, D. J. Lowe, B. C. Baldwin, and S. L. Kelly. 1997. The mutation T315A in *Candida albicans* sterol 14 α -demethylase causes reduced enzyme activity and fluconazole resistance through reduced affinity. *J. Biol. Chem.* **272**:5682–5688.
- Loeffler, R. S. T., and A. L. Hayes. 1990. Sterols of the plant pathogenic fungi *Botrytis cinera* and *pyrenophora teres*. *Phytochemistry* **29**:3424–3425.
- Loeffler, R. S. T., and A. L. Hayes. 1992. Effects of sterols biosynthesis inhibitor fungicides on growth and sterol composition of *Ustilago maydis*, *Botrytis cinera* and *pyrenophora teres*. *Pestic. Sci.* **35**:7–17.
- Marichal, P., J. Gorrens, M.-C. Coene, L. LeJeune, and H. Vanden Bossche. 1995. Origin in differences in susceptibility of *Candida krusei* to azole antifungal agents. *Mycoses* **38**:111–117.
- McQuillen, D. P., B. S. Zingman, F. Meunier, and S. M. Levitz. Invasive infections due to *Candida krusei*: report of ten cases of fungemia that include three cases of endophthalmitis. *Clin. Infect. Dis.* **14**:472–478.
- National Committee for Clinical Laboratory Standards. 1995. Reference method for broth dilution antifungal susceptibility testing of yeast. Tentative standard. Document M27-T. National Committee for Clinical Laboratory Standards, Villanova, Pa.
- Omura, T., and R. Sato. 1964. The carbon monoxide-binding pigment of liver microsomes. I. Evidence for its hemoprotein nature. *J. Biol. Chem.* **239**:2370–2378.
- Parkinson, T., D. J. Falconer, and C. A. Hitchcock. 1995. Fluconazole resistance due to energy-dependent drug efflux in *Candida glabrata*. *Antimicrob. Agents Chemother.* **39**:1696–1699.
- Quail, M. A., A. Arnoldi, D. J. Moore, M. W. Goosey, and S. L. Kelly. 1993. Ketoconazole mediated growth inhibition in *Botrytis cinera* and *Saccharomyces cerevisiae*. *Phytochemistry* **32**:273–280.
- Sanglard, D., F. Ischer, L. Koymans, and J. Bille. 1998. Amino acid substitutions in the cytochrome P-450 lanosterol 14 α -demethylase (CYP51A1) from azole-resistant *Candida albicans* clinical isolates contribute to resistance to azole antifungal agents. *Antimicrob. Agents Chemother.* **42**:241–253.
- Sanglard, D., O. Käppeli, and A. Fiechter. 1986. The distinction of different types of cytochromes P-450 from the yeasts *Candida tropicalis* and *Saccharomyces uvarum*. *Arch. Biochem. Biophys.* **251**:276–286.
- Sanglard, D., K. Kuchler, F. Ischer, J.-L. Pagani, M. Monod, and J. Bille. 1995. Mechanisms of resistance to azole antifungal agents in *Candida albicans* isolates from AIDS patients involve specific multidrug transporters. *Antimicrob. Agents Chemother.* **39**:2378–2386.
- Sorkhoh, N. A., M. A. Ghannoum, A. S. Ibrahim, R. J. Stretton, and S. S. Radwan. 1991. Growth of *Candida albicans* in the presence of hydrocarbons: a correlation between sterol concentration and hydrocarbon uptake. *Appl. Microbiol. Biotechnol.* **34**:509–512.
- Vandenheuvel, F. A., and A. S. Court. 1968. Reference high-efficiency non-polar packed column for the gas-liquid chromatography of nanogram amounts of sterols. Part I. Retention time data. *J. Chromatogr.* **38**:439–459.
- Venkateswarlu, K., D. W. Denning, and S. L. Kelly. 1997. Inhibition and interaction of cytochrome P450 of *Candida krusei* with azole antifungal drugs. *J. Med. Vet. Mycol.* **35**:19–25.
- Venkateswarlu, K., D. W. Denning, N. J. Manning, and S. L. Kelly. 1996.

- Comparison of D0870, a new triazole antifungal agent, to fluconazole for inhibition of *Candida albicans* cytochrome P-450 by using in vitro assays. *Antimicrob. Agents Chemother.* **40**:1382–1386.
31. **Venkateswarlu, K., D. W. Denning, N. J. Manning, and S. L. Kelly.** 1996. Reduced accumulation of drug in *Candida krusei* accounts for itraconazole resistance. *Antimicrob. Agents Chemother.* **40**:2443–2446.
32. **White, T. C.** 1997. The presence of an R467K amino acid substitution and loss of allelic variation correlate with an azole-resistant lanosterol 14 α demethylase in *Candida albicans*. *Antimicrob. Agents Chemother.* **41**:1488–1494.
33. **Wingard, J. R., W. G. Merz, M. G. Rinaldi, T. R. Johnson, J. E. Karp, and R. Saral.** 1991. Increase in *Candida krusei* infection among patients with bone marrow transplantation and neutropenia treated prophylactically with fluconazole. *N. Engl. J. Med.* **325**:1274–1277.