

Alterations in GyrA and ParC Associated with Fluoroquinolone Resistance in *Enterococcus faecium*

NAGWA EL AMIN, SHAH JALAL, AND BENGT WRETLIND*

Division of Clinical and Oral Bacteriology, Department of Immunology, Microbiology, Pathology, and Infectious Diseases, Huddinge University Hospital, S-14186 Huddinge, Sweden

Received 5 October 1998/Returned for modification 10 November 1998/Accepted 28 January 1999

High-level quinolone resistance in *Enterococcus faecium* was associated with mutations in both *gyrA* and *parC* genes in 10 of 11 resistant strains. One low-level resistant strain without such mutations may instead possess an efflux mechanism or alterations in the other subunits of the gyrase or topoisomerase IV genes. These findings are similar to those for other gram-positive bacteria, such as *Enterococcus faecalis*.

Enterococci are important pathogens and a common cause of serious nosocomial infections such as bacteremia and endocarditis (21). *Enterococcus faecalis* is the most common isolate among enterococci. However, *Enterococcus faecium* has become increasingly important after the emergence of ampicillin and vancomycin resistance. Clinical isolates resistant to vancomycin are usually resistant to quinolones and almost all currently available antibiotics (1, 2, 10), but new fluoroquinolones may still have therapeutic value for enterococcal infections (12). At Huddinge Hospital, where this study was conducted, the frequency of quinolone-resistant enterococci has increased from 11% in 1994 to 25% in 1997.

Recent studies showed that fluoroquinolone resistance in gram-negative organisms is associated with mutations in the *gyrA* gene (subunit of DNA gyrase) or with a reduced accumulation of the drug (3, 9, 18, 22, 23, 29, 30). In gram-positive bacteria, fluoroquinolone resistance can be associated with mutations in the *gyrA* or *parC* (subunit of topoisomerase IV) gene (6, 7, 16, 17, 19, 26), and in some gram-positive organisms, such as *Staphylococcus aureus*, efflux pumps have been identified (6, 7, 13, 16, 17, 19, 26, 31).

The nucleotide sequence encoding 41 amino acids in the gyrase A protein, corresponding to *Escherichia coli* amino acids 67 to 106 in the GyrA protein (5, 8, 25, 27, 28), was identified in both gram-negative and gram-positive organisms as the quinolone resistance determining region.

In this study we have analyzed clinical quinolone-resistant *E. faecium* isolates for mutations in the genes encoding subunits of the quinolone target enzymes GyrA and ParC. Twelve clinical isolates of *E. faecium*, collected in different Swedish hospitals, were studied. The isolates were identified by using API 20 STREP (bioMérieux, Marcy l'Etoile, France). The MICs of ciprofloxacin and trovafloxacin were determined by using the E test (Biodisk, Solna, Sweden) on Iso-Sensitest agar (Oxoid) or, for those strains for which MICs of ciprofloxacin were ≥ 32 $\mu\text{g/ml}$, the agar dilution method on PDM-ASM agar (Biodisk) with 5% horse blood (4). The antibiotic susceptibilities are shown in Table 1. Of the 12 isolates, 10 were resistant (here defined as any strain for which the MIC of ciprofloxacin is > 8 $\mu\text{g/ml}$), one was susceptible, and for another the MICs of both drugs were slightly higher than for the reference strain. The type strain *E. faecium* ATCC 19434 was used as a reference.

Genomic DNA was extracted from the 11 isolates and the type strain, by using the guanidium thiocyanate method (24). The target regions were amplified by using PCR with primers derived from *E. faecalis* sequences. A DNA fragment of 241 bp from the *gyrA* gene corresponding to the quinolone resistance determining region was amplified with the 23-mer oligonucleotide primers 5'-CGG GAT GAA CGA ATT GGG TGT GA-3' and 5'-AAT TTT ACT CAT ACG TGC TTC GG-3', equivalent to nucleotide positions 150 to 172 and 368 to 390 of the *E. coli gyrA* gene (15), respectively, and a 191-bp *parC* fragment was amplified with the 20-mer oligonucleotide primers 5'-AAT GAA TAA AGA TGG CAA TA-3' and 5'-CGC CAT CCA TAC TTC CGT TG-3' (positions 10 to 29 and 181 to 200 of the *E. faecalis parC* gene, respectively) (14). One-step PCR was performed for each gene in a 50- μl reaction mixture containing 1.25 U of *Taq* DNA polymerase, 200 μM concentrations of each deoxynucleoside triphosphate, 45 pmol of each primer, 1.5 mM MgCl_2 , and 10 \times PCR buffer (Sigma). Each reaction was run for 30 cycles with the following temperature profile: denaturation at 94°C for 1 min, annealing at 55°C for *gyrA* and 48.7°C for *parC* for 1 min, and then extension at 72°C for 1 min. The amplified fragments were processed for cycle sequencing by using the ABI PRISM Big-dye terminator kit,

TABLE 1. Alterations in *gyrA* and *parC* genes in clinical isolates of *E. faecium*

ATCC strain	MIC ($\mu\text{g/ml}$) ^a		Amino acid (codon) at position ^b		
	CIP	TRV	GyrA		ParC, 80
			83	87	
19434	4	1	Ser (AGT)	Glu (GAG)	Ser (AGC)
3797	1	0.125			
216	8	4			
214	32	8		Lys (AAG)	Ile (ATC)
211	32	8		Lys (AAG)	Ile (ATC)
209	32	16		Lys (AAG)	Ile (ATC)
210	256	8		Lys (AAG)	Ile (ATC)
3784	>256	4		Lys (AAG)	Ile (ATC)
208	64	16	Arg (CGT)		Ile (ATC)
221	256	32	Ile (ATT)		Ile (ATC)
215	256	8	Tyr (TAT)		Ile (ATC)
213	>256	16	Tyr (TAT)		Ile (ATC)
212	>256	32	Tyr (TAT)		Ile (ATC)

^a CIP, ciprofloxacin; TRV, trovafloxacin.

^b Positions correspond to *E. coli*. For those strains with no amino acid shown, the codon is identical to that of the type strain.

* Corresponding author. Mailing address: Division of Clinical and Oral Bacteriology, Huddinge Hospital F82, S-14186 Huddinge, Sweden. Phone: 46-8-5858 7841. Fax: 46-8-711 3918. E-mail: bengt.wretlind@impi.ki.se.

- Activity of two novel fluoroquinolones (DU-6859a and DV-7751a) tested against glycopeptide-resistant enterococcal isolates. *Diagn. Microbiol. Infect. Dis.* **23**:123–127.
13. **Kaatz, G. W., S. M. Seo, and C. A. Ruble.** 1993. Efflux-mediated fluoroquinolone resistance in *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **37**:1086–1094.
 14. **Kanematsu, E., T. Deguchi, M. Yasuda, T. Kawamura, Y. Nishino, and Y. Kawada.** 1998. Alterations in the GyrA subunit of DNA gyrase and the ParC subunit of DNA topoisomerase IV associated with quinolone resistance in *Enterococcus faecalis*. *Antimicrob. Agents Chemother.* **42**:433–435.
 15. **Korten, V., W. M. Huang, and B. E. Murray.** 1994. Analysis by PCR and direct DNA sequencing of *gyrA* mutations associated with fluoroquinolone resistance in *Enterococcus faecalis*. *Antimicrob. Agents Chemother.* **38**:2091–2094.
 16. **Muñoz, R., and A. G. de la Campa.** 1996. ParC subunit of DNA topoisomerase IV of *Streptococcus pneumoniae* is a primary target of fluoroquinolones and cooperates with DNA gyrase A subunit in forming resistance phenotype. *Antimicrob. Agents Chemother.* **40**:2252–2257.
 17. **Ng, E. Y., M. Trucksis, and D. Hooper.** 1996. Quinolone resistance mutations in topoisomerase IV: relationship to *flqA* locus and genetic evidence that topoisomerase IV is the primary target and DNA gyrase is the secondary target of fluoroquinolones in *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **40**:1881–1888.
 18. **Oram, M., and L. M. Fisher.** 1991. 4-Quinolone resistance mutations in the DNA gyrase of *Escherichia coli* clinical isolates identified by using the polymerase chain reaction. *Antimicrob. Agents Chemother.* **35**:387–389.
 19. **Pan, X.-S., J. Ambler, S. Mehtar, and L. M. Fisher.** 1994. Involvement of topoisomerase IV and DNA gyrase as ciprofloxacin targets in *Streptococcus pneumoniae*. *Antimicrob. Agents Chemother.* **40**:2321–2326.
 20. **Pan, X.-S., and L. M. Fisher.** 1997. Targeting of DNA gyrase in *Streptococcus pneumoniae* by sparfloxacin: selective targeting of gyrase or topoisomerase IV by quinolones. *Antimicrob. Agents Chemother.* **41**:471–474.
 21. **Patterson, J. E., A. H. Sweeney, M. Simms, N. Carley, R. Mangi, J. Sabetta, and R. W. Lyons.** 1995. An analysis of 110 serious enterococcal infections. Epidemiology, antibiotic susceptibility, and outcome. *Medicine* **74**:191–200.
 22. **Piddock, L. J.** 1991. Mechanism of quinolone uptake into bacterial cells. *J. Antimicrob. Chemother.* **27**:399–403.
 23. **Piddock, L. J., D. J. Griggs, M. C. Hall, and Y. F. Jin.** 1993. Ciprofloxacin resistance in clinical isolates of *Salmonella typhimurium* obtained from two patients. *Antimicrob. Agents Chemother.* **37**:662–666.
 24. **Pitcher, D. G., N. A. Saunders, and R. J. Owen.** 1989. Rapid extraction of bacterial genomic DNA with guanidium thiocyanate. *Lett. Appl. Microbiol.* **8**:151–156.
 25. **Takenouchi, T., C. Ishii, M. Sugawara, Y. Tokue, and S. Ohya.** 1995. Incidence of various *gyrA* mutants in 451 *Staphylococcus aureus* strains isolated in Japan and their susceptibilities to 10 fluoroquinolones. *Antimicrob. Agents Chemother.* **39**:1414–1418.
 26. **Tankovic, J., B. Perichon, J. Duval, and P. Courvalin.** 1996. Contribution of mutations in *gyrA* and *parC* genes to fluoroquinolone resistance of mutants of *Streptococcus pneumoniae* obtained in vivo and in vitro. *Antimicrob. Agents Chemother.* **40**:2505–2510.
 27. **Truong, Q. C., J.-C. Nguyen Van, D. Shlaes, L. Gutmann, and N. J. Moreau.** 1997. A novel, double mutation in DNA gyrase A of *Escherichia coli* conferring resistance to quinolone antibiotics. *Antimicrob. Agents Chemother.* **41**:85–90.
 28. **Waters, B., and J. Davies.** 1997. Amino acid variation in the GyrA subunit of bacteria potentially associated with natural resistance to fluoroquinolone antibiotics. *Antimicrob. Agents Chemother.* **41**:2766–2769.
 29. **Yoshida, H., M. Nakamura, M. Bogaki, and S. Nakamura.** 1990. Proportion of DNA gyrase mutants among quinolone-resistant strains of *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **34**:1273–1275.
 30. **Yoshida, H., M. Nakamura, M. Bogaki, H. Ito, T. Kojima, H. Hattori, and S. Nakamura.** 1993. Mechanism of action of quinolones against *Escherichia coli* DNA gyrase. *Antimicrob. Agents Chemother.* **37**:839–845.
 31. **Yoshida, H., M. Bogaki, S. Nakamura, K. Ubukata, and M. Konno.** 1990. Nucleotide sequence and characterization of the *Staphylococcus aureus* *norA* gene, which confers resistance to quinolones. *J. Bacteriol.* **172**:6942–6949.