

Correlation between the Resistance Genotype Determined by Multiplex PCR Assays and the Antibiotic Susceptibility Patterns of *Staphylococcus aureus* and *Staphylococcus epidermidis*

FRANCIS MARTINEAU,^{1,2} FRANÇOIS J. PICARD,¹ NICOLAS LANSAC,^{1,2} CHRISTIAN MÉNARD,¹
PAUL H. ROY,^{1,3} MARC OUELLETTE,^{1,2} AND MICHEL G. BERGERON^{1,2*}

Centre de Recherche en Infectiologie de l'Université Laval,¹ Division de Microbiologie,² Faculté de Médecine,
Université Laval, Québec G1V 4G2, and Département de Biochimie, Université Laval, Québec G1K 7P4,³ Canada

Received 22 March 1999/Returned for modification 27 June 1999/Accepted 22 October 1999

Clinical isolates of *Staphylococcus aureus* (a total of 206) and *S. epidermidis* (a total of 188) from various countries were tested with multiplex PCR assays to detect clinically relevant antibiotic resistance genes associated with staphylococci. The targeted genes are implicated in resistance to oxacillin (*mecA*), gentamicin [*aac(6')*-*aph(2'')*], and erythromycin (*ermA*, *ermB*, *ermC*, and *msrA*). We found a nearly perfect correlation between genotypic and phenotypic analysis for most of these 394 strains, showing the following correlations: 98% for oxacillin resistance, 100% for gentamicin resistance, and 98.5% for erythromycin resistance. The discrepant results were (i) eight strains found to be positive by PCR for *mecA* or *ermC* but susceptible to the corresponding antibiotic based on disk diffusion and (ii) six strains of *S. aureus* found to be negative by PCR for *mecA* or for the four erythromycin resistance genes targeted but resistant to the corresponding antibiotic. In order to demonstrate *in vitro* that the eight susceptible strains harboring the resistance gene may become resistant, we subcultured the susceptible strains on media with increasing gradients of the antibiotic. We were able to select cells demonstrating a resistant phenotype for all of these eight strains carrying the resistance gene based on disk diffusion and MIC determinations. The four oxacillin-resistant strains negative for *mecA* were PCR positive for *blaZ* and had the phenotype of β -lactamase hyperproducers, which could explain their borderline oxacillin resistance phenotype. The erythromycin resistance for the two strains found to be negative by PCR is probably associated with a novel mechanism. This study reiterates the usefulness of DNA-based assays for the detection of antibiotic resistance genes associated with staphylococcal infections.

Nosocomial infections caused by multiresistant staphylococci are a growing problem for many health care institutions (26, 42, 50). Of all species of staphylococci, *Staphylococcus epidermidis* and *S. aureus* have the greatest pathogenic potential. *S. epidermidis* is widely recognized as one of the etiologic agents of bacteremia, postoperative cardiac infections and endocarditis, osteomyelitis, urinary tract infections, and peritonitis caused by ambulatory dialysis, with a frequent association with colonization of intravascular catheters and orthopedic devices (26, 50). As for *S. aureus*, it is responsible for diseases caused by exotoxin production (toxic shock and staphylococcal scalded-skin syndromes) and by direct invasion and systemic dissemination (bacteremia, septic shock syndrome, skin infections, and abscesses) (7, 54).

Methicillin-resistant staphylococci (MRS) are resistant to all penicillins, including semisynthetic penicillinase-resistant congeners, penems, carbapenems, and cephalosporins. The basis of this resistance is conferred by an additional penicillin-binding protein, PBP-2' (or PBP-2a), which is absent in methicillin-susceptible strains (11, 15). Plasmid-mediated aminoglycoside-modifying enzymes of all three classes (aminoglycoside phosphotransferases, acetyltransferases, and nucleotidyltransferases) have been found in staphylococci (53). The bifunctional enzyme AAC(6')/APH(2''), encoded by the *aac(6')*-*aph(2'')* gene, inactivates a broad range of clinically useful aminoglycosides such as gentamicin, tobramycin, netilmicin,

and amikacin (20, 49) and is the most frequently encountered aminoglycoside resistance mechanism among staphylococcal isolates (8). Resistance to erythromycin in staphylococci is usually associated with resistance to other macrolides, to the lincosamides, and to type B streptogramin (MLS). This resistance is mediated by a single alteration in the ribosome, the N⁶-dimethylation of an adenine residue in the 23S rRNA. This dimethylation leads to a conformational change in the ribosome, rendering the strain resistant to most antibiotic of the MLS group. Three genes (*ermA*, *ermB*, and *ermC*) encoding methylases have been found in staphylococci (18, 25, 55). Another mechanism of inducible resistance to erythromycin is conferred by the gene *msrA*, which encodes an ATP-dependent efflux pump (47, 48, 65).

The aim of this study was to develop rapid multiplex PCR assays for the detection of clinically relevant antibiotic resistance genes in staphylococci and the identification of the staphylococcal species and to compare those PCR assays with standard microbiological methods for susceptibility testing and microbial identification. In this study, a panel of 206 strains of *S. aureus* and 188 strains of *S. epidermidis* from various sources were tested by (i) conventional susceptibility testing methods and (ii) PCR for the antibiotic resistance genes *mecA*, *aac(6')*-*aph(2'')*, *ermA*, *ermB*, *ermC*, and *msrA*.

(This study was presented in part at the 98th General Meeting of the American Society for Microbiology, Atlanta, Ga., May 1998.)

* Corresponding author. Mailing address: Centre de Recherche en Infectiologie, CHUQ (Pavillon CHUL), 2705 Blvd. Laurier, Ste-Foy, Québec, Canada G1V 4G2. Phone: (418) 654-2705. Fax: (418) 654-2715. E-mail: Michel.G.Bergeron@crchul.ulaval.ca.

MATERIALS AND METHODS

Bacterial strains. A total of 206 *S. aureus* and 188 *S. epidermidis* strains were used in this study. These isolates were obtained from the American Type Culture

TABLE 1. Antibiotic resistance gene-specific and species-specific primers used in this study

Target gene	Accession number (GenBank)	Primer sequence	Annealing position
<i>aac(6')-aph(2'')</i>	M13771	5'-TTG GGA AGA TGA AGT TTT TAG A-3'	159–180
		5'-CCT TTA CTC CAA TAA TTT GGC T-3'	311–332
<i>blaZ</i>	M60253	5'-ACT TCA ACA CCT GCT GCT TTC-3'	511–531
		5'-TGA CCA CTT TTA TCA GCA ACC-3'	663–683
<i>ermA</i>	K02987	5'-TAT CTT ATC GTT GAG AAG GGA TT-3'	370–392
		5'-CTA CAC TTG GCT TAG GAT GAA A-3'	487–508
<i>ermB</i>	U35228	5'-CTA TCT GAT TGT TGA AGA AGG ATT-3'	366–389
		5'-GTT TAC TCT TGG TTT AGG ATG AAA-3'	484–507
<i>ermC</i>	M17990	5'-CTT GTT GAT CAC GAT AAT TTC C-3'	214–235
		5'-ATC TTT TAG CAA ACC CGT ATT C-3'	382–403
Internal control (16S rRNA gene)	J01859	5'-GGA GGA AGG TGG GGA TGA CG-3'	1244–1263
		5'-ATG GTG TGA CGG GCG GTG TG-3'	1469–1489
<i>mecA</i>	X52594	5'-AAC AGG TGA ATT ATT AGC ACT TGT AAG-3'	1059–1085
	X52593	5'-ATT GCT GTT AAT ATT TTT TGA GTT GAA-3'	1206–1232
<i>msrA</i>	X52085	5'-TCC AAT CAT TGC ACA AAA TC-3'	891–910
		5'-AAT TCC CTC TAT TTG GTG GT-3'	1034–1053
<i>S. aureus</i> ^a	AF033191	5'-AAT CTT TGT CGG TAC ACG ATA TTC TTC ACG-3'	5–34
		5'-CGT AAT GAG ATT TCA GTA GAT AAT ACA ACA-3'	83–112
<i>S. epidermidis</i> ^a	NA ^b	5'-ATC AAA AAG TTG GCG AAC CTT TTC A-3'	21–45
		5'-CAA AAG AGC GTG GAG AAA AGT ATC A-3'	121–145

^a The *S. aureus*- and *S. epidermidis*-specific primer pairs were derived from genomic DNA fragments of unknown coding potential (34, 35).

^b NA, not applicable.

Collection (ATCC) (10 strains), the microbiology laboratory of the Centre Hospitalier Universitaire de Québec, The Pavillon Centre Hospitalier de l'Université Laval (CHUL) (Ste-Foy, Québec, Canada) (184 strains), the Laboratoire de Santé Publique du Québec (LSPQ) (Sainte-Anne-de-Bellevue, Québec, Canada) (80 strains), the microbiology laboratory of Hôpital Laval (Ste-Foy, Québec, Canada) (91 strains), the Mount Sinai Hospital (Toronto, Ontario, Canada) (5 strains), the Huashan Hospital (Shanghai, China) (21 strains), and the Institut Pasteur (Paris, France) (3 strains). The 184 staphylococcal strains from the Pavillon CHUL were identified by using the MicroScan Autoscan-4 system equipped with the Positive BP Combo Panel Type 6 (Dade Diagnostics, Mississauga, Ontario, Canada). A reconfirmation of the staphylococcal species identification for the 210 remaining clinical isolates was performed by using the MicroScan Autoscan-4 system. There was no known relationship between any of the patients. These isolates were implicated in a variety of classical staphylococcal diseases. Duplicate isolates from the same patients, even if the site of infection was different, were excluded from this study. Strains were cultured on sheep blood agar or in brain heart infusion (BHI). Stock cultures were stored frozen (–80°C) in BHI containing 10% glycerol.

Susceptibility testing. (i) **Disk diffusion.** Disk diffusion tests were performed for each of the 394 isolates previously identified as *S. aureus* or *S. epidermidis* by following the method recommended by the National Committee for Clinical Laboratory Standards (NCCLS) (39). Disks (Becton Dickinson Microbiology Systems, Cockeysville, Md.) containing 1 µg of oxacillin, 10 µg of gentamicin, or 15 µg of erythromycin were added to inoculated Mueller-Hinton agar plates. Subsequently, they were incubated 24 h at 35°C. Any growth, including pinpoint colonies, within the 10-mm-diameter zone for oxacillin, the 12-mm-diameter zone for gentamicin, and the 13-mm-diameter zone for erythromycin was considered indicative of resistance. *S. aureus* ATCC 29213 was used as negative control.

(ii) **MIC determinations.** MICs of oxacillin, gentamicin, and erythromycin (ranging for each antibiotic from 0 to 64 µg/ml, by serial twofold dilutions) were determined for each discordant result between genotype and phenotype by the broth microdilution methodology as recommended by the NCCLS (38). The

trays were incubated at 35°C and were read for turbidity with indirect light after 24 h. *S. aureus* ATCC 29213 was used as negative control.

(iii) **Breakpoint determination.** Breakpoints for oxacillin, gentamicin, and erythromycin were determined by using the automated microbial identification system MicroScan Autoscan-4 equipped with the Positive BP Combo Panel Type 6 (Dade Diagnostics). *S. aureus* ATCC 29213 was used as negative control.

(iv) **Nitrocefin test.** The chromogenic cephalosporin nitrocefin disk test was used as recommended by the manufacturer (Becton Dickinson) on each of the four *S. aureus* strains found to be negative for *mecA* by PCR assays but resistant to oxacillin. Also these strains were tested for susceptibility to amoxicillin and clavulanic acid (20 and 10 µg, respectively) by disk diffusion according to NCCLS guidelines. *S. aureus* ATCC 43300 was used as a positive control, and *S. aureus* ATCC 25923 was used as a negative control.

Multiplex PCR. *S. aureus*- and *S. epidermidis*-specific PCR assays used in this study have been previously described by us (34, 35). PCR primers were chosen from the antibiotic resistance genes *mecA*, *aac(6')-aph(2'')*, *ermA*, *ermB*, *ermC*, and *msrA* (Table 1). Primers were designed with the help of the Oligo Primer Analysis software version 4.0 (National Biosciences, Inc., Plymouth, Minn.) and synthesized by using a model 391 DNA synthesizer (Perkin-Elmer Corp./Applied Biosystems Division, Foster City, Calif.).

Multiplex PCR assays were all performed directly from a bacterial suspension whose turbidity was adjusted to that of a 0.5 McFarland standard, which corresponds to approximately 1.5×10^8 bacteria per ml. Then, 1 µl of the standardized bacterial suspension was transferred directly to a 20-µl PCR mixture containing 50 mM KCl, 10 mM Tris-HCl (pH 9.0), 0.1% Triton X-100, 2.5 mM MgCl₂, 0.4 µM concentrations (each) of the specific primers, 200 µM concentrations (each) of the four deoxynucleoside triphosphates, and 0.5 U of *Taq* DNA polymerase (Promega). In order to reduce the formation of nonspecific extension products, a "hot-start" protocol was used (34). The PCR mixtures were subjected to thermal cycling (3 min at 96°C and then 30 cycles of 1 s at 95°C for the denaturation step and 30 s at 55°C for the annealing-extension step) with a PTC-200 thermal cycler (MJ Research, Inc., Watertown, Mass.). All multiplex PCR assays (Table 2) also included a primer pair specific to conserved regions of

TABLE 2. Multiplex PCR assays

Multiplex no. ^a	PCR primer pairs combined (amplicon sizes)
1.....	<i>S. aureus</i> (108 bp) + <i>S. epidermidis</i> (125 bp) + <i>mecA</i> (174 bp)
2.....	<i>S. aureus</i> (108 bp) + <i>S. epidermidis</i> (125 bp) + <i>aac(6')-aph(2'')</i> (174 bp)
3.....	<i>S. aureus</i> (108 bp) + <i>S. epidermidis</i> (125 bp) + <i>ermA</i> (139 bp)
4.....	<i>S. aureus</i> (108 bp) + <i>S. epidermidis</i> (125 bp) + <i>ermB</i> (142 bp)
5.....	<i>S. aureus</i> (108 bp) + <i>S. epidermidis</i> (125 bp) + <i>ermC</i> (190 bp)
6.....	<i>S. aureus</i> (108 bp) + <i>S. epidermidis</i> (125 bp) + <i>msrA</i> (163 bp)
7.....	<i>S. aureus</i> (108 bp) + <i>S. epidermidis</i> (125 bp) + <i>blaZ</i> (173 bp)

^a All multiplex PCR assays included an internal control (241 bp).




FIG. 1. Example of multiplex PCR amplifications with the *mecA*-, *S. aureus*-, *S. epidermidis*-, and internal control-specific primer pairs. PCR assays were performed either from 10 pg of purified genomic DNA (lanes 2 to 14) or from 1 µl of a bacterial suspension whose turbidity was adjusted to that of a 0.5 McFarland standard (lanes 15 to 18). The content of each lane is as follows: 2, *S. aureus* ATCC 43300 (Oxa^r); 3, *S. aureus* ATCC 25923 (Oxa^s); 4, *S. epidermidis* ATCC 35983 (Oxa^r); 5, *S. epidermidis* ATCC 14990 (Oxa^s); 6, *S. capitis* ATCC 49326 (Oxa^r); 7, *S. haemolyticus* ATCC 29970 (Oxa^s); 8, *S. hominis* ATCC 27844 (Oxa^s); 9, *S. saprophyticus* ATCC 15305 (Oxa^s); 10, *S. simulans* ATCC 27848 (Oxa^s); 11, *S. warneri* ATCC 27836 (Oxa^r); 12, *Enterococcus faecalis* ATCC 29212; 13, *Enterococcus faecium* ATCC 51559; 14, *Streptococcus pneumoniae* ATCC 27336; 15, *S. aureus* ATCC 43300 (Oxa^r); 16, *S. aureus* ATCC 25923 (Oxa^s); 17, *S. epidermidis* ATCC 35983 (Oxa^r); and 18, *S. epidermidis* ATCC 14990 (Oxa^s). Lanes 1 and 19, controls to which no DNA was added; lane M, 50-bp ladder molecular size standard.

the 16S rRNA gene (241-bp amplicon) which was used to provide an internal control (34). This allowed us to control the efficiency of the quick protocol for bacterial lysis of the PCR assays and to ensure that significant PCR inhibition was absent. Of the PCR-amplified reaction mixture, 10 µl was resolved by electrophoresis through a 2% agarose gel containing 0.5 µg of ethidium bromide per ml in Tris-borate-EDTA buffer (89 mM Tris, 89 mM boric acid, 2 mM EDTA) at 12 V/cm for 30 min (34) (Fig. 1). The gels were visualized under 254-nm UV lights. The sizes of the amplification products were estimated by comparison with a 50-bp molecular size standard ladder. The total time for the PCR assays from a standardized bacterial suspension was about 1.5 h.

In vitro selection of resistant cells. In order to select resistant cells from susceptible strains carrying the corresponding antibiotic resistance gene, we subcultured them onto medium containing increasing concentration gradients of the antibiotic of interest (9, 57). For medium preparation, 10 ml of autoclaved antibiotic medium 1 (Difco Laboratories) without antibiotic was poured into a round petri plate and allowed to cool with one edge elevated so that the agar level just reached the intersection of the bottom and side of the plate. Subsequently, the plate was leveled and an additional 10 ml of the same medium containing low (1 µg/ml), intermediate (5 µg/ml), or high (10 µg/ml) concentrations of oxacillin or erythromycin was added. After 18 h at room temperature, a gradient of antibiotic concentrations ranging from near 0 µg/ml at one edge to the maximal concentration (near the antibiotic concentration in the added medium) at the opposite edge was established. Inoculated plates were incubated for 18 to 24 h at 35°C. For subculturing, colonies were always picked in the area of highest antibiotic concentrations and streaked onto an agar plate containing the same or an increased gradient of antibiotic concentrations. The resistant cells selected from the plates with the highest concentrations of antibiotic were submitted to MIC determinations to confirm their resistance phenotype. They were also tested with the multiplex PCR assays to verify that their resistance genotype was unchanged. To verify the stability of the selected antibiotic resistance, MICs were reevaluated after three passages in medium without antibiotic. The *S.*

TABLE 3. Correlation between gentamicin resistance and the presence of the resistance gene *aac(6')-aph(2'')*

Species	No. of strains (type) tested by disk diffusion with gentamicin	PCR results for <i>aac(6')-aph(2'')</i>	
		No. positive	No. negative
<i>S. aureus</i>	41 (resistant)	41	0
<i>S. aureus</i>	165 (susceptible)	0	165
<i>S. epidermidis</i>	131 (resistant)	131	0
<i>S. epidermidis</i>	57 (susceptible)	0	57

TABLE 4. Correlation between oxacillin resistance and the presence of the resistance gene *mecA*

Species	No. of strains (type) tested by disk diffusion with oxacillin	PCR results for <i>mecA</i>	
		No. positive	No. negative
<i>S. aureus</i>	66 (resistant)	62	4
<i>S. aureus</i>	140 (susceptible)	1	139
<i>S. epidermidis</i>	146 (resistant)	146	0
<i>S. epidermidis</i>	42 (susceptible)	3	39

aureus strain ATCC 29213, which (i) is susceptible to oxacillin and erythromycin and (ii) does not carry any of the antibiotic resistance gene tested in this study, was always used in parallel as a negative control.

RAPD assay. Randomly amplified polymorphic DNA (RAPD) assay (66) was performed by using the Operon 10-base kit "AD" which contains 20 different oligonucleotides (Operon Technologies, Inc., Alameda, Calif.). For all bacterial species tested, amplification was performed directly from a bacterial suspension whose turbidity was adjusted to that of a 0.5 McFarland standard. The PCR mixtures (50 mM KCl, 10 mM Tris-HCl [pH 9.0], 0.1% Triton X-100, 2.5 mM MgCl₂, a 1.5 µM concentration of a single 10-nucleotide primer, 200 µM concentrations (each) of the four deoxynucleoside triphosphates, and 0.5 U of *Taq* DNA polymerase [Promega]) were subjected to thermal cycling (3 min at 96°C and then 40 cycles of 1 min at 94°C for the denaturation step, 1 min at 32°C for the annealing step, and 2 min at 72°C for the extension step) by using a PTC-200 thermal cycler (MJ Research, Inc.). A 7-min extension step at 72°C was performed to allow the completion of DNA synthesis. Amplification products were analyzed by electrophoresis in 1.8% agarose gels containing 0.5 µg of ethidium bromide per ml.

RESULTS

Correlation between susceptibility testing and the multiplex PCR assays. We have compared gentamicin, oxacillin, and erythromycin susceptibility results determined by the disk diffusion method for 394 staphylococcal clinical isolates with those obtained by the multiplex PCR assays for the detection of antibiotic resistance genes. The multiplex PCR assays allowed us also to verify the identity of all *S. aureus* and *S. epidermidis* strains by using the species-specific PCR assays previously described by our group (34, 35). The various multiplex PCR assays developed in this study are given in Table 2. A comparison of the multiplex PCR-based assays with conventional susceptibility testing and identification methods (disk diffusion, MIC determination, and the MicroScan identification system) showed the following correlations: 100% for gentamicin resistance (Table 3), 98% for oxacillin resistance (Table 4), 98.5% for erythromycin resistance (Table 5), and 100% for staphylococcal species identification.

The correlation between gentamicin resistance and the presence of *aac(6')-aph(2'')* is summarized in Table 3. Of 206 *S. aureus* strains, 165 (80%) were susceptible to gentamicin as determined by the disk diffusion method and negative by PCR for *aac(6')-aph(2'')*. The remaining 41 *S. aureus* strains were

TABLE 5. Correlation between erythromycin resistance and the presence of the resistance gene *ermA*, *ermB*, *ermC*, or *msrA*

Species	No. of strains (type) tested by disk diffusion with erythromycin	No. of strains positive by PCR for:				No. of PCR-negative strains
		<i>ermA</i>	<i>ermB</i>	<i>ermC</i>	<i>msrA</i>	
<i>S. aureus</i>	73 (resistant)	43	5	21	2	2
<i>S. aureus</i>	133 (susceptible)	0	0	4	0	129
<i>S. epidermidis</i>	142 (resistant)	9	1	124	8	0
<i>S. epidermidis</i>	46 (susceptible)	0	0	0	0	46

TABLE 6. Characteristics of the four strains resistant to oxacillin but negative for *mecA* by PCR

Strains	Oxacillin MIC ($\mu\text{g/ml}$)	Nitrocefin test result	Disk diffusion test result (mm) amoxicillin with clavulanic acid ^a	<i>blaZ</i> (PCR detection)
<i>S. aureus</i> R620	4	+	24.0	+
<i>S. aureus</i> R626	8	+	22.0	+
<i>S. aureus</i> R627	4	+	24.5	+
<i>S. aureus</i> STA572	8	+	23.3	+
<i>S. aureus</i> ATCC 43300 ^b	64	+	15.2	+
<i>S. aureus</i> ATCC 25923 ^b	1	–	28.3	–

^a Strains were considered β -lactamase hyperproducers when (i) the diameter of the inhibition zone with the amoxicillin-clavulanic acid (20 and 10 μg , respectively) disk exceeded 20 mm and (ii) they were resistant to oxacillin (2).

^b *S. aureus* ATCC 43300 is a nonhyperproducing strain carrying *mecA*, while *S. aureus* ATCC 25923 is a β -lactamase-negative strain not carrying *mecA*.

resistant to gentamicin while harboring the *aac(6')-aph(2'')*. Interestingly, the situation is quite different for *S. epidermidis*. Of 188 *S. epidermidis* strains, 131 (70%) were resistant to gentamicin and harbor the *aac(6')-aph(2'')* resistance gene. The remaining 57 strains were negative by PCR for *aac(6')-aph(2'')* and susceptible to gentamicin.

The correlation between oxacillin resistance and the presence of the *mecA* gene is summarized in Table 4. There were 139 strains (68%) of *S. aureus* which were both negative by PCR for *mecA* and susceptible to oxacillin. However, there was one *S. aureus* strain which was oxacillin susceptible and positive for *mecA*. This strain had an MIC of 1 $\mu\text{g/ml}$. Therefore, although present, the *mecA* gene did not confer a detectable level of oxacillin resistance. However, as described below, it was possible to select resistant cells by exposing this strain to increasing concentrations of oxacillin. Of 66 oxacillin-resistant *S. aureus* strains (32%), 62 were harboring *mecA* based on PCR analysis. Importantly, four oxacillin-resistant *mecA*-negative *S. aureus* strains showed a borderline level of resistance based on susceptibility testing by disk diffusion and MIC determination. In fact, these strains all had MICs ranging from 4 to 8 $\mu\text{g/ml}$ for oxacillin. Further testing was performed on these four *S. aureus* strains to verify if they were β -lactamase hyperproducers, a mechanism which may mediate *S. aureus* resistance to methicillin (2). They were characterized by testing with (i) nitrocefin disks, (ii) a PCR assay for the detection of *blaZ*, and (iii) amoxicillin-clavulanic acid disks. Positive reaction to nitrocefin and the presence of the *blaZ* gene revealed by PCR confirmed that these strains were β -lactamase producers. In vitro testing with oxacillin and amoxicillin-clavulanic acid disks suggests that these four strains are β -lactamase hyperproducers, which could explain their borderline oxacillin-resistant phenotype (Table 6).

As expected, there was far more resistance to oxacillin in *S. epidermidis* than in *S. aureus*. There were 146 *S. epidermidis* strains (78%) which were both *mecA* positive and oxacillin resistant. Of the 42 oxacillin-susceptible strains (22%), there were 39 not harboring the *mecA* gene. The three *S. epidermidis* oxacillin-susceptible *mecA*-positive strains had MICs ranging from 0.25 to 1 $\mu\text{g/ml}$. Therefore, as for *S. aureus*, the presence of *mecA* was not sufficient to confer a resistance phenotype in these three *S. epidermidis* strains. Again, as described below, it was possible to select resistant cells from these oxacillin-susceptible strains.

Table 5 shows the correlation between erythromycin resistance and the presence of *ermA*, *ermB*, *ermC*, and *msrA*. As for oxacillin and gentamicin, there was far more resistance to erythromycin in *S. epidermidis* than in *S. aureus*. For *S. aureus*, of the 133 susceptible strains (64%), there were 129 not harboring any of the four genes associated with erythromycin resistance, based on multiplex PCR assays. The four erythro-

mycin-susceptible *S. aureus* strains harboring an erythromycin resistance gene were all positive for *ermC* only (Table 7). Therefore, as observed for oxacillin resistance, the presence of *ermC* did not confer a detectable level of resistance. Again, it was possible to select resistant cells from these four strains. Regarding the *S. aureus* that was resistant to erythromycin, of 73 strains (35%) there were 71 harboring one of the four erythromycin resistance gene tested (Table 4). The incidences of the different genotypes were 21% for *ermA*, 2.4% for *ermB*, 10% for *ermC*, and 1% for *msrA*. None of the resistant strains harbored more than one resistance gene. The erythromycin resistance in the two *S. aureus* strains not carrying any of the four resistance genes tested is likely mediated by as-yet-unknown mechanisms. We are now investigating the mechanisms of such resistance.

For *S. epidermidis*, only 46 strains (24%) were both susceptible to erythromycin and did not contain any erythromycin resistance gene. The incidences of the three erythromycin ribosomal methylase genes tested were 4.8% for *ermA*, 0.5% for *ermB*, and 66% for *ermC*. *msrA* was present in 4.3% of *S. epidermidis* strains. No discrepant results were found between the genotypes and the phenotypes in these 188 *S. epidermidis* clinical strains.

In vitro selection of resistant cells. As shown in Table 7, there were 8 susceptible staphylococcal strains carrying the corresponding antibiotic resistance gene. Four strains (one *S. aureus* strain and three *S. epidermidis* strains) were PCR positive for *mecA* but susceptible for oxacillin, and four strains of *S. aureus* were positive for *ermC* but susceptible to erythromycin. These strains were all initially subcultured twice on plates containing a gradient of oxacillin or erythromycin ranging from 0 to 1 $\mu\text{g/ml}$ in order to reach growth confluence. Subsequently, strains were subcultured twice on plates with gradients ranging from 0 to 5 $\mu\text{g/ml}$ and then twice again on plates with gradients ranging from 0 to 10 $\mu\text{g/ml}$ in order to attain growth confluence.

The culture on media with increasing gradients of the target antibiotic allowed for the selection of cells having MICs greater than 64 $\mu\text{g/ml}$ for both oxacillin and erythromycin. This is a major increase, considering that the MICs for the original susceptible strains ranged from 0.5 to 1 $\mu\text{g/ml}$ for these two antibiotics. Furthermore, we have confirmed by PCR testing that the genotypes of these strains harboring *mecA* or *ermC* remained unchanged after the selection process. Interestingly, even after three passages in antibiotic-free media, the MICs remained unchanged. Importantly, with the control strain ATCC 29213, it was not possible to select cells resistant to oxacillin or erythromycin, thereby confirming that the resistance gene must be present to allow for the rapid selection of resistant cells. The negative control strain always remained

TABLE 7. *S. aureus* ($n = 11$) and *S. epidermidis* ($n = 3$) strains for which the resistance genotype was discordant with the resistance phenotype

Group and strain ^a	Profile	MIC ($\mu\text{g/ml}$)	MIC after selection ^b ($\mu\text{g/ml}$)
Group 1			
<i>S. aureus</i> R632	<i>mecA</i> , Oxa ^s	2	>64
<i>S. epidermidis</i> R588	<i>mecA</i> , Oxa ^s	0.125	>64
<i>S. epidermidis</i> 184	<i>mecA</i> , Oxa ^s	1	>64
<i>S. epidermidis</i> 3348	<i>mecA</i> , Oxa ^s	0.25	>64
<i>S. aureus</i> ATCC 29213	<i>mecA</i> negative, Oxa ^s	0.5	0.5
<i>S. aureus</i> 18	<i>ermC</i> , Ery ^s	1	>64
<i>S. aureus</i> 75	<i>ermC</i> , Ery ^s	1	>64
<i>S. aureus</i> 93	<i>ermC</i> , Ery ^s	1	>64
<i>S. aureus</i> 5919	<i>ermC</i> , Ery ^s	1	>64
<i>S. aureus</i> ATCC 29213	<i>ermC</i> negative, Ery ^s	0.5	0.5
Group 2			
<i>S. aureus</i> R572	<i>mecA</i> negative, Oxa ^r	8	
<i>S. aureus</i> R620	<i>mecA</i> negative, Oxa ^r	4	
<i>S. aureus</i> R626	<i>mecA</i> negative, Oxa ^r	8	
<i>S. aureus</i> R627	<i>mecA</i> negative, Oxa ^r	4	
<i>S. aureus</i> MA-5091	<i>ermABC</i> negative, <i>msrA</i> negative, Ery ^r	>32	
<i>S. aureus</i> R566	<i>ermABC</i> negative, <i>msrA</i> negative, Ery ^r	>32	

^a Group 1, resistance gene present, susceptible; group 2, resistance gene absent, resistant. MIC column data for group 1 strains are the initial MICs.

^b The selection of resistant cells was done by plating on media with increasing gradients of the target antibiotic.

low, at 0.5 $\mu\text{g/ml}$. These experiments strongly suggest that susceptible strains harboring a resistance gene have the potential to develop resistance upon in vivo selection by the appropriate antimicrobial agent.

In order to further confirm that the resistant cells selected were derived from the same susceptible initial strain, RAPD-PCR was performed with genomic DNA purified from the four oxacillin-susceptible *mecA*-positive and the four erythromycin-susceptible *ermC*-positive strains as well as with the resistant cells derived from these eight strains (five *S. aureus* and three *S. epidermidis*) after in vitro selection. The RAPD assay was performed with three different oligonucleotide primers shown to be highly discriminatory for these strains. For the four staphylococcal oxacillin-susceptible *mecA*-positive strains (Table 7), three distinct RAPD patterns were obtained for each oligonucleotide primer tested. When used in combination, these three primers produced unique amplification patterns for each of those four strains. We found that the amplification patterns with the three selected RAPD primers were identical for both the initial susceptible strain and the selected resistant strain (data not shown). These findings strongly suggest that the selected resistant cells were derived from the corresponding original susceptible strain. Similarly, distinct RAPD patterns were obtained for each of the four *S. aureus* erythromycin-susceptible *ermC*-positive strains (Table 7). Again, identical amplification patterns were obtained with both the initial susceptible strain and the selected resistant strain.

DISCUSSION

We describe here PCR primers that can be used to survey clinically relevant antibiotic resistance genes frequently encountered in staphylococci. We have compared multiplex PCR assays for the detection of antibiotic resistance genes with classical methods for the determination of susceptibility to antibiotics. Overall, we found correlations between these two methods of 98% for oxacillin resistance, 100% for gentamicin resistance, 98.5% for erythromycin resistance, and 100% for species identification.

The *mecA* gene is a 2.4-kb chromosomal determinant en-

coding the PBP-2' protein which is not subjected to dissemination among staphylococcal strains via plasmid spread. Expression of PBP-2' is under control of the negative regulation elements *mecI* and *mecRI* (11, 27, 40). Five phenotypic methods are commonly used by clinical laboratories to detect MRS. For all screening methods, oxacillin is used because it is the most sensitive member of the penicillinase-resistant semisynthetic β -lactam agents for the detection of resistance. The easiest and among the most accurate test is the use of oxacillin screening agar (60). Broth dilution methods are generally accurate when cation-adjusted Mueller-Hinton broth containing 2% NaCl is used and when the results are interpreted by visual inspection of broth turbidity (21). Agar disk diffusion is the most used antibiotic susceptibility assay in clinical microbiology laboratories, but it is not particularly accurate because it may demonstrate a lack of reproducibility and sensitivity (2). The Crystal MRSA (Becton Dickinson) is a reliable and rapid method (4 to 5 h) with a fluorescent indicator to detect methicillin-resistant *S. aureus* (MRSA) (46). Finally, the E test is an excellent quantitative system for laboratories that do not wish to stock the more cumbersome MIC trays (41). Although these culture-based methods are generally reliable for detecting MRS, the detection of *mecA* is now considered the gold standard method, mainly because (i) phenotypic methods may be difficult to interpret and (ii) some isolates do not express their *mecA* gene unless selective pressure via antibiotic treatment is applied.

Several studies deal with the detection by PCR of the *mecA* gene only (36, 44, 59) or combined in multiplex with *S. aureus*-specific amplification assays (3, 6, 19, 51, 62). Overall, we found in this study that 30.6% of *S. aureus* and 79.3% of *S. epidermidis* strains were carrying the *mecA* gene. When restricted to strains isolated from Canada (obtained from CHUL, LSPQ, Laval Hospital, and Mount Sinai Hospital), the proportion of strains carrying *mecA* falls to 25.2% for *S. aureus* and 77.1% for *S. epidermidis*. It should be noted that this high percentage of MRS is not representative of the true Canadian incidence because the *S. aureus* isolates obtained from LSPQ and Mount Sinai Hospital were selected for their phenotypic resistance to

oxacillin. When considering only isolates from the CHUL and Laval Hospital, which were not selected for resistant strains, we observed an incidence of 2% of *S. aureus* harboring *mecA*. This is comparable to the study from the SENTRY antimicrobial Surveillance Program for *S. aureus* isolates from Canada, in which an incidence of 2.5% of oxacillin-resistant *S. aureus* was reported (43).

The *aac(6′)-aph(2′)* is the gene coding for the most frequently encountered aminoglycoside modifying enzyme (AME) in gram-positive bacteria, AAC(6′)-APH(2′) (61). This bifunctional enzyme inactivates a broad range of clinically useful aminoglycosides, especially gentamicin and tobramycin, because it catalyzes both acetyltransferase and phosphotransferase reactions. Two other genes associated with aminoglycoside resistance in staphylococci may also be found. These genes, *aphA3* [coding for APH(3′)III enzyme], and *aadC*, coding for ANT(4′,4′) enzyme, are much less frequently encountered than *aac(6′)-aph(2′)* and are not clinically relevant because they mediate resistance to aminoglycosides not usually prescribed to treat staphylococcal infections (53). DNA amplification by PCR has been shown to be a reliable tool for the specific detection of AME genes in gram-negative bacteria as well as for the detection of *aac(6′)-aph(2′)* in epidemic strains of MRSA (61). DNA dot blot analysis of AME genes has also been described (8). In this study, we found *aac(6′)-aph(2′)* in 20% of *S. aureus* isolates and in 69.7% of *S. epidermidis* isolates, with no discrepant results with the resistance phenotypes obtained by the disk diffusion method. The absence of discrepant results between the resistance genotypes and phenotypes suggests that there were no strains harboring (or at least not expressing) aminoglycoside resistance genes other than *aac(6′)-aph(2′)*. Therefore, all isolates harboring the *aac(6′)-aph(2′)* gene were clearly resistant to gentamicin in susceptibility tests, and all isolates not harboring this resistance gene were susceptible to gentamicin. These findings are in agreement with other studies in which it was reported that all aminoglycoside-resistant strains were carrying *aac(6′)-aph(2′)* (16, 31, 32, 45, 61).

Several genes are implicated in erythromycin resistance, especially in staphylococci and streptococci. Simultaneous resistance to macrolides, lincosamides, and type-B streptogramins (MLS resistance) in clinical isolates is a form of acquired resistance due to several evolutionary variants of *erm* genes, which encodes a 23S rRNA methylase (23, 64). The inducible gene *ermA* is found on the transposon Tn554 and has a single specific site for insertion into the *S. aureus* chromosome (37). The *ermB* gene is found on the transposon Tn551 of a penicillinase plasmid (25). The *ermC* gene is responsible for constitutive or inducible resistance to erythromycin and is generally located on small plasmids (18, 63, 64). Staphylococcal strains resistant to macrolides and type-B streptogramins also frequently harbor *msrA*, which encodes an ATP-dependent efflux pump and mediates the macrolide-streptogramin B (MS) resistance (18). Several studies concerning the epidemiological distribution of genes encoding erythromycin ribosomal methylases and efflux pumps have been performed by dot blot or Southern hybridization (18, 23, 65), and detection of erythromycin resistance determinants by PCR has been performed with staphylococci and streptococci (55).

In this study, the incidences of *ermA* in erythromycin-resistant staphylococci were 21% for *S. aureus* and 4.8% for *S. epidermidis*. These findings are in agreement with the study by Eady et al. (18) conducted with coagulase-negative staphylococci (CoNS) in the United Kingdom in which an incidence of 5.9% for *ermA* was reported. In a study performed in Denmark (64), 16% of *S. aureus* strains were carrying *ermA*, while only

3% of CoNS strains had this gene. These observations are also in agreement with our data. On the other hand, Thakker-Varia et al. (58) reported a higher incidence for *ermA*. They found that 31% of *S. aureus* and 19% of CoNS from three New Jersey hospitals were harboring the *ermA* gene. Regarding *ermB*, we found that this gene was less frequently encountered than *ermA* in erythromycin-resistant staphylococci with 1.9% of *S. aureus* and 0.5% of *S. epidermidis* strains carrying *ermB*. In the United Kingdom, an incidence of 7.2% for *ermB* in CoNS has been reported (18). Interestingly, in that study it was observed that *ermB* was found exclusively in animal isolates of *S. intermedius*, *S. xylosum*, and *S. hyicus* but was absent in CoNS of human origin. This observation could explain why our incidence for *ermB* in CoNS is much lower because our CoNS strains were exclusively of human origin. For *ermC*, we found an incidence of 10.2% for *S. aureus* strains and 66% for *S. epidermidis* strains. Eady et al. (18) have also reported a high incidence of CoNS strains carrying *ermC* (i.e., detected in 112 of 221 [50.6%] CoNS isolates). In New Jersey hospitals, an even higher incidence for the *ermC* gene (i.e., 69% for *S. aureus* and 81% for CoNS isolates) has been reported (58). A high incidence of *S. aureus* carrying *ermC* (i.e., 84%) has also been reported in Denmark (64). The much lower prevalence of *ermC*-positive strains found in Canada suggests that the selective pressure for *ermC*-positive *S. aureus* is weaker in this country. These variations may be associated with variable use of erythromycin in each country. Finally, among erythromycin-resistant strains, we found *msrA* in only 1% of *S. aureus* strains and 4.3% of *S. epidermidis* strains. A much higher incidence for *msrA* in MRS (i.e., 33%) has been reported in the United Kingdom (18). Again, these results suggest a lower selective pressure on populations of *msrA*-positive staphylococci in Canada.

Among the staphylococcal strains showing discrepancies between the genotype and the phenotype (four oxacillin-susceptible *mecA*-positive strains and four erythromycin-susceptible *ermC*-positive strains), we were able to select cells demonstrating a resistant phenotype from all of them. Others have also reported *S. aureus* strains carrying the *mecA* gene but susceptible to oxacillin (28, 40). The heterogeneous nature of methicillin and erythromycin resistances suggests that numerous factors could explain the sensitive phenotype in these strains (1, 10, 22, 29, 30, 40, 56). Such factors include (i) the regulation of the expression of *mecA* or *ermC* and (ii) the absence of host factors associated with the phenotypic expression of methicillin resistance. The fact that we were able to select resistant cells from originally susceptible strains demonstrates that upon in vitro selection in the presence of increasing gradients of the antimicrobial agent, it is possible to select for resistance. Furthermore, once induced, the resistance phenotype was shown to be stable. Kolbert et al. (28) were also able to select for oxacillin resistance for 6 of 10 *mecA*-positive and oxacillin-sensitive isolates. Our ability to select resistant cells from all *mecA*-positive and oxacillin-susceptible strains is probably explained by the fact that we used plates with medium containing an increasing concentration gradient of oxacillin (9, 57) as opposed to the method used by Kolbert et al., in which a series of plates containing increasing concentrations of oxacillin were inoculated. Consequently, the use of plates with an increasing concentration gradient of antibiotic appear more efficient to select for resistant cells. To our knowledge, the selection of erythromycin-resistant cells from originally susceptible bacterial strains has not been previously reported. From a clinical perspective, these findings suggest that a susceptible strain harboring but not expressing an antibiotic resistance gene should be regarded as potentially resistant to that antibiotic.

Well-documented clinical observations of this phenomenon have not been demonstrated with erythromycin or a macrolide-like antibiotic. Investigators using teicoplanin have denoted a failure of treatment associated with an increase in MIC during therapy of *S. aureus* septicemia (12, 24, 33).

The second type of discrepancy was encountered in four *mecA*-negative *S. aureus* strains which were borderline oxacillin resistant based on MIC determinations. Since these four strains were positive for the nitrocefin test and positive by PCR for the *blaZ* gene and were sensitive to amoxicillin-clavulanic acid (i.e., inhibition zone of >20 mm), they appeared to be β -lactamase hyperproducers, which could explain their borderline resistance to oxacillin. Others (2, 14, 51) have also used this criteria to determine the phenotype of β -lactamase hyperproducers. However, we cannot exclude the possibility that the methicillin resistance in these strains, which is not mediated by *mecA*, is associated with other mechanisms of resistance (2).

We also reported two *S. aureus* strains fully resistant (>32 μ g/ml) to erythromycin but negative for *ermA*, *ermB*, *ermC*, and *msrA* by PCR. Based on these results, we postulate that this resistance is probably associated with a novel mechanism not yet characterized in staphylococci. As reported recently by Huovinen et al. (52), a novel erythromycin methylase resistance gene, *ermTR*, was discovered in *Streptococcus*. The nucleotide sequence of *ermTR* is 82.5% identical to the staphylococcal *ermA*. Another finding concerns a novel macrolide efflux pump from *Streptococcus agalactiae*, called *mreA* (13). This putative efflux determinant is distinct from the multicomponent MsrA pump found in staphylococci. Consequently, it is possible that novel erythromycin resistance gene(s) in staphylococci will be found in the future.

This study reiterates the usefulness of DNA-based assays for the detection of antibiotic resistance genes associated with staphylococcal infections. However, the approach presented in this study would not be appropriate for detecting the clinically important resistance to fluoroquinolones because it is associated with many point mutations in two different genes (i.e., *gyrA* and *grlA*) (17). On the other hand, such complex resistance genotypes could be detected by PCR amplification of the mutated genetic target, followed by hybridization on a DNA chip or rapid automated DNA sequencing.

In the clinical setting, the simultaneous identification of the bacteria and determination of its susceptibility to antibiotics generally require 48 h (4, 5). Yet in the choice of empiric antibiotic therapy for suspected staphylococcal sepsis, the clinician must know rapidly which species is involved and its susceptibility to antibiotics. The multiplex PCR assays developed in this study could be adapted for direct detection from positive blood cultures or from clinical specimens (e.g., detection of MRSA from nasal swabs or infected wounds), thereby allowing a much faster diagnosis than conventional culture methods. We believe that a direct impact of such rapid PCR assays is that they should allow for a faster establishment of effective antibiotic therapy and a reduction of empirical treatments with broad-spectrum antibiotics, which are associated with high costs and toxicity. Utilization of the multiplex PCR technology in the clinical laboratory in combination with the recommended NCCLS guidelines would better enable physicians to prescribe appropriate antibiotic therapy, leading to therapeutic success, more prudent antibiotic usage, and conditions less conducive to staphylococcal resistance selection (5).

ACKNOWLEDGMENTS

We thank Louise Côté, director of the Microbiology Laboratory of CHUL, for free access to the laboratory and for providing the staphy-

lococcal isolates. We also thank Louise Jetté (Laboratoire de Santé Publique du Québec), Pierre Auclair (Laval Hospital), Donald E. Low (Mount Sinai Hospital), Wang Fu (Huashan Hospital), and Nevine El Solh (Institut Pasteur) for providing staphylococcal strains. We also thank Ann Huletsky and Maurice Boissinot for helpful suggestions.

F. Martineau has a scholarship from le Fonds de Recherche en Santé du Québec. M. Ouellette is an MRC Scientist and a Burroughs-Wellcome Fund New Investigator in molecular parasitology. This research project was supported by grant PA-15586 from the Medical Research Council of Canada and Infectio Diagnostic (IDI), Inc., Sainte-Foy, Québec, Canada.

REFERENCES

1. Archer, G. L., D. M. Niemeyer, J. A. Thanassi, and M. J. Pucci. 1994. Dissemination among staphylococci of DNA sequences associated with methicillin resistance. *Antimicrob. Agents Chemother.* **38**:447-454.
2. Baron, E. J. 1995. Genetic aspects of methicillin resistance in *Staphylococcus aureus* and methods used for its detection in clinical laboratories in the United States. *J. Chemother.* **7**(Suppl. 3):87-92.
3. Barski, P., L. Piechowicz, J. Galinski, and J. Kur. 1996. Rapid assay for detection of methicillin-resistant *Staphylococcus aureus* using multiplex PCR. *Mol. Cell. Probes* **10**:471-475.
4. Bergeron, M. G., and M. Ouellette. 1995. Diagnosing bacterial infectious diseases in one hour: an essential upcoming revolution. *Infection* **23**:69-72.
5. Bergeron, M. G., and M. Ouellette. 1998. Preventing antibiotic resistance through rapid genotypic identification of bacteria and of their antibiotic resistance genes in the clinical microbiology laboratory. *J. Clin. Microbiol.* **36**:2169-2172.
6. Brakstad, O. G., J. A. Maeland, and Y. Tveten. 1993. Multiplex polymerase chain reaction for detection of genes for *Staphylococcus aureus* thermonuclease and methicillin resistance and correlation with oxacillin resistance. *APMIS* **101**:681-688.
7. Brumfitt, W., and J. Hamilton-Miller. 1989. Methicillin-resistant *Staphylococcus aureus*. *N. Engl. J. Med.* **320**:1188-1196.
8. Busch-Sorensen, C., N. Frimodt-Moller, G. H. Miller, and F. Espersen. 1996. Aminoglycoside resistance among Danish blood culture isolates of coagulase-negative staphylococci. *APMIS* **104**:873-880.
9. Carlton, B. C., and B. J. Brown. 1981. Antibiotic resistance (gradient plate), p. 229-230. *In* P. Gerhardt, R. Murray, R. Costillow, E. Nester, W. Wood, N. Krieg, and G. Briggs Phillips (ed.), *Manual of methods for general bacteriology*. American Society for Microbiology, Washington, D.C.
10. Catchpole, L., C. Thomas, A. Davies, and K. G. H. Dykes. 1988. The nucleotide sequence of *Staphylococcus aureus* plasmid pT48 conferring inducible resistance and comparison with similar plasmids expressing constitutive resistance. *J. Gen. Microbiol.* **134**:697-709.
11. Chambers, H. F. 1997. Methicillin resistance in staphylococci: molecular and biochemical basis and clinical implications. *Clin. Microbiol. Rev.* **10**:781-791.
12. Chomarar, M., D. Espinouse, and J. P. Flandrois. 1991. Coagulase-negative staphylococci emerging during teicoplanin therapy and problems in the determination of their sensitivity. *J. Antimicrob. Chemother.* **27**:475-480.
13. Clancy, J., F. Dib-Hajj, P. J. W., and W. Yuan. 1997. Cloning and characterization of a novel macrolide efflux gene, *mreA*, from *Streptococcus agalactiae*. *Antimicrob. Agents Chemother.* **41**:2719-2723.
14. Cloney, L., C. Marlowe, A. Wong, R. Chow, and R. Bryan. 1999. Rapid detection of *mecA* in methicillin resistant *Staphylococcus aureus* using cycling probe technology. *Mol. Cell. Probes* **13**:191-197.
15. de Lencastre, H., B. L. de Jonge, P. R. Matthews, and A. Tomasz. 1994. Molecular aspects of methicillin resistance in *Staphylococcus aureus*. *J. Antimicrob. Chemother.* **33**:7-24.
16. Dornbusch, K., G. H. Miller, R. S. Hare, and K. J. Shaw. 1990. Resistance to aminoglycoside antibiotics in gram-negative bacilli and staphylococci isolated from blood. Report from a European collaborative study. The ESGAR study group (European Study Group on Antibiotic Resistance). *J. Antimicrob. Chemother.* **26**:131-144.
17. Dubin, D. T., J. E. Fitzgibbon, M. D. Nahvi, and J. F. John. 1999. Topoisomerases sequences of coagulase-negative staphylococcal isolates resistant to ciprofloxacin or trovafloxacin. *Antimicrob. Agents Chemother.* **43**:1631-1637.
18. Eady, E. A., J. I. Ross, J. L. Tipper, C. E. Walters, J. H. Cove, and W. C. Noble. 1993. Distribution of genes encoding erythromycin ribosomal methylases and an erythromycin efflux pump in epidemiologically distinct groups of staphylococci. *J. Antimicrob. Chemother.* **31**:211-217.
19. Geha, D. L., J. R. Uhl, C. A. Gustafarro, and D. H. Persing. 1994. Multiplex detection for identification of methicillin-resistant staphylococci in the clinical laboratory. *J. Clin. Microbiol.* **32**:1768-1772.
20. Hodel-Christian, S. L., and B. E. Murray. 1991. Characterization of the gentamicin resistance transposon Tn5281 from *Enterococcus faecalis* and comparison to staphylococcal transposons Tn4001 and Tn4031. *Antimicrob. Agents Chemother.* **35**:1147-1152.
21. Huang, M. B., T. E. Gay, C. N. Baker, S. N. Banerjee, and F. C. Tenover. 1993. Two percent sodium chloride is required for susceptibility testing of

- staphylococci with oxacillin when using agar-based dilution methods. *J. Clin. Microbiol.* **31**:2683–2688.
22. Hürliemann-Dalel, R. L., C. Ryffel, F. H. Kayser, and B. Berger-Bächli. 1992. Survey of the methicillin resistance-associated genes *mecA*, *mecR1-mecI*, and *femA-femB* in clinical isolates of methicillin-resistant *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **36**:2617–2621.
 23. Jenssen, W. D., S. Thakker Varia, D. T. Dubin, and M. P. Weinstein. 1987. Prevalence of macrolides-lincosamides-streptogramin B resistance and *erm* gene classes among clinical strains of staphylococci and streptococci. *Antimicrob. Agents Chemother.* **31**:883–888.
 24. Kaatz, G. W., S. M. Seo, N. J. Dorman, and S. A. Lerner. 1990. Emergence of teicoplanin resistance during therapy of *Staphylococcus aureus* endocarditis. *J. Infect. Dis.* **162**:103–108.
 25. Khan, S. A., and R. P. Novick. 1980. Terminal nucleotide sequences of Tn551, a transposon specifying erythromycin resistance in *Staphylococcus aureus*: homology with Tn3. *Plasmid* **4**:148–154.
 26. Kloos, W. E., and T. L. Bannerman. 1994. Update on clinical significance of coagulase-negative staphylococci. *Clin. Microbiol. Rev.* **7**:117–140.
 27. Kobayashi, N., K. Taniguchi, K. Kojima, S. Urasawa, N. Uehara, Y. Omizu, Y. Kishi, A. Yagihashi, I. Kurokawa, and N. Watanabe. 1996. Genomic diversity of *mec* regulator genes in methicillin-resistant *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Epidemiol. Infect.* **117**:289–295.
 28. Kolbert, C. P., J. Arruda, P. Varga-Delmore, X. Zheng, M. Lewis, J. Kolberg, and D. H. Persing. 1998. Branched-DNA assay for detection of the *mecA* gene in oxacillin-resistant and oxacillin-sensitive staphylococci. *J. Clin. Microbiol.* **36**:2640–2644.
 29. Kuwahara-Arai, K., N. Kondo, and S. Hori. 1996. Suppression of methicillin resistance in a *mecA*-containing pre-methicillin-resistant *Staphylococcus aureus* strain is caused by the *mecI*-mediated repression of PBP2' production. *Antimicrob. Agents Chemother.* **40**:2680–2685.
 30. Lampson, B. C., and J. T. Parisi. 1986. Naturally occurring *Staphylococcus epidermidis* plasmid expressing constitutive macrolides-lincosamides-streptogramin B resistance contains a deleted attenuator. *J. Bacteriol.* **166**:479–483.
 31. Lovering, A. M., M. J. Bywater, H. A. Holt, H. M. Champion, and D. S. Reeves. 1988. Resistance of bacterial pathogens to four aminoglycosides and six other antibacterials and prevalence of aminoglycosides modifying enzymes. *J. Antimicrob. Chemother.* **22**:823–839.
 32. Madsen, O. R., J. S. Lorentzen, N. Frimodt-Møller, I. Mortensen, and V. T. Rosdahl. 1991. Mechanisms of aminoglycoside resistance in Danish *Staphylococcus aureus* strains during the years 1979–1987. *APMIS* **99**:537–540.
 33. Manquat, G., J. Croizé, J. P. Stahl, M. Meyran, P. Hirtz, and M. Micoud. 1992. Failure of teicoplanin treatment associated with an increase in MIC during therapy of *Staphylococcus aureus* septicaemia. *J. Antimicrob. Chemother.* **29**:731–732.
 34. Martineau, F., F. J. Picard, P. H. Roy, M. Ouellette, and M. G. Bergeron. 1998. Species-specific and ubiquitous DNA-based assays for rapid identification of *Staphylococcus aureus*. *J. Clin. Microbiol.* **36**:618–623.
 35. Martineau, F., F. J. Picard, P. H. Roy, M. Ouellette, and M. G. Bergeron. 1996. Species-specific and ubiquitous DNA-based assays for rapid identification of *Staphylococcus epidermidis*. *J. Clin. Microbiol.* **34**:2888–2893.
 36. Murakami, K., W. Minamide, K. Wada, E. Nakamura, H. Teraoka, and S. Watanabe. 1991. Identification of methicillin-resistant strains of staphylococci by polymerase chain reaction. *J. Clin. Microbiol.* **29**:2240–2244.
 37. Murphy, E. 1985. Nucleotide sequence of *ermA*, a macrolide-lincosamide-streptogramin B determinant in *Staphylococcus aureus*. *J. Bacteriol.* **162**:633–640.
 38. National Committee for Clinical Laboratory Standards. 1990. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically. Approved standard M7-A2. National Committee for Clinical Laboratory Standards, Villanova, Pa.
 39. National Committee for Clinical Laboratory Standards. 1990. Performance standards for antimicrobial disk susceptibility tests. Approved standard M2-4A. National Committee for Clinical Laboratory Standards, Villanova, Pa.
 40. Niemeyer, D. M., M. J. Pucci, J. A. Thanassi, V. K. Sharma, and G. L. Archer. 1996. Role of *mecA* transcriptional regulation in the phenotypic expression of methicillin resistance in *Staphylococcus aureus*. *J. Bacteriol.* **178**:5464–5471.
 41. Petersson, A. C., H. Miorner, and C. Kamme. 1996. Identification of *mecA*-related oxacillin resistance in staphylococci by the E test and the broth microdilution method. *J. Antimicrob. Chemother.* **37**:445–456.
 42. Pfaller, M. A., and L. A. Herwaldt. 1988. Laboratory, clinical and epidemiological aspects of coagulase-negative staphylococci. *Clin. Microbiol. Rev.* **1**:281–299.
 43. Pfaller, M. A., R. N. Jones, G. V. Doern, K. Kugler, and The SENTRY Participants Group. 1998. Bacterial pathogens isolated from patients with bloodstream infection: frequencies of occurrence and antimicrobial susceptibility patterns from the SENTRY antimicrobial surveillance program (United States and Canada, 1997). *Antimicrob. Agents Chemother.* **42**:1762–1770.
 44. Predari, S. C., M. Ligozzi, and R. Fontana. 1991. Genotypic identification of methicillin-resistant coagulase-negative staphylococci by polymerase chain reaction. *Antimicrob. Agents Chemother.* **35**:2568–2573.
 45. Price, K. E., P. A. Kresel, L. A. Farchione, S. B. Siskin, and S. A. Karpow. 1981. Epidemiological studies of aminoglycoside resistance in the U.S.A. *J. Antimicrob. Chemother.* **8**(Suppl. A):89–105.
 46. Qadri, S. M. H., Y. Ueno, H. Imambaccus, and E. Almodovar. 1994. Rapid detection of methicillin-resistant *Staphylococcus aureus* by Crystal MRSA ID System. *J. Clin. Microbiol.* **32**:1830–1832.
 47. Ross, J. I., E. A. Eady, J. H. Cove, and S. Baumberg. 1995. Identification of a chromosomally encoded ABC-transport system with which the staphylococcal erythromycin exporter *MsrA* may interact. *Gene* **153**:93–98.
 48. Ross, J. I., E. A. Eady, J. H. Cove, W. J. Cunliffe, S. Baumberg, and J. C. Wootton. 1990. Inducible erythromycin resistance in staphylococci is encoded by a member of the ATP-binding transport super-gene family. *Mol. Microbiol.* **4**:1207–1214.
 49. Rouch, D. A., M. E. Byrne, Y. C. Kong, and R. A. Skurray. 1987. The *aacA-aphD* gentamicin and kanamycin resistance determinant of Tn4001 from *Staphylococcus aureus*: expression and nucleotide sequence analysis. *J. Gen. Microbiol.* **133**:3039–3052.
 50. Rupp, M. E., and G. L. Archer. 1994. Coagulase-negative staphylococci: pathogens associated with medical progress. *Clin. Infect. Dis.* **19**:231–245.
 51. Salisbury, S. M., L. M. Sabatini, and C. A. Spiegel. 1997. Identification of methicillin-resistant staphylococci by multiplex polymerase chain reaction. *Am. J. Clin. Pathol.* **107**:368–373.
 52. Seppälä, H., M. Skurnik, H. Soini, M. Roberts, and P. Huovinen. 1998. A novel erythromycin resistance methylase gene (*ermTR*) in *Streptococcus pyogenes*. *Antimicrob. Agents Chemother.* **42**:257–262.
 53. Shaw, K. J., P. N. Rather, R. S. Hare, and G. H. Miller. 1993. Molecular genetics of aminoglycoside resistance genes and familial relationships of the aminoglycoside-modifying enzymes. *Microbiol. Rev.* **57**:138–163.
 54. Sheagren, J. N. 1984. *Staphylococcus aureus*: the persistent pathogen. *N. Engl. J. Med.* **310**:1368–1442.
 55. Sutcliffe, J., T. Grebe, A. Taitkramradt, and L. Wondrack. 1996. Detection of erythromycin-resistant determinants by PCR. *Antimicrob. Agents Chemother.* **40**:2562–2566.
 56. Suzuki, E., K. Kuwahara-Arai, and J. F. Richardson. 1993. Distribution of *mec* regulator genes in methicillin-resistant *Staphylococcus* clinical strains. *Antimicrob. Agents Chemother.* **37**:1219–1226.
 57. Szybalski, W. 1952. Microbial selection. I. Gradient plate technique for study of bacterial resistance. *Science* **116**:46–48.
 58. Thakker-Varia, S., W. D. Jenssen, L. Moon-McDermott, M. P. Weinstein, and D. T. Dubin. 1987. Molecular epidemiology of macrolides-lincosamides-streptogramin B resistance in *Staphylococcus aureus* and coagulase-negative staphylococci. *Antimicrob. Agents Chemother.* **31**:735–743.
 59. Únal, S., J. Hoskins, J. E. Flokowsitch, C. Y. E. Wu, D. A. Preston, and P. L. Skatrud. 1992. Detection of methicillin-resistant staphylococci by using the polymerase chain reaction. *J. Clin. Microbiol.* **30**:1685–1691.
 60. Únal, S., K. Werner, P. De Girolami, F. Barsant, and G. Eliopoulos. 1994. Comparison of tests for detection of methicillin-resistant *Staphylococcus aureus* in a clinical microbiology laboratory. *Antimicrob. Agents Chemother.* **38**:345–347.
 61. Vanhoof, R., C. Godard, J. Content, H. J. Nyssen, and E. Hannecart-Pokorni. 1994. Detection by polymerase chain reaction of genes encoding aminoglycoside-modifying enzymes in methicillin-resistant *Staphylococcus aureus* isolates of epidemic phage types. *J. Med. Microbiol.* **41**:282–290.
 62. Vannuffel, P., J. Gigi, H. Ezzedine, B. Vandercam, M. Delmee, G. Wauters, and J.-L. Gala. 1995. Specific detection of methicillin-resistant *Staphylococcus* species by multiplex PCR. *J. Clin. Microbiol.* **33**:2864–2867.
 63. Weisblum, B. 1995. Erythromycin resistance by ribosome modification. *Antimicrob. Agents Chemother.* **39**:577–585.
 64. Westh, H., D. M. Hougaard, J. Vuust, and V. T. Rosdahl. 1995. *Erm* genes in erythromycin-resistant *Staphylococcus aureus* and coagulase-negative staphylococci. *APMIS* **103**:225–232.
 65. Westh, H., D. M. Hougaard, J. Vuust, and V. T. Rosdahl. 1995. Prevalence of *erm* gene classes in erythromycin-resistant *Staphylococcus aureus* strains isolated between 1959 and 1988. *Antimicrob. Agents Chemother.* **39**:369–373.
 66. Williams, J. G., A. R. Kubelik, K. J. Livak, J. A. Rafalski, and S. V. Tingey. 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res.* **25**:6531–6535.