

# Antipneumococcal Activities of GAR-936 (a New Glycylcycline) Compared to Those of Nine Other Agents against Penicillin-Susceptible and -Resistant Pneumococci

DIANNE B. HOELLMAN,<sup>1</sup> GLENN A. PANKUCH,<sup>1</sup> MICHAEL R. JACOBS,<sup>2</sup>  
AND PETER C. APPELBAUM<sup>1\*</sup>

*Department of Pathology, Hershey Medical Center, Hershey, Pennsylvania 17033,<sup>1</sup> and Department of Pathology, Case Western Reserve University, Cleveland, Ohio 44106<sup>2</sup>*

Received 25 June 1999/Returned for modification 26 November 1999/Accepted 16 January 2000

**GAR-936, a new glycylcycline, had lower MICs ( $\leq 0.016$  to  $0.125$   $\mu\text{g/ml}$ ) for 201 penicillin- and tetracycline-susceptible and -resistant pneumococcal strains than tetracycline ( $\leq 0.06$  to  $128$   $\mu\text{g/ml}$ ), minocycline ( $\leq 0.06$  to  $16.0$   $\mu\text{g/ml}$ ), or doxycycline ( $\leq 0.06$  to  $32.0$   $\mu\text{g/ml}$ ). GAR-936 was also bactericidal against 11 of 12 strains tested at the MIC after 24 h, with significant kill rates at earlier time points.**

The incidence of pneumococcal resistance to penicillin G and other  $\beta$ -lactam and non- $\beta$ -lactam compounds has increased worldwide at an alarming rate (1, 6, 7). In the United States, a recent survey of 1,476 strains showed that 49.6% were penicillin susceptible, 17.9% were of intermediate resistance (penicillin intermediate), and 32.5% were penicillin resistant. Macrolide MICs rose with those of penicillin G, with an overall macrolide resistance level of 33% (8).

Resistance to conventional oral tetracyclines (including minocycline and doxycycline) occurs with increased frequency among penicillin-intermediate and -resistant pneumococci, and genes coding for macrolide and tetracycline resistance are carried on the same transposon (6, 7). GAR-936 is a new broad-spectrum parenteral glycylcycline. Preliminary studies have shown that this drug is active against penicillin-susceptible and -resistant pneumococci (13, 15).

This study compares the activity of GAR-936 against 201 penicillin-susceptible, -intermediate, or -resistant pneumococcal strains with those of three other tetracyclines, four  $\beta$ -lactams, clarithromycin, and vancomycin. Additionally, the activities of these compounds against 12 penicillin-susceptible and -resistant strains of pneumococci were determined by the broth MIC and time-kill methods.

For determination of MICs by agar dilution, 51 penicillin-susceptible, 72 penicillin-intermediate, and 78 penicillin-resistant strains of pneumococci were utilized. For time-kill studies, four penicillin-susceptible, four penicillin-intermediate, and four penicillin-resistant strains were tested; of these, five were tetracycline susceptible (MIC,  $\leq 2.0$   $\mu\text{g/ml}$ ), 1 was tetracycline intermediate (MIC,  $4.0$   $\mu\text{g/ml}$ ), and six were tetracycline resistant (MIC,  $\geq 8.0$   $\mu\text{g/ml}$ ).

The agar dilution method used in this study to determine MICs of various antimicrobial agents for 201 pneumococcal strains has been used routinely for many years by our group (6, 7); Mueller-Hinton agar (BBL Microbiology Systems, Cockeysville, Md.) supplemented with 5% sheep blood was employed. Microbroth MIC determinations for 12 strains were

performed in accordance with the protocol recommended by the National Committee for Clinical Laboratory Standards (9). Standard quality control strains, including *Streptococcus pneumoniae* ATCC 49619 (9), were included in each run of agar and broth dilution MIC determinations. All MIC studies were performed in air. All strains, recently isolated from stocks stored at  $-70^\circ\text{C}$  and subcultured twice, grew in air and did not require added  $\text{CO}_2$  for growth. The susceptibility patterns of these strains are representative of those commonly encountered in the United States (8).

For time-kill studies, methods described previously were used (11, 12). The final inoculum was  $5 \times 10^5$  to  $5 \times 10^6$  CFU/ml. Growth controls with inoculum but no antibiotic were included with each experiment (11, 12). Viable-cell counts of antibiotic-containing suspensions were performed according to standard methods (11, 12). Colony counts were performed on plates yielding 30 to 300 colonies. The lower limit of sensitivity of colony counts was 300 CFU/ml. As mentioned above, all incubations were in ambient air without added  $\text{CO}_2$ .

Time-kill assays were analyzed by determining the number of strains which yielded a  $\Delta\log_{10}$  CFU/ml of  $-1$ ,  $-2$ , and  $-3$  at 0, 4, 6, 12, and 24 h compared to counts at time zero. Antimicrobial agents were considered bactericidal at the lowest concentration that reduced the original inoculum by  $\geq 3 \log_{10}$  CFU/ml (99.9%) at each of the time periods, and they were considered bacteriostatic if the inoculum was reduced by 0 to  $3 \log_{10}$  CFU/ml. With the sensitivity threshold and inocula used in these studies, no problems were encountered in delineating 99.9% killing when it occurred. The problem of bacterial carryover was addressed by dilution, as described previously (11, 12). For tetracycline and clarithromycin time-kill testing, only strains with MICs of  $<8.0$  and  $<4.0$   $\mu\text{g/ml}$ , respectively, were tested. In all cases, growth controls grew satisfactorily, without spontaneous autolysis.

The results of agar dilution MIC testing are presented in Table 1. GAR-936 had lower MICs ( $\leq 0.016$  to  $0.125$   $\mu\text{g/ml}$ ) than tetracycline ( $\leq 0.06$  to  $128$   $\mu\text{g/ml}$ ), minocycline ( $\leq 0.06$  to  $16.0$   $\mu\text{g/ml}$ ), and doxycycline ( $\leq 0.06$  to  $32.0$   $\mu\text{g/ml}$ ) against all strains. MICs of all  $\beta$ -lactams rose with those of penicillin G, with imipenem being the most active, followed by meropenem, ceftriaxone, and amoxicillin. Macrolide and tetracycline resistance occurred mainly in penicillin-intermediate

\* Corresponding author. Mailing address: Department of Pathology, Hershey Medical Center, Hershey, PA 17033. Phone: (717) 531-5113. Fax: (717) 531-7953. E-mail: pappelbaum@psghs.edu.

TABLE 1. Agar dilution MICs for 201 pneumococcal isolates

Drug	Penicillin susceptibility of strain <sup>a</sup>	MIC (μg/ml)		
		Range	MIC <sub>50</sub> <sup>b</sup>	MIC <sub>90</sub> <sup>c</sup>
Penicillin	S	≤0.008–0.06	≤0.016	0.03
	I	0.125–1.0	0.25	1.0
	R	2.0–>8.0	2.0	4.0
GAR-936	S	≤0.016–0.125	0.03	0.125
	I	≤0.016–0.125	0.03	0.06
	R	≤0.016–0.125	0.06	0.125
Tetracycline	S	≤0.06–64.0	0.25	32.0
	I	≤0.06–64.0	1.0	64.0
	R	≤0.25–128.0	32.0	64.0
Minocycline	S	≤0.06–16.0	0.06	8.0
	I	≤0.06–16.0	0.25	16.0
	R	≤0.06–16.0	4.0	16.0
Doxycycline	S	0.03–16.0	0.125	8.0
	I	≤0.06–32.0	0.5	8.0
	R	0.125–16.0	4.0	8.0
Amoxicillin	S	≤0.008–0.25	0.016	0.03
	I	≤0.016–2.0	0.125	1.0
	R	0.5–4.0	2.0	2.0
Ceftriaxone	S	≤0.008–1.0	0.016	0.03
	I	≤0.016–2.0	0.25	1.0
	R	0.25–8.0	1.0	2.0
Clarithromycin	S	≤0.016–>128.0	0.03	8.0
	I	≤0.016–>128.0	0.06	>128.0
	R	≤0.016–>128.0	1.0	>128.0
Imipenem	S	≤0.008–0.25	≤0.008	0.016
	I	≤0.008–0.25	0.06	0.25
	R	0.03–0.5	0.25	0.5
Meropenem	S	≤0.008–0.5	0.016	0.03
	I	≤0.016–0.5	0.125	0.5
	R	0.125–2.0	0.5	1.0
Vancomycin	S	≤0.06–0.5	0.25	0.25
	I	≤0.06–0.5	0.25	0.25
	R	0.125–0.5	0.25	0.25

<sup>a</sup> S, penicillin susceptible; I, penicillin intermediate; R, penicillin resistant.  
<sup>b</sup> MIC<sub>50</sub>, MIC at which 50% of the isolates tested were inhibited.  
<sup>c</sup> MIC<sub>90</sub>, MIC at which 90% of the isolates tested were inhibited.

and -resistant strains, and all strains were susceptible to vancomycin.

Microbroth MICs of the 12 strains tested by the time-kill method were either identical to or differed by one dilution from those obtained by agar dilution (Table 2). Time-kill results are presented in Table 3. GAR-936 was bactericidal at the MIC after 24 h for 11 of the 12 strains tested; the remaining strain (for which the GAR-936 MIC was 0.06 μg/ml) was killed

at eight times the MIC (0.5 μg/ml). GAR-936 showed significant kill rates at earlier time periods: at 12 h, 90% killing of all 12 strains occurred at twice the MIC, while after 6 h, 90% killing of all strains occurred at four times the MIC. The killing rates of other tetracyclines for the six susceptible strains were slightly lower than those of GAR-936. The bactericidal activities of β-lactams were similar, with uniform bactericidal activity being exhibited at twice the MIC for amoxicillin, ceftriaxone, and imipenem and at eight times the MIC for meropenem after 24 h. Clarithromycin was bactericidal against six of seven strains with clarithromycin MICs of <4.0 μg/ml at twice the MIC after 24 h; vancomycin was bactericidal against 11 of 12 strains at twice the MIC at the same time point. The kill kinetics of GAR-936 against a tetracycline-resistant strain is depicted in Fig. 1.

The glycylicyclines are semisynthetic tetracyclines exhibiting a broad spectrum of activity compared to conventional tetracyclines, with activity against gram-positive and -negative aerobes, anaerobes, chlamydiae, and mycoplasmas (2–5, 10, 14, 16, 17). GAR-936 is a newly described 9-*t*-butylglycylamido derivative of minocycline. This compound possesses activity against isolates possessing the two major determinants of tetracycline resistance, ribosomal protection and efflux (13, 15).

In the present study, MICs of GAR-936 against all pneumococci, irrespective of their susceptibility to β-lactams, macrolides, or other tetracyclines, ranged between ≤0.016 and 0.125 μg/ml. These results are similar to those obtained by Petersen et al (13). Bactericidal activity was also found after 24 h for all 12 strains tested at ≤0.5 μg/ml. At the MIC, after 24 h, GAR-936 showed better kill kinetics, relative to the MIC, than other tetracyclines and was bactericidal for more strains than were the β-lactams tested. It should be emphasized, however, that the relatively small number of strains tested makes interpretation of relatively small differences in kill kinetics problematic. More strains must be tested before such conclusions can be considered valid. Although tetracyclines are traditionally thought to be bacteriostatic agents, the present study shows that GAR-936 (and other tetracyclines, relative to their MICs) may be bactericidal after 24 h.

Of the other tetracyclines tested, minocycline and doxycycline had lower MICs than tetracycline; the significance of this phenomenon is unknown at this time. The MICs of and time-kills results for the other compounds tested are similar to those reported by others (7, 11, 12).

Although the results of toxicologic and pharmacokinetic studies are awaited, GAR-936 appears to be a promising

TABLE 2. Microbroth MICs for 12 strains tested by the time-kill method

Drug	MIC (μg/ml) for strain <sup>a</sup> :											
	1(S)	2(S)	3(S)	4(S)	5(I)	6(I)	7(I)	8(I)	9(R)	10(R)	11(R)	12(R)
Penicillin	0.06	0.016	0.03	0.03	0.5	0.5	0.25	0.25	4.0	2.0	4.0	4.0
GAR-936	0.06	0.03	0.03	0.06	0.125	0.06	0.03	0.03	0.03	0.03	0.06	0.06
Tetracycline	64.0	0.5	64.0	0.5	4.0	64.0	0.125	0.25	32.0	32.0	1.0	32.0
Minocycline	16.0	0.125	8.0	0.125	0.5	64.0	0.06	0.06	16.0	8.0	0.25	4.0
Doxycycline	8.0	0.25	16.0	0.25	1.0	16.0	0.06	0.125	8.0	8.0	1.0	8.0
Amoxicillin	0.03	0.03	0.016	0.016	0.06	0.125	0.125	0.06	1.0	1.0	2.0	2.0
Ceftriaxone	0.06	0.016	0.016	0.016	0.125	1.0	0.06	0.06	2.0	2.0	1.0	2.0
Clarithromycin	>64.0	0.03	0.016	0.03	>64.0	0.008	0.016	0.03	>64.0	>64.0	0.03	>64.0
Imipenem	0.008	0.008	0.008	0.008	0.03	0.06	0.03	0.03	0.25	0.25	0.5	0.25
Meropenem	0.016	0.016	0.016	0.008	0.06	0.125	0.125	0.06	1.0	0.5	1.0	1.0
Vancomycin	0.25	0.25	0.5	0.5	0.25	0.5	0.5	0.125	0.5	0.25	0.25	0.25

<sup>a</sup> S, penicillin susceptible; I, penicillin intermediate; R, penicillin resistant.

TABLE 3. Results of time-kill assays of 12 pneumococcal strains

Drug and concn	No. of strains killed by time point, with $\Delta\log_{10}$ CFU/ml (from time zero value) of:											
	3 h			6 h			12 h			24 h		
	-1	-2	-3	-1	-2	-3	-1	-2	-3	-1	-2	-3
<b>GAR-936</b>												
8× MIC	7	3	1	12	6	4	12	11	7	12	12	12
4× MIC	6	2	1	12	6	4	12	10	6	12	12	11
2× MIC	5	2	1	11	5	3	12	9	6	12	11	11
MIC	4	1	1	7	5	2	10	8	4	12	11	11
0.5× MIC	0	0	0	3	0	0	3	2	1	5	3	2
<b>Tetracycline<sup>a</sup></b>												
8× MIC	3	1	0	5	3	3	5	4	3	6	6	5
4× MIC	2	0	0	4	3	0	5	4	2	6	6	5
2× MIC	2	0	0	3	2	0	5	2	2	6	5	5
MIC	1	0	0	2	0	0	3	2	0	3	3	1
0.5× MIC	0	0	0	0	0	0	1	0	0	0	0	0
<b>Minocycline<sup>a</sup></b>												
8× MIC	4	2	1	4	4	2	5	4	4	6	5	5
4× MIC	3	1	1	4	3	2	5	4	3	6	5	5
2× MIC	2	1	0	4	2	2	5	4	3	6	5	4
MIC	2	0	0	3	2	0	5	3	2	6	4	3
0.5× MIC	1	0	0	1	0	0	0	0	0	0	0	0
<b>Doxycycline<sup>a</sup></b>												
8× MIC	4	2	0	5	3	3	5	5	3	6	5	5
4× MIC	3	1	0	5	3	0	5	5	3	6	5	5
2× MIC	1	0	0	4	1	0	5	4	1	6	5	4
MIC	1	0	0	2	1	0	2	1	1	2	2	2
0.5× MIC	0	0	0	1	0	0	1	0	0	0	0	0
<b>Amoxicillin</b>												
8× MIC	10	4	0	12	10	6	12	12	10	12	12	12
4× MIC	9	2	0	11	9	3	12	12	8	12	12	12
2× MIC	8	3	0	11	8	2	12	12	7	12	12	12
MIC	5	2	0	11	4	2	10	5	3	10	8	8
0.5× MIC	1	0	0	2	1	0	1	1	0	2	1	1
<b>Ceftriaxone</b>												
8× MIC	8	0	0	12	10	0	12	12	10	12	12	12
4× MIC	8	1	0	12	9	1	12	11	8	12	12	12
2× MIC	8	0	0	10	7	0	12	10	8	12	12	12
MIC	5	1	0	9	6	0	10	7	1	10	10	8
0.5× MIC	2	0	0	2	0	0	1	1	0	0	0	0
<b>Clarithromycin<sup>b</sup></b>												
8× MIC	4	2	0	7	4	0	7	4	4	7	7	6
4× MIC	3	1	0	6	3	0	7	4	4	7	6	6
2× MIC	3	1	0	4	3	1	7	4	4	7	6	6
MIC	2	1	0	4	2	0	6	4	3	7	6	4
0.5× MIC	1	0	0	2	0	0	2	1	0	2	1	1
<b>Imipenem</b>												
8× MIC	12	4	0	12	10	2	12	11	10	12	12	12
4× MIC	11	5	0	12	9	2	12	11	10	12	12	12
2× MIC	11	3	0	12	9	4	12	11	10	12	12	12
MIC	10	2	0	12	8	2	12	10	8	11	11	9
0.5× MIC	8	0	0	6	3	0	4	2	0	3	3	2
<b>Meropenem</b>												
8× MIC	11	3	1	12	5	2	12	12	9	12	12	12
4× MIC	9	2	0	12	4	2	12	12	7	12	12	11
2× MIC	6	4	0	12	7	3	12	11	8	12	12	11
MIC	6	2	1	11	5	2	11	9	5	10	9	8
0.5× MIC	1	0	0	2	0	0	1	0	0	2	1	1
<b>Vancomycin</b>												
8× MIC	7	1	0	12	6	1	12	11	9	12	12	12
4× MIC	7	1	0	12	6	2	12	10	9	12	12	11
2× MIC	7	1	0	12	6	2	12	10	8	12	12	11
MIC	6	1	0	12	4	0	11	9	5	10	9	7
0.5× MIC	0	0	0	2	0	0	2	0	0	2	2	0

<sup>a</sup> Only strains with tetracycline MICs of <8.0 µg/ml were tested.<sup>b</sup> Only strains with clarithromycin MICs of <4.0 µg/ml were tested.




FIG. 1. Graphical depiction of GAR-936 kill kinetics versus a tetracycline-resistant (MIC, 64.0  $\mu\text{g/ml}$ ) strain of *S. pneumoniae*.

new agent for treatment of infections caused by drug-resistant pneumococci.

This study was supported by a grant from Wyeth-Ayerst Research Laboratories, Pearl River, N.Y.

#### REFERENCES

- Appelbaum, P. C. 1992. Antimicrobial resistance in *Streptococcus pneumoniae*—an overview. *Clin. Infect. Dis.* **15**:77–83.
- Bergeron, J., M. Ammirati, D. Danley, L. James, M. Norcia, J. Retsema, C. A. Strick, W.-G. Su, J. Sutcliffe, and L. Wondrack. 1996. Glycylcyclines bind to the high-affinity tetracycline ribosomal binding site and evade Tet(M)- and Tet(O)-mediated ribosomal protection. *Antimicrob. Agents Chemother.* **40**:2226–2228.
- Eliopoulos, G. M., C. B. Wennersten, G. Cole, and R. C. Moellering. 1994. In vitro activities of two glycylcyclines against gram-positive bacteria. *Antimicrob. Agents Chemother.* **38**:534–541.
- Goldstein, F. W., M. D. Kitzis, and J. F. Acar. 1994. *N,N*-Dimethylglycyl-amido derivative of minocycline and 6-demethyl-6-desoxytetracycline, two new glycylcyclines highly effective against tetracycline-resistant gram-positive cocci. *Antimicrob. Agents Chemother.* **38**:2218–2220.
- Hamilton-Miller, J. M. T., and S. Shah. 1996. Activity of glycylcyclines CL 329998 and CL 331002 against minocycline-resistant and other strains of methicillin-resistant *Staphylococcus aureus*. *J. Antimicrob. Chemother.* **37**:1171–1175.
- Jacobs, M. R. 1992. Treatment and diagnosis of infections caused by drug-resistant *Streptococcus pneumoniae*. *Clin. Infect. Dis.* **15**:119–127.
- Jacobs, M. R., and P. C. Appelbaum. 1995. Antibiotic-resistant pneumococci. *Rev. Med. Microbiol.* **6**:77–93.
- Jacobs, M. R., S. Bajaksouzian, A. Zilles, G. Lin, G. A. Pankuch, and P. C. Appelbaum. 1999. Susceptibilities of *Streptococcus pneumoniae* and *Haemophilus influenzae* to 10 oral antimicrobial agents based on pharmacodynamic parameters: 1997 U.S. surveillance study. *Antimicrob. Agents Chemother.* **43**:1901–1908.
- National Committee for Clinical Laboratory Standards. 1997. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically, 3rd ed. Approved standard M7-A4. National Committee for Clinical Laboratory Standards, Villanova, Pa.
- Nord, C. E., A. Lindmark, and I. Persson. 1993. In vitro activity of DMG-Mino and DMG-DM Dot, two new glycylcyclines, against anaerobic bacteria. *Eur. J. Clin. Microbiol. Infect. Dis.* **12**:784–786.
- Pankuch, G. A., M. R. Jacobs, and P. C. Appelbaum. 1994. Study of comparative antipneumococcal activities of penicillin G, RP 59500, erythromycin, sparflaxacin, ciprofloxacin, and vancomycin by using time-kill methodology. *Antimicrob. Agents Chemother.* **38**:2065–2072.
- Pankuch, G. A., C. Lichtenberger, M. R. Jacobs, and P. C. Appelbaum. 1996. Antipneumococcal activities of RP 59500 (quinupristin-dalfopristin), penicillin G, erythromycin, and sparflaxacin determined by MIC and rapid time-kill methodologies. *Antimicrob. Agents Chemother.* **40**:1653–1656.
- Petersen, P. J., N. V. Jacobus, W. J. Weiss, P. E. Sum, and R. T. Testa. 1999. In vitro and in vivo antibacterial activities of a novel glycylcycline, the 9-*t*-butylglycylamido derivative of minocycline (GAR-936). *Antimicrob. Agents Chemother.* **43**:738–744.
- Testa, R. T., P. J. Petersen, N. V. Jacobus, P.-E. Sum, V. J. Lee, and F. P. Tally. 1993. In vitro and in vivo antibacterial activities of the glycylcyclines, a new class of semisynthetic tetracyclines. *Antimicrob. Agents Chemother.* **37**:2270–2277.
- Van Ogtrop, M. L., D. Andes, W. A. Craig, and O. Vesga. 1998. In vivo activity of two glycylcyclines (GAR-936 and WAY 152,288) against tetracycline-sensitive and -resistant bacteria, abstr. F-133, p. 266. *In Abstracts of the 38th Interscience Conference on Antimicrobial Agents and Chemotherapy*. American Society for Microbiology, Washington, D.C.
- Weiss, W. J., N. V. Jacobus, P. J. Petersen, and R. T. Testa. 1995. Susceptibility of enterococci, methicillin-resistant *Staphylococcus aureus* and *Streptococcus pneumoniae* to the glycylcyclines. *J. Antimicrob. Chemother.* **36**:225–230.
- Wise, R., and J. M. Andrews. 1994. In vitro activities of two glycylcyclines. *Antimicrob. Agents Chemother.* **38**:1096–1102.