

## Mefloquine and New Related Compounds Target the $F_0$ Complex of the $F_0F_1$ $H^+$ -ATPase of *Streptococcus pneumoniae*

Antonio Javier Martín-Galiano,<sup>1</sup> Begoña Gorgojo,<sup>1</sup> Calvin M. Kunin,<sup>2</sup> and Adela G. de la Campa<sup>1\*</sup>

Unidad de Genética Bacteriana (Consejo Superior de Investigaciones Científicas), Centro Nacional de Microbiología, Instituto de Salud Carlos III, 28220 Majadahonda, Madrid, Spain,<sup>1</sup> and Department of Internal Medicine, The Ohio State University, Columbus, Ohio<sup>2</sup>

Received 26 November 2001/Returned for modification 7 February 2002/Accepted 26 February 2002

**The activities of mefloquine (MFL) and related compounds against previously characterized *Streptococcus pneumoniae* strains carrying defined amino acid substitutions in the *c* subunit of the  $F_0F_1$   $H^+$ -ATPase were studied. In addition, a series of MFL-resistant (MFL<sup>r</sup>) strains were isolated and characterized. A good correlation was observed between inhibition of growth and inhibition of the membrane-associated  $F_0F_1$   $H^+$ -ATPase activity. MFL was about 10-fold more active than optochin and about 200-fold more active than quinine in inhibiting both the growth and the ATPase activities of laboratory pneumococcal strain R6. Mutant strains were inhibited by the different compounds to different degrees, depending on their specific mutations in the *c* subunit. The resistant strains studied had point mutations that changed amino acid residues in either the *c* subunit or the *a* subunit of the  $F_0$  complex. Changes in the *c* subunit were located in one of the two transmembrane  $\alpha$  helices: residues M13, G14, G20, M23, and N24 of helix 1 and residues M44, G47, V48, A49, and V57 of helix 2. Changes in the *a* subunit were also found in either of the transmembrane  $\alpha$  helices, helix 5 or 6: residue L186 of helix 5 and residues W206, F209, and S214 of helix 6. These results suggest that the transmembrane helices of the *c* and *a* subunits interact and that the mutated residues are important for the structure of the  $F_0$  complex and proton translocation.**

The prevalence of *Streptococcus pneumoniae* strains resistant to common antimicrobial agents (10, 26) and the emergence of resistance to new drugs like the fluoroquinolones (4, 20) have highlighted the need for novel antimicrobial agents directed at new targets. The membrane-associated  $F_0F_1$   $H^+$ -ATPase could be such a target, given that the unique sensitivity of *S. pneumoniae* to the amino alcohol antimalarial agents optochin (OPT) and quinine (QIN) resides in the characteristics of its  $F_0$  complex (6, 24). The  $H^+$ -ATPase could be essential for the viability of the pneumococcus, as demonstrated in *Lactococcus lactis*, a genetically related species (17). The primary role of this enzyme in *S. pneumoniae* is to create a proton gradient with the energy provided by ATP hydrolysis and to maintain the intracellular pH via proton extrusion (21), as in other related bacteria (7, 14). However, in bacteria with a respiratory chain such as *Escherichia coli*, the primary role of this enzyme is to synthesize ATP from the proton gradient of the respiratory chain. The bacterial  $F_0F_1$  ATPases consist of cytoplasmic sector  $F_1$  (with  $\alpha$ ,  $\beta$ ,  $\delta$ ,  $\gamma$ , and  $\xi$  subunits), which contains the catalytic sites for ATP hydrolysis, and cytoplasmic membrane sector  $F_0$  (with *a*, *c*, and *b* subunits), which forms the proton channel. Proton transport through  $F_0$  drives the release of ATP on  $F_1$  by long-range conformational changes. The protonation state of an acidic residue of the *c* subunit produces conformational changes that drive the rotation of an oligomeric ring of these subunits. This rotation, in turn, causes the rotation of the attached  $\gamma$  and  $\xi$  subunits, producing conformational changes

in the  $\beta$  subunits (28, 33) that result in ATP synthesis (1). Conversely, hydrolysis of ATP in  $F_1$  generates rotation of  $\gamma$  and  $\xi$ , resulting in the pumping of protons back across the membrane through  $F_0$ . In *S. pneumoniae* (21), as well as in other related bacteria (2, 15–18), the activity of the  $F_0F_1$  ATPase has been seen to increase as the pH of the growth medium decreases. The regulation of this pH-inducible phenotype occurs in *S. pneumoniae* at the level of transcription initiation (21).

Although OPT was used at the beginning of the 20th century for the treatment of pneumococcal infections (23), nowadays its use is restricted to diagnostic purposes due to its high toxicity. However, new related, less toxic compounds have been synthesized (19). Given the special characteristics of the pneumococcal  $F_0F_1$  ATPase and its role in the maintenance of intracellular pH, this enzyme could be considered a target for the design of new antibacterial agents. Moreover, the pneumococcus could also be considered a model system for the study of the putative antimalarial activities of new compounds (24).

In the present study, we describe the molecular basis of the antipneumococcal behavior of mefloquine (MFL) and related compounds. MFL has been found to have bactericidal activities against gram-positive bacteria, including staphylococci, pneumococci, and enterococci, but showed poor activities against gram-negative enteric bacteria and yeasts (19). However, the MFL-related compounds used in this work showed enhanced activities against staphylococci, enterococci, and yeasts, having activities similar to those of amphotericin B against *Candida* spp. and *Cryptococcus neoformans* (19). The activities of these compounds on the growth and membrane-associated ATPase activities of *S. pneumoniae* strains with de-

\* Corresponding author. Mailing address: Unidad de Genética Bacteriana (Consejo Superior de Investigaciones Científicas), Centro Nacional de Microbiología, Instituto de Salud Carlos III, 28220 Majadahonda, Madrid, Spain. Phone: (341) 509-7904. Fax: (341) 509-7919. E-mail: agcampa@isciii.es.

TABLE 1. Susceptibilities of *S. pneumoniae* strains to amino alcohol antimalarials

Strain	MIC ( $\mu\text{g/ml}$ ) <sup>a</sup>									
	OPT	QIN	MFL	OSU8	OSU95	OSU99	OSU199	OSU207	OSU255	OSU259
ATCC 6303	0.75	25	0.15	0.15	0.15	0.15	0.15	0.15	0.07	0.15
ATCC 49619	1.5	50	0.15	0.31	0.62	0.62	0.31	0.15	0.15	0.15
R6	1.5	50	0.15	0.31	0.15	0.62	0.15	0.15	0.62	1.25
MJ11	24	72	0.15	0.31	0.15	0.62	0.15	0.07	0.62	1.25
MJ2	24	230	0.62	0.62	0.31	0.31	0.31	0.62	1.25	1.25
MJQ3	3	144	5	5	0.31	0.31	2.5	5	1.25	1.25
MJQ4	18	180	2.5	2.5	0.31	0.31	0.62	0.62	1.25	1.25
MJM3	6	200	2.5	—	—	—	5	—	—	—
MJM5	48	200	5	—	—	—	5	—	—	—
MJM6	12	100	2.5	—	—	—	0.62	—	—	—
MJM7	3	200	1.25	—	—	—	0.62	—	—	—
MJM8	6	100	5	—	—	—	2.5	—	—	—
MJM9	12	200	0.62	—	—	—	1.25	—	—	—
MJM11	12	100	2.5	—	—	—	1.25	—	—	—
MJM13	24	400	10	—	—	—	10	—	—	—
MJM15	12	200	5	—	—	—	2.5	—	—	—
MJM21	6	200	2.5	—	—	—	2.5	—	—	—
MJM22	24	200	2.5	—	—	—	2.5	—	—	—

<sup>a</sup> —, not determined.

defined mutations in the *c* and *a* subunits of the  $F_0$  complex of the  $F_0F_1$  ATPase have been studied. In addition, MFL-resistant ( $Mfl^r$ ) strains have been isolated and characterized.

#### MATERIALS AND METHODS

**Bacterial strains, growth, and transformation of bacteria.** The following strains of *S. pneumoniae* were used in this study: common laboratory strain R6; OPT- and QIN-resistant R6 derivatives MJ11 and MJ2 (6) and MJQ3 and MJQ4 (24); and strains ATCC 6303 and ATCC 49619, which were used as quality controls for antimalarials (18). Unless stated otherwise, strain R6 was grown in liquid C medium with added 0.08% yeast extract, and transformation was carried out as described by Tomasz (34). Cultures containing  $9 \times 10^9$  CFU per ml were treated with DNA at 0.1  $\mu\text{g/ml}$  for 40 min at 30°C and then at 37°C for 90 min before they were plated onto plates with medium containing 0.3  $\mu\text{g}$  of MFL per ml.

**Susceptibility tests.** MICs were determined by the microdilution method with cation-adjusted Mueller-Hinton broth supplemented with 2.5% lysed horse blood recommended by the National Committee for Clinical Laboratory Standards (25). The inoculum was prepared by suspension of several colonies from an overnight blood agar culture in Mueller-Hinton broth and adjusting the turbidity to that of a 0.5 McFarland standard (ca.  $10^8$  CFU/ml). The suspension was further diluted to provide a final bacterial concentration of  $10^5$  CFU/ml in each well of the microdilution trays. The plates were covered with plastic wrap and incubated in the ambient atmosphere at 37°C for 20 to 24 h. The MIC was defined as the lowest concentration of drug that inhibited visible growth. *S. pneumoniae* strains ATCC 49619 and ATCC 6303 were used for quality control.

**PCR amplification and DNA sequence determination.** *S. pneumoniae* chromosomal DNA was obtained as described previously (6). PCR amplifications were performed with 1 U of *Thermus thermophilus* thermostable DNA polymerase (Biotools), 1  $\mu\text{g}$  of chromosomal DNA, each of the synthetic oligonucleotide primers at a concentration of 0.4  $\mu\text{M}$ , each deoxynucleoside triphosphate at a concentration of 0.2 mM, and 2 mM  $\text{MgCl}_2$  in a final volume of 50  $\mu\text{l}$ . To amplify the *atpC* and *atpA* genes, oligonucleotides atporf1 (5'-ggtcggaatTC TGGATG GCTAACTTGA ACTTG-3'; corresponding to nucleotides -143 to -120 upstream of *atpC*) and atpB56 (5'-GACGGGCTTCTTCAGCTCTGTC-3'; the complementary strand of the primer coding for the DRAEEAR amino acid sequence of subunit *b*), respectively, were used. The 5' end of atporf1 contained an additional sequence (lowercase) that included an *EcoRI* restriction site, which is underlined in the atporf1 oligonucleotide sequence above. Amplification was achieved with an initial cycle of 1 min of denaturation at 94°C and 30 cycles of 30 s at 94°C, 90 s at 55°C, and 75 s of a polymerase extension step at 72°C, with a final 8-min extension step at 72°C and slow cooling at 4°C. The PCR products used in control transformation experiments were obtained with 2.6 U of a proofreading enzyme (Expand High Fidelity PCR system; Boehringer Mann-

heim). The *parE* fragment used in these experiments was amplified with oligonucleotides parE105 (5'-GGAGGGAAATTCGGTCAAGGTGGCTA-3'; this sequence codes for the GGKFGNGGY amino acid sequence) and parE483 (9). Gel electrophoresis of the PCR products was carried out in agarose gels as described previously (32). The DNA fragments were purified with MicroSpin S400 HR columns (Amersham Pharmacia Biotech), and both strands were sequenced with an Applied Biosystems Prism 377 DNA sequencer, according to the protocols provided by the manufacturer, with both the primers used for the PCR amplifications and internal primers.

**ATPase activity determinations.** Membrane-associated ATPase activity was measured as described previously (24). One unit of activity was defined as the amount of enzyme that resulted in the release of 1  $\mu\text{mol}$  of inorganic phosphate per ml per 30 min. The 50% inhibitory concentrations ( $IC_{50}$ s; the means of two independent determinations) were defined as the concentration of the antimalarial agent required for a 50% reduction of enzymatic activity.

**Chemicals.** OPT and QIN were purchased from Sigma Chemical Co., St. Louis, Mo. MFL (Ro 21-5998-000) was provided by Roche Laboratories, Basel, Switzerland. Other antimalarial drugs are identified by The Ohio State University (OSU) code numbers (18). New OSU compounds used were OSU207 and OSU255, which have molecular weights of 427.4 and 444.35, respectively.

**Nucleotide sequence accession numbers.** The sequences reported here have been deposited in GenBank under accession numbers AY061860 to AY061873.

#### RESULTS

**Antimicrobial activities of MFL and OSU compounds.** To test if MFL and the OSU compounds specifically target the  $F_0F_1$  ATPase of *S. pneumoniae*, their MICs for OPT-resistant ( $Opt^r$ ) and QIN-resistant ( $Qin^r$ ) strains were determined (Table 1). The chemical structures of these compounds are shown in Fig. 1. The strains used were R6 mutants with defined point mutations in *atpC* and were selected according to either their OPT resistance (strains MJ11 and MJ2) (6) or QIN resistance (strains MJQ3 and MJQ4) (24). Susceptible strains used as controls were R6, ATCC 6303, and ATCC 49619. MFL and the OSU compounds were 1.2- to 10-fold more active than OPT and 40- to >300-fold more active than QIN in inhibiting the growth of the last three *S. pneumoniae* strains. When the MICs for R6 were compared to those for the MJ2, MJ11, MJQ3, and MJQ4 mutants, significant increases (variations of two or more times the MIC) in the levels of resistance to MFL, OSU8, and




FIG. 1. Chemical structures of the amino alcohol antimalarial agents used in this study.

TABLE 2. Mutational substitutions of the *c* and *a* subunits of *S. pneumoniae* strains resistant to amino alcohol antimalarial agents

Strain	Amino acid change (codon change) from the R6 amino acid sequence <sup>a</sup>			
	<i>c</i> subunit		<i>a</i> subunit	
	$\alpha$ helix 1	$\alpha$ helix 2	$\alpha$ helix 5	$\alpha$ helix 6
R6	<b>13</b> MGVSVGEGLLMN <sup>24</sup>	<b>44</b> MFLGVAFIEGTFV <sup>57</sup>	<b>178</b> GEVMTSLLLL <sup>187</sup>	<b>205</b> AWTAFSVFIS <sup>214</sup>
MJ11		A49T (GCC→ACC)		
MJ2		V48L (GTT→CTT)		
MJQ3	M23I (ATG→ATA)			
MJQ4	G20A (GGT→GCT)			
MJM3				W206G (TGG→GGG)
MJM5	G14S (GGC→AGC)			
MJM6				S214P (TCC→CCC)
MJM7		V57L (GTA→CTA)		
MJM8	M13T (ATG→ACG)			
MJM9		M44V (ATG→GTG)		
MJM10	M23I (ATG→ATT)			
MJM11				F209S (TTT→TCT)
MJM13	N24Y (AAT→TAT)			
MJM14	G20A (GGT→GCT)			
MJM15		G47A (GGT→GCT)		
MJM21			L186P (CTT→CCT)	
MJM22				W206C (TGG→TGT)

<sup>a</sup> The amino acid positions indicated refer to the coordinates for *S. pneumoniae* R6 (GenBank accession no. Z26851) (see reference 6). Residues that changed in the mutant strains indicated are shown in boldface. The E residues possibly involved in proton translocation are showed underlined in the  $\alpha$  helices of the *c* subunit.

OSU199 were observed for strains MJQ3 and MJQ4; and strain MJQ3 also showed a 15-fold increase in its level of resistance to OSU199 and OSU207. However, strain MJ2 showed an increased level of resistance only to MFL and OSU207, whereas there was no difference in the resistance of strain MJ11 to any of the compounds assayed. None of the mutant strains had increased levels of resistance to OSU compounds 95, 99, 255, and 259 (Table 1). These results together suggest that the *c* subunit of the F<sub>0</sub>F<sub>1</sub> complex of the F<sub>0</sub>F<sub>1</sub> ATPase is the target for MFL and OSU compounds 8, 199, and 207 and that for each mutant there is a characteristic pattern of inhibition by the various compounds.

**Characterization of Mfl<sup>r</sup> *S. pneumoniae* strains.** Mfl<sup>r</sup> strains were obtained by plating  $2 \times 10^{10}$  R6 cells in 0.3  $\mu$ g of MFL per ml (twofold the MIC). One isolate, designated isolate MJM1, was obtained. The *atpC* and *atpA* genes of this strain were amplified by PCR with oligonucleotides atporf1 and atpB56 and sequenced. A single mutation was observed in *atpA*; i.e., a TTT codon in strain R6 was changed to a TCT codon, which would produce an F209S change in the predicted amino acid sequence of the *a* subunit of the MJM1 F<sub>0</sub>F<sub>1</sub> ATPase (data not shown). To provide genetic evidence that this change was indeed involved in MFL resistance, the PCR product obtained from strain MJM1 DNA carrying *atpCA* was used to transform competent R6 cells, and the same PCR product, but from R6 DNA, was used as a control. As expected, the PCR product from strain MJM1 transformed R6 to MFL resistance with a high degree of efficiency (greater than  $10^5$  transformants/ml). However, the R6 PCR product was also able to transform R6 to MFL resistance, although at a much lower efficiency ( $10^3$  transformants/ml). Two colonies selected from the R6 transformation with the MJM1 PCR product were found to carry the F209S amino acid change in the *a* subunit, indicating that this mutation is indeed involved in MFL resistance. One of these transformants was named MJM11 (Table

2) and was used for further studies. Several colonies were selected from cultures of isolates used in the transformation experiment with the R6 PCR product, and the sequences of their *atpCA* genes were determined. These strains carried point mutations either in the *atpC* gene (strains MJM5, MJM7, MJM8, MJM9, MJM10, MJM13, MJM14, and MJM15) or in the *atpA* gene (strains MJM3, MJM6, MJM11, MJM21, and MJM22) (Table 2). Genetic evidence showing that the mutations carried by these strains were indeed involved in resistance was obtained by genetic transformation. The PCR products carrying those mutations were able to transform MFL-susceptible (Mfl<sup>s</sup>) strain R6 to MFL resistance with a high degree of efficiency (greater than  $10^5$  transformants/ml). With the exception of strains MJM10 and MJM14, which had amino acid substitutions identical to those in MJQ3 and MJQ4, respectively, the susceptibilities of these strains to MFL and the OSU compounds were analyzed. The MICs of MFL for mutant strains selected for resistance to MFL were significantly increased (from 4- to 64-fold), as were the MICs of OSU199 (4- to 64-fold). All strains except MJM7 had increased levels of resistance to OPT (4- to 32-fold); and all strains except MJM6, MJM8, and MJM11 had increased levels of resistance to QIN (4- to 8-fold) (Table 1).

The appearance of Mfl<sup>r</sup> colonies in the transformation with the 1,277-bp PCR product carrying the R6 *atpCA* genes might be attributed to the error rate of the polymerase. This rate is of 1 error/10 kbp; therefore, one error in 10 molecules of 1 kb would be expected. From our results, about 1% ( $10^{-2}$ ) of the putative nucleotide changes would result in a residue change that confers MFL resistance. Since  $9 \times 10^6$  cells were used in the transformation experiments, the expected number of clones with mutations for MFL resistance would be  $9 \times 10^3$ , a figure consistent with the results obtained. Furthermore, when a 1,158-bp fragment of an unrelated gene (*parE*, coding for the ParE subunit of DNA topoisomerase IV) was amplified and



**c subunit****a subunit**

FIG. 2. Comparisons of the amino acid sequences and secondary structures of the *c* and *a* subunits of *S. pneumoniae* (SPN) and *E. coli* (ECO). The predicted  $\alpha$  helices are shown above the sequence for *S. pneumoniae* and below the sequence for *E. coli*; gray bars indicate predictions made by the PHD method and solid bars indicate predictions made according to Miller et al. (22) (for the *c* subunit) or Valiyaveetil and Fillingame (35) (for the *a* subunit). Residues mutated in the resistant pneumococcal strains are shown in boldface and are underlined. Residues of the *E. coli* *a* subunit involved in interactions with the *c* subunit (11) are shown in boldface and are double underlined. Residues involved in proton translocation in *E. coli* and the equivalent residues in *S. pneumoniae* are boxed.

used in transformation experiments, the frequency of Mfl<sup>r</sup> colonies was equivalent to that obtained in the control experiment with no DNA (no colonies appeared in 0.1 ml of the transformation mixture, which gives a frequency of less than 10 transformants/ml). In addition, similar numbers ( $5 \times 10^2$  transformants/ml) of Mfl<sup>r</sup> colonies were observed when a proofreading enzyme was used in the amplification of the *atpCA* genes, and no Mfl<sup>r</sup> colonies appeared when *parE* was amplified with this enzyme.

**Characteristics of the mutations affecting the *c* and *a* subunits of *S. pneumoniae* F<sub>0</sub>F<sub>1</sub> ATPase.** Predictions of the secondary structures of the *c* and *a* subunits were made by the PHD method (29–31) by using a server available over the Internet (<http://www.embl-heidelberg.de>). Those calculations predicted two transmembrane  $\alpha$  helices for the *c* subunits and

six transmembrane  $\alpha$  helices for the *a* subunits of *S. pneumoniae* and *E. coli* (Fig. 2). Among the 13 Mfl<sup>r</sup> strains obtained by selection with MFL (named MJMX, where the second M represents MFL; Table 2), 8 carried mutations in *atpC* and 5 carried mutations in *atpA*. All amino acid residues that changed in the Mfl<sup>r</sup> stains were located in a predicted transmembrane  $\alpha$  helix (Table 2; Fig. 2). Five strains showed changes in helix 1 in the *c* subunit, and three showed changes in helix 2 in the *c* subunit, while the changes in the *a* subunit were at residues located in helix 5 (one strain) and helix 6 (four strains).

**Inhibition of F<sub>0</sub>F<sub>1</sub> ATPase by MFL and OSU compounds.** Biochemical evidence for the targeting of the F<sub>0</sub>F<sub>1</sub> ATPase by MFL and related compounds was obtained by testing the inhibition of the ATPase activities of membranes obtained from




FIG. 3. Correlations between inhibition of ATPase activity and inhibition of growth of the membranes of the indicated strains by MFL and the OSU compounds (A) and inhibition of growth of the membrane of MJQ4 by the different compounds (B).

strain R6 and the mutant strains. All compounds inhibited the R6 ATPase activity, including those compounds (OSU95, OSU99, OSU255, and OSU259) that inhibited the growth of both sensitive and resistant strains (Table 1). The levels of inhibition of the ATPase activities of the strains by MFL, OSU8, OSU199, and OSU207 ( $IC_{50}$  values) showed good correlations with their MICs (Fig. 3A), indicating that the primary target for these compounds in the cell is the membrane ATPase. A good correlation was also obtained when the  $IC_{50}$ s of the different compounds were plotted against the MICs for strain MJQ4, a strain that showed intermediate sensitivities to most of the compounds tested (Fig. 3B), indicating that all compounds share the same target in the cell.

## DISCUSSION

Combined genetic and biochemical studies support a hairpinlike structure for the *c* subunit, with two  $\alpha$  helices that transverse the membrane in an antiparallel fashion and that are separated by a conserved polar loop region that forms the  $F_1$  binding region (14). The Mfl<sup>r</sup> pneumococcal strains had point mutations that change the amino acid residues located in one of the two transmembrane  $\alpha$  helices of the *c* subunit at positions close to the two acidic residues E19 and E52 (Table 2; Fig. 2). Although previously characterized Opt<sup>r</sup> strains, selected according to their OPT resistance (strains MJ2 and MJ11), had mutations in helix 1 (6) and strains selected according to their QIN resistance (strains MJQ3 and MJQ4) had mutations in helix 2 (24), the strains selected according to their MFL resistance had mutations in either helix 1 or helix 2 (Table 2; Fig. 2). However, independently of the compound used for mutant selection, cross-resistance to those three antimalarial agents was observed (Table 1; Fig. 2 and 3), suggesting that these chemically similar drugs interact with the *c* subunit in the same region of the protein, the ion-binding pocket. In the *E. coli* pocket, the D61 side chain is positioned within a four-helix bundle formed by the front and back faces of two adjacent monomers, with D61 of helix 2 of one monomer packing in close proximity to residues A24 and I28 of helix 1 and with A62 of helix 2 of the neighboring monomer (5, 13) being in a cavity created by the absence of side chains at residues G23, G27, and G58 (8). Interestingly, two of the *S. pneumoniae*-resistant strains (strains MJ11 and MJM5) had amino acid changes at positions equivalent to G23 and G58 of *E. coli* (Table 2; Fig. 2), and G-to-A changes are found in strains MJQ4 and MJM15. All these results suggest a structure for the oligomeric subunit *c* ring of *S. pneumoniae* compatible with that of *E. coli*. However, whereas the *E. coli* *c* subunit has only one carboxyl group (D61 of  $\alpha$  helix 2), the *S. pneumoniae* *c* subunit has two carboxyl groups, one each in a centered position in each  $\alpha$  helix (residues E19 and E52, equivalent to residues I28 and D61, respectively, of *E. coli*), suggesting that both carboxylic groups would be involved in proton translocation. In agreement with this hypothesis, functional ATPases from *E. coli* mutants in which the essential carboxyl group of the *c* subunit has been switched to position 24 of helix 1 (equivalent to V15 of *S. pneumoniae*) (22) or in which a second carboxyl group has been introduced at position 28 (equivalent to E19 of *S. pneumoniae*) (12) have been described.

In addition to mutations in the *c* subunit, five of the Mfl<sup>r</sup>

strains carried mutations in two  $\alpha$  helices of the *a* subunit: position L186 of the penultimate helix and positions W206, F209, and S214 of the ultimate helix (Fig. 2). These two helices are also present in the *E. coli* *a* subunit (Fig. 2). Studies of cysteine cross-linking (11) and nuclear magnetic resonance analysis (28) have established that helix 2 of subunit *c* makes contact with the penultimate helix of subunit *a* (Fig. 2), with residues S207, N214, A217, I221, I223, L224, and I225 lying in one face of the transmembrane helix. These residues then interact with helix 2 of subunit *c*. This proximity of the helices supports the possibility of an interaction between residue R210 of the *a* subunit and residue D61 of the *c* subunit during proton translocation. The location of the L186P change in *S. pneumoniae* MJM21 is in accordance with this model of the  $F_0$  complex of *E. coli*.

Four residue changes present in the resistant *S. pneumoniae* strains are located in the last helix of subunit *a* (Fig. 2), suggesting an interaction between this helix of the *a* subunit and any of the helices of the *c* subunit. Although no cross-link formation has been observed between helix 1 of the *E. coli* subunit *c* and the last helix of subunit *a* (11), a mutant strain with an H245D substitution showed a 45% reduction of normal proton translocation activity (3), and structural studies have defined one hydrophilic pathway between R210 of the *E. coli* *a* subunit and the periplasmic surface, including residues N238, H245, and Q252 of the last helix of this protein (11).

The location and nature of the amino acid changes found in the *c* subunit of the R6 descendants reported here are in accordance with those observed in pneumococcal Opt<sup>r</sup> clinical isolates, which showed G20S, M23I, A49T, V48F, and F50L changes (6, 27). In addition, a change in the *a* subunit (W206S) that corresponds to that found in strain MJM3 has also been recently reported in a clinical isolate of *S. pneumoniae* (27).

## ACKNOWLEDGMENTS

We thank E. García for critical reading of the manuscript. The technical assistance of A. Rodríguez-Bernabé is acknowledged.

A.J.M.-G. received a fellowship from Comunidad Autónoma de Madrid. This study was supported by grant 1274/01 from Instituto de Salud Carlos III.

## REFERENCES

- Abrahams, J. P., A. G. W. Leslie, and J. E. Walker. 1994. Structure at 2.8 Å resolution of  $F_1$ -ATPase from bovine heart mitochondria. *Nature* **370**:621–628.
- Bender, G., S. Sutton, and R. Marquis. 1986. Acid tolerance, proton permeabilities, and membrane ATPases of oral streptococci. *Infect. Immun.* **53**:331–338.
- Cain, B. D., and R. D. Simoni. 1988. Interaction between Glu-219 and His-245 within the *a* subunit of  $F_1F_0$ -ATPase in *Escherichia coli*. *J. Biol. Chem.* **263**:6606–6612.
- Chen, D. K., A. McGeer, J. C. de Azavedo, and D. E. Low. 1999. Decreased susceptibility of *Streptococcus pneumoniae* to fluoroquinolones in Canada. *N. Engl. J. Med.* **341**:233–239.
- Dmitrev, O. Y., P. C. Jones, and R. H. Fillingame. 1999. Structure of the subunit *c* oligomer in the  $F_1F_0$  ATP synthase: model derived from solution structure of the monomer and cross-linking in the native enzyme. *Proc. Natl. Acad. Sci. USA* **96**:7785–7790.
- Fenoll, A., R. Muñoz, E. García, and A. G. de la Campa. 1994. Molecular basis of the optochin-sensitive phenotype of pneumococcus: characterization of the genes encoding the  $F_0$  complex of the *Streptococcus pneumoniae* and *Streptococcus oralis*  $H^+$ -ATPases. *Mol. Microbiol.* **12**:587–598.
- Futai, M., T. Noumi, and M. Maeda. 1989. ATP synthase ( $H^+$ -ATPase): results by combined biochemical and molecular biological approaches. *Annu. Rev. Biochem.* **58**:111–136.
- Girvin, M. E., V. K. Rastogi, F. Abridgaard, J. L. Markley, and R. H. Fillingame. 1998. Solution structure of the transmembrane  $H^+$ -translocating subunit *c* of the  $F_1F_0$  ATP synthase. *Biochemistry* **37**:8817–8824.

9. González, I., M. Georgiou, F. Alcáide, D. Balas, J. Liñares, and A. G. de la Campa. 1998. Fluoroquinolone resistance mutations in the *parC*, *parE*, and *gyrA* genes of clinical isolates of viridans group streptococci. *Antimicrob. Agents Chemother.* **42**:2792–2798.
10. Hoffman, J., M. S. Cetron, M. M. Farley, W. S. Baughman, R. R. Facklam, J. A. Elliot, K. A. Deaver, and R. F. Breiman. 1995. The prevalence of drug-resistant *Streptococcus pneumoniae* in Atlanta. *N. Engl. J. Med.* **333**:481–486.
11. Jiang, W., and R. H. Fillingame. 1998. Interacting helical faces of subunits *a* and *c* in the  $F_1F_0$  ATP synthase of *Escherichia coli* defined by disulfide cross-linking. *Proc. Natl. Acad. Sci. USA* **95**:6607–6612.
12. Jones, P. C. 2001. Introduction of a carboxyl group in the first transmembrane helix of *Escherichia coli*  $F_1F_0$  ATPase subunit *c* and cytoplasmic pH regulation. *J. Bacteriol.* **183**:1524–1530.
13. Jones, P. C., W. Jiang, and R. H. Fillingame. 1998. Arrangement of the multicopy  $H^+$ -translocating subunit *c* in the membrane sector of the *Escherichia coli*  $F_1F_0$  ATP synthase. *J. Biol. Chem.* **273**:1718–1785.
14. Kakinuma, Y. 1998. Inorganic cation transport and energy transduction in *Enterococcus hirae* and other streptococci. *Microbiol. Mol. Biol. Rev.* **62**:1021–1045.
15. Kobayashi, H., T. Suzuki, N. Kinoshita, and T. Unemoto. 1984. Amplification of the *Streptococcus faecalis* proton-translocating ATPase by a decrease in cytoplasmic pH. *J. Bacteriol.* **158**:1157–1160.
16. Kobayashi, H., T. Suzuki, and T. Unemoto. 1986. Streptococcal cytoplasmic pH is regulated by changes in amount and activity of a proton-translocating ATPase. *J. Biol. Chem.* **261**:627–630.
17. Koebmann, B. J., D. Nilsson, O. P. Kuipers, and P. R. Jensen. 2000. The membrane-bound  $H^+$ -ATPase complex is essential for growth of *Lactococcus lactis*. *J. Bacteriol.* **182**:4738–4743.
18. Kullen, M. J., and T. R. Klaenhammer. 1999. Identification of the pH-inducible, proton-translocating  $F_1F_0$ -ATPase (*atpBEFHAGDC*) operon of *Lactobacillus acidophilus* by differential display: gene structure, cloning and characterization. *Mol. Microbiol.* **33**:1152–1161.
19. Kunin, C. M., and W. Y. Ellis. 2000. Antimicrobial activities of mefloquine and a series of related compounds. *Antimicrob. Agents Chemother.* **44**:848–852.
20. Liñares, J., A. G. de la Campa, and R. Pallarés. 1999. Fluoroquinolone resistance in *Streptococcus pneumoniae*. *N. Engl. J. Med.* **341**:1546–1548.
21. Martín-Galiano, A. J., M. J. Ferrándiz, and A. G. de la Campa. 2001. The promoter of the operon encoding the  $F_0F_1 H^+$ -ATPase of *Streptococcus pneumoniae* is inducible by pH. *Mol. Microbiol.* **6**:1327–1338.
22. Miller, M. J., M. Oldenburg, and R. H. Fillingame. 1990. The essential carboxyl group in subunit *c* of the  $F_1F_0$  ATP synthase can be moved and  $H^+$ -translocation retained. *Proc. Natl. Acad. Sci. USA* **87**:4900–4904.
23. Moore, H. F., and A. M. Chesney. 1917. A study of ethylhydrocuprein (optochin) in the treatment of acute lobar pneumonia. *Arch. Intern. Med.* **19**:611–682.
24. Muñoz, R., E. García, and A. G. de la Campa. 1996. Quinine specifically inhibits the proteolipid subunit of the  $F_0F_1 H^+$ -ATPase of *Streptococcus pneumoniae*. *J. Bacteriol.* **178**:2455–2458.
25. National Committee for Clinical Laboratory Standards. 2000. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically. Approved standard M7-A4. National Committee for Clinical Laboratory Standards, Villanova, Pa.
26. Pallarés, R., J. Liñares, M. Vadillo, C. Cabellos, F. Manresa, P. F. Viladrich, R. Martín, and F. Gudiol. 1995. Resistance to penicillin and cephalosporin and mortality from severe pneumococcal pneumonia in Barcelona, Spain. *N. Engl. J. Med.* **333**:474–480.
27. Píakis, A., J. M. Campos, W. J. Rodríguez, and J. M. Keith. 2001. Optochin resistance in *Streptococcus pneumoniae*: mechanism, significance, and clinical implications. *J. Infect. Dis.* **184**:582–590.
28. Rastogi, V. K., and M. E. Girvin. 1999. Structural changes linked to proton translocation by subunit *c* of the ATP synthase. *Nature* **402**:263–268.
29. Rost, B. 1996. PHD: predicting one-dimensional protein structure by profile-based neural networks. *Methods Enzymol.* **266**:525–539.
30. Rost, B., P. Fariselli, and R. Casadio. 1996. Topology prediction for helical transmembrane proteins at 86% accuracy. *Protein Sci.* **5**:1704–1718.
31. Rost, B., and C. Sander. 1993. Prediction of protein secondary structure at better than 70% accuracy. *J. Mol. Biol.* **232**:584–599.
32. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
33. Stock, D., A. G. W. Leslie, and J. E. Walker. 1999. Molecular architecture of the rotary motor in ATP synthase. *Science* **286**:1700–1705.
34. Tomasz, A. 1970. Cellular metabolism in genetic transformation of pneumococci: requirement for protein synthesis during induction of competence. *J. Bacteriol.* **101**:860–871.
35. Valiyaveetil, F. I., and R. H. Fillingame. 1998. Transmembrane topography of subunit *a* in the *Escherichia coli*  $F_1F_0$  ATP synthase. *J. Biol. Chem.* **273**:16241–16247.