

## *Enterococcus faecium* Low-Affinity *pbp5* Is a Transferable Determinant

Louis B. Rice,<sup>1,2,3\*</sup> Lenore L. Carias,<sup>3</sup> Susan Rudin,<sup>3</sup> Viera Laktičová,<sup>2</sup>  
Aaron Wood,<sup>3</sup> and Rebecca Hutton-Thomas<sup>1</sup>

Medical and Research Services, Louis Stokes Cleveland VA Medical Center,<sup>1</sup> the Cleveland VA Research  
and Education Foundation,<sup>2</sup> and Department of Medicine, Case Medical School,<sup>3</sup> Cleveland, Ohio

Received 28 June 2005/Returned for modification 8 August 2005/Accepted 19 September 2005

Using 15 unrelated *Enterococcus faecium* isolates as donors, we demonstrated that ampicillin resistance was transferable to an *E. faecium* recipient containing a *pbp5* deletion for all but four strains. The transfers occurred at low frequencies (generally ca.  $10^{-9}$  transconjugants/recipient CFU), consistent with chromosome-to-chromosome transfer. *pbp5* transfer occurred within large genetic regions, and insertion into the recipient genome occurred most commonly into the recipient *Sma*I restriction fragment that had been created by the previous *pbp5* deletion. Restriction mapping of the region upstream of *pbp5* revealed a commonality of fragment sizes among the clinical isolates from the United States which differed significantly from those of three strains that were isolated from turkey feces. These data prove conclusively that *E. faecium pbp5* is a transferable determinant, even in the absence of a coresiding vancomycin resistance mobile element. They also suggest that the spread of high-level ampicillin resistance among U.S. *E. faecium* strains is due in part to the transfer of low-affinity *pbp5* between clinical isolates.

Among the more intriguing aspects of the emergence and spread of glycopeptide resistance in *Enterococcus faecium* is the almost universal association of vancomycin resistance with clinically significant levels of resistance to ampicillin (16). The mechanisms for resistance to glycopeptides and ampicillin are distinct. Resistance to glycopeptides is most often conferred by the acquisition of glycopeptide-resistant VanA or VanB operons, both of which are incorporated within mobile elements (7). Resistance to ampicillin is conferred most commonly by expression of low-affinity penicillin-binding protein 5 (PBP5) (20). The genetic determinant encoding PBP5 (*pbp5*) is located in the bacterial chromosome and considered to be intrinsic to the species. Elevated levels of ampicillin resistance found in clinical *E. faecium* isolates are most commonly associated with point mutations in PBP5 that lower the affinities for  $\beta$ -lactam antibiotics (11, 15, 21).

Although monoclonal outbreaks of glycopeptide-resistant *E. faecium* have been reported, many areas of endemicity of highly resistant strains report several distinct clones expressing resistance (5, 10). Polyclonal glycopeptide resistance has often been attributed to the movement of transposons Tn1546 (1, 17) and Tn5382/1549 (2, 6, 9), which encode VanA- and VanB-type resistance, respectively. The movement of these transposons has undoubtedly contributed to the rapidly rising prevalence of vancomycin resistance in U.S. hospitals. The nearly universal association of vancomycin resistance with ampicillin resistance in the United States implies that ampicillin-resistant *E. faecium* strains are polyclonal as well. The existence of multiple clones of ampicillin-resistant *E. faecium* raises the possibility that the chromosomally encoded *pbp5* of *E. faecium* is transferable.

In prior work, we demonstrated the transferability of *pbp5* from *E. faecium* strains in which the VanB mobile element Tn5382/1549 was inserted immediately downstream of *pbp5* (2). As Tn5382/1549 bears significant similarities to conjugative transposons, the possibility remains that this transposon in some way fueled the transfer of both vancomycin and ampicillin resistance. The present study was undertaken to determine whether we could demonstrate the transferability of *pbp5* alone from additional *E. faecium* strains of diverse origins, including those that are susceptible to vancomycin.

### MATERIALS AND METHODS

**Bacterial strains.** The *E. faecium* strains used in this study were derived from a variety of sources and are detailed in Table 1. Strain *E. faecium* TX0016 is the clinical isolate used to determine the sequence of the *E. faecium* genome (incomplete sequence available from the Baylor College of Medicine [ftp://ftp.hgsc.bcm.tmc.edu/pub/data/Efaecium/] and the Joint Genome Institute [http://genome.jgi-psf.org/draft\_microbes/entfa/entfa.home.html]). It was kindly provided to us by Barbara E. Murray. The recipient strain for transfer of *pbp5* was *E. faecium* strain D344SRF (13), an ampicillin-susceptible derivative of *E. faecium* strain D344R in which *pbp5* was spontaneously deleted from the chromosome and was subsequently selected for resistance to fusidic acid and rifampin by sequential plating on high concentrations of each antibiotic. MICs for ampicillin and vancomycin were determined by agar dilution using techniques as defined in the CLSI guidelines, except that brain heart infusion agar was used instead of Mueller-Hinton agar. Transconjugants expressing resistance to ampicillin or vancomycin exceeding that for strain D344SRF (0.25 to 0.5  $\mu$ g/ml and 2 to 4  $\mu$ g/ml, respectively) were deemed resistant to those antibiotics.

**Conjugation experiments.** Matings were performed using a technique designed to detect low-frequency events as previously described (12) using ampicillin (2 to 10  $\mu$ g/ml), fusidic acid (25  $\mu$ g/ml), and rifampin (50  $\mu$ g/ml) in the agar for the selection of transconjugants. Plates were examined daily for 4 days to detect the appearance of colonies. Colonies that appeared on the mating plates were restreaked on selective plates to confirm ampicillin resistance. Matings were performed one to three times per mating pair. Transfer frequencies were calculated as numbers of transconjugants per recipient CFU. Selected colonies were subjected to detailed analysis by pulsed-field gel electrophoresis (PFGE) to confirm that they were indeed transconjugants. Specific rates were calculated only if colonies from a mating were shown by PFGE to be true transconjugants.

**Genetic techniques.** Lysis of *E. faecium* cells for PFGE was performed as previously described (2). Digestion of genomic DNA with the restriction enzyme *Sma*I or *Bst*ZI (Promega, Madison, Wis.) used 40 U of restriction enzyme in

\* Corresponding author. Mailing address: Medical Service 111(W), Louis Stokes Cleveland VA Medical Center, 10701 East Blvd., Cleveland, OH 44106. Phone: (216) 791-3800. Fax: (216) 231-3289. E-mail: louis.rice@med.va.gov.

TABLE 1. Bacterial strains and plasmids used in these experiments

Bacterial strain or plasmid	Relevant antimicrobial resistance	Description
<i>E. faecium</i> strains		
C68	Ap <sup>r</sup> Vm <sup>r</sup>	Clinical isolate initially used to define sequences downstream of <i>pbp5</i> (2)
D344R	Ap <sup>r</sup> Vm <sup>s</sup> , Ery <sup>r</sup> , Tc <sup>r</sup>	Ampicillin-resistant clinical isolate (21)
D344SRF	Ap <sup>s</sup> ( <i>pbp5</i> negative) Ery <sup>r</sup> Tc <sup>r</sup> , Fus <sup>r</sup> Rif <sup>r</sup>	PBP5 <sup>-</sup> , ampicillin-susceptible derivative of strain D344R (11)
TX0016	Low-level Ap <sup>r</sup> , Vm <sup>s</sup>	Ampicillin-resistant clinical isolate whose partial genome sequence is available (Baylor College of Medicine [ftp://ftp.hgsc.bcm.tmc.edu/pub/data/Efaecium]; Joint Genome Institute [http://genome.jgi-psf.org/draft_microbes/entfa/entfa.home.html])
D14	Ap <sup>r</sup> Vm <sup>s</sup>	Ampicillin-resistant clinical isolate originally isolated from Providence, R.I. (4), clonally distinct from strains D24, D25, and D29
D24	Ap <sup>r</sup> Vm <sup>s</sup>	Ampicillin-resistant clinical isolate originally isolated from Richmond, Va. (4), clonally distinct from strains D14, D25, and D29
D25	Ap <sup>r</sup> Vm <sup>s</sup>	Ampicillin-resistant clinical isolate originally isolated from Durham, N.C. (4), clonally distinct from strains D14, D24, and D29
D29	Ap <sup>r</sup> Vm <sup>s</sup>	Ampicillin-resistant clinical isolate originally isolated from Calif. (4), clonally distinct from strains D14, D24, and D25
TJ386	Ap <sup>r</sup> Vm <sup>r</sup>	Ampicillin- and vancomycin-resistant clinical isolate originally isolated from Philadelphia, Pa. (kindly provided by Henry Fraimow [18])
WB312	Ap <sup>r</sup> Vm <sup>r</sup>	Ampicillin- and vancomycin-resistant clinical isolate originally isolated from Scranton, Pa. (kindly provided by Henry Fraimow [18])
WC176	Ap <sup>r</sup> Vm <sup>r</sup>	Ampicillin- and vancomycin-resistant clinical isolate originally isolated in Westchester, N.Y. (kindly provided by Henry Fraimow [18])
T420	Low-level Ap <sup>r</sup> , Vm <sup>s</sup>	Strain derived from turkey feces (did not yield transconjugants) (19)
T450	Low-level Ap <sup>r</sup> , Vm <sup>s</sup>	Strain derived from turkey feces (did not yield transconjugants) (19)
T471	Low-level Ap <sup>r</sup> , Vm <sup>s</sup>	Strain derived from turkey feces (did yield transconjugants) (19)
T636	Low-level Ap <sup>r</sup> , Vm <sup>s</sup>	Strain derived from turkey feces (did yield transconjugants) (19)
T642	Low-level Ap <sup>r</sup> , Vm <sup>s</sup>	Strain derived from turkey feces (did yield transconjugants) (19)
GE-1	Ap <sup>s</sup> ( <i>pbp5</i> negative) Fus <sup>r</sup> Rif <sup>r</sup>	Recipient for enterococcal mating experiments, referenced here because it was the recipient strain used in the mating experiment that created CV133 (2)
CV133	Ap <sup>r</sup> Vm <sup>r</sup>	Ampicillin- and vancomycin-resistant transconjugant resulting from mating between C68 and GE-1 (2); genomic DNA was the source for probe 4 as depicted in Fig. 1; in this region, it is identical to the genome of C68
Plasmids		
PBCSK(-)	Cm <sup>r</sup>	Cloning vector (Stratagene)
pCWR570	Cm <sup>r</sup>	18-Kb PstI fragment of CV133 genomic DNA cloned into pBCSK-
PCWR576	Cm <sup>r</sup>	5-Kb XbaI digestion fragment of insert from pCWR570 cloned into pBCSK-

400  $\mu$ l of designated restriction buffer. The program for the separation of restriction digests used the autoalgorithm function (Bio-Rad, Hercules, Calif.) with the following details: separation was set for a low of 20 kb and a high of 500 kb (20 kb to 100 kb for BstZI) to run over 24 h, the calibration factor was 1.0, gels consisted of 1% PFC agarose and were run in a bath of 0.5 $\times$  Tris-borate-EDTA at 14°C with a gradient of 6.0 V/cm, the included angle was 120°, the initial switch time was 2.98 s, and the final switch time was 44.69 s (8.53 s for BstZI) with linear ramping. Extraction of genomic DNA from *E. faecium* strains for routine restriction digestion was performed as previously described (14). Southern transfer, hybridization using digoxigenin-labeled probes, and detection with chemiluminescence assays were performed by standard techniques as previously described (2). Probes used to hybridize transferred DNA were created by incorporating digoxigenin into PCRs as previously described (2) using the following primers for *pbp5*: PBP5-2004-1 (5'-GAATCCAGAATTAAGCAGTAATGG-3'), PBP5-2004-2 (5'-CGCAACAGTTGATCCAGC-3'), probe 1 from Fig. 1, 523-F (5'-CTGGGTTCCATCAACGACTG-3'), 523-G (5'-TCATTGATAAGCGCG AGTGC-3'), probe 2 from Fig. 1, 536-1 (5'-GACGTCGAAGATGCGGTA-3'), 536-2 (5'-AGTATTGCGATGGGGAATGC-3'), probe 3 from Fig. 1, lipase-1 (5'-GAAGGACCGACTGAAGGGATT-3'), and lipase-2 (5'-CGGCTTTTT GTCCCTTCTG-3'). The probes were based on sequence analysis of *pbp5* and its upstream region from *E. faecium* strain C68 and its transconjugant, strain CV133. The locations of the probes upstream of *pbp5* are shown in Fig. 1. The single exception was probe 4, which consisted of a 1.5-kb fragment derived from pCWR576 as detailed in Table 1 and indicated in Fig. 1. This restriction fragment was extracted from the gel and labeled with digoxigenin by using the Klenow fragment as previously described (14).

## RESULTS AND DISCUSSION

**Transfer of *pbp5*.** Conjugation experiments were performed between the 15 strains listed in Table 2 and strain D344SRF, with selection on ampicillin, fusidic acid, and rifampin. Frequencies of transfer were universally low (Table 2), ranging from a low of  $5.3 \times 10^{-10}$  to a high of  $2.56 \times 10^{-7}$ . Four of the 11 donor strains from which transconjugants were isolated expressed vancomycin resistance. In three of these strains, transfer of ampicillin resistance occurred without transfer of vancomycin resistance. Interestingly, one of the three strains from which vancomycin resistance was not transferred along with resistance to ampicillin was C68, in which VanB transposon Tn5382/1549 was integrated immediately downstream of *pbp5* and from which the two determinants had been shown to transfer together in the past (vancomycin selection was used in previous experiments). The absence of vancomycin resistance from the transconjugant suggests that Tn5382/1549 may have been excised from its location in the process of transfer (2, 6).

Two clinical strains (D14, D25) and two strains isolated from turkeys (T420, T450) yielded no transconjugants. One clinical




FIG. 1. Map of the region extending approximately 62 kb upstream of *pbp5* in strains C68 and CV133. Locations of relevant restriction sites are shown. The precise locations of the probes used in the mapping experiments are indicated by the vertical arrows. The expected restriction fragments based on the C68/CV133 map that were hybridized are in white. *pbp5* is located at the extreme right end of the map. Just beyond the right end of the map in C68 and CV133 lies Tn5382/1549.

strain, D14, yielded no colonies on mating plates during any mating. The second donor, D25, yielded colonies on mating plates initially, but these colonies failed to grow repeatedly when restreaked on identical ampicillin concentrations. Neither T420 nor T450 produced any colonies on mating plates. In fact, one strain (T420) yielded no colonies on plates designed to determine recipient numbers, suggesting that the donor produced a substance that was lethal to the recipient. The 11 remaining strains all produced transconjugants that were verified as such by PFGE.

PFGE of eight donor and transconjugant pairs is shown in Fig. 2A. Comparison of the PFGE patterns seen in the eight donor strains (D344R is not well shown in Fig. 2A) confirmed that they are genetically unrelated to each other (a difference of greater than five bands). All of the transconjugants had digestion patterns similar to that of D344SRF, confirming that they are true transconjugants. In all but one transconjugant, a ca.-170-kb *SmaI* digestion fragment (Fig. 2A) of D344SRF had disappeared and larger bands of various lengths had appeared. In a previous work, we showed that this 170-kb band is created as a result of the deletion of *pbp5* and surrounding regions

from the D344R genome (11a). A clearer picture of restriction digests of D344R, D344SRF, and their transconjugant CV333 is shown in Fig. 2B. These data suggest that integration of the transferred *pbp5* region occurs within the same *SmaI* restriction fragment from which the *pbp5* region had been excised, consistent with either site-specific integration of the transferred region or recombination across homologous genomic regions. Similar results were seen with the three donor strains not shown on this gel (for strain D24, disappearance of the band in one of two transconjugants; for strain T636, disappearance of the band in one of one transconjugant; and for strain T642, disappearance of the band in five of five transconjugants).

Southern hybridization of DNA transferred from the PFGE gel using an internal amplification product of *pbp5* as a probe is shown in Fig. 2A. Similar-sized *pbp5*-hybridizing *SmaI* restriction fragments were seen in five of the transconjugants, which corresponded in size to the *pbp5*-hybridizing *SmaI* fragments in three of the donor strains (TX0016, D29, TJ386). These include the transconjugant resulting from the mating in which C68 was the donor strain. In this instance, the *pbp5*-

TABLE 2. Characteristics of donors and transconjugants and frequency of transfer of ampicillin resistance

<i>E. faecium</i> donor strain <sup>a</sup> (drug resistance)	Transconjugant (drug resistance)	Transfer frequency
C68 (Ap <sup>r</sup> Vm <sup>s</sup> )	CV384 (Ap <sup>r</sup> Vm <sup>s</sup> )	3.3 × 10 <sup>-8</sup>
TX0016 (Ap <sup>r</sup> Vm <sup>s</sup> )	CV477 (Ap <sup>r</sup> Vm <sup>s</sup> )	5.3 × 10 <sup>-10</sup>
D14 (Ap <sup>r</sup> Vm <sup>s</sup> )	None	<5 × 10 <sup>-10</sup>
D24 (Ap <sup>r</sup> Vm <sup>s</sup> )	CV465 (Ap <sup>r</sup> Vm <sup>s</sup> )	7.14 × 10 <sup>-9</sup>
D25 (Ap <sup>r</sup> Vm <sup>s</sup> )	Colonies appeared on plates but did not regrow on ampicillin	Unknown
D29 (Ap <sup>r</sup> Vm <sup>s</sup> )	CV420 (Ap <sup>r</sup> Vm <sup>s</sup> )	1.35 × 10 <sup>-7</sup>
TJ386 (Ap <sup>r</sup> Vm <sup>r</sup> )	CV426 (Ap <sup>r</sup> Vm <sup>s</sup> )	5.7 × 10 <sup>-9</sup>
WB312 (Ap <sup>r</sup> Vm <sup>r</sup> )	CV433 (Ap <sup>r</sup> Vm <sup>r</sup> )	1 × 10 <sup>-9</sup>
WC176 (Ap <sup>r</sup> Vm <sup>r</sup> )	CV437 (Ap <sup>r</sup> Vm <sup>s</sup> )	2.56 × 10 <sup>-7</sup>
T420 (low-level Ap <sup>r</sup> , Vm <sup>s</sup> )	None	Unknown <sup>b</sup>
T450 (low-level Ap <sup>r</sup> , Vm <sup>s</sup> )	None	<5 × 10 <sup>-10</sup>
T471 (low-level Ap <sup>r</sup> , Vm <sup>s</sup> )	CV343 (low-level Ap <sup>r</sup> , Vm <sup>s</sup> )	3.3 × 10 <sup>-9</sup>
T636 (low-level Ap <sup>r</sup> , Vm <sup>s</sup> )	CV470 (low-level Ap <sup>r</sup> , Vm <sup>s</sup> )	3.08 × 10 <sup>-9</sup>
T642 (low-level Ap <sup>r</sup> , Vm <sup>s</sup> )	CV472 (low-level Ap <sup>r</sup> , Vm <sup>s</sup> )	3.28 × 10 <sup>-9</sup>
D344R (Ap <sup>r</sup> Vm <sup>s</sup> )	CV333 (Ap <sup>r</sup> Vm <sup>s</sup> )	3.3 × 10 <sup>-8</sup>

<sup>a</sup> Recipient strain for all matings was *E. faecium* D344SRF.

<sup>b</sup> No recipients grew on colony-counting plates, suggesting a donor-killing recipient.



## BstZI Digestion of Genomic DNA




FIG. 3. PFGE agarose separation of genomic DNA from strains CV437 (lanes A) and CV426 (lanes B). Both strains revealed minimal to no hybridization of restriction fragments with the *pbp5* probe after digestion with *Sma*I (Fig. 2A), implying possible supercoiling of the transferred fragment, which remained in the well. In both cases, *Bst*ZI digestion resulted in entry of the hybridizing fragment into the gel, consistent with a supercoiled form successfully digested with *Bst*ZI. Probes used were for *pbp5*, along with probes 3 and 4 as described in the Materials and Methods. CV426 demonstrated no hybridization to probe 4, indicating that this region of DNA was not involved in the transfer.

to the DNA in the well. In order to assess the possibility that hybridization to the well in these strains reflected a supercoiled state (without an internal *Sma*I site), we digested the two transconjugants with restriction enzyme *Bst*ZI and hybridized them with *pbp5* as well as probe 3 and probe 4. Results of these experiments are shown in Fig. 3. In both instances, strong hybridization to bands within the gel was observed. For transconjugant CV437, all three probes hybridized to the same restriction fragment. In contrast, CV426 demonstrated hybridization to two distinct fragments with both *pbp5* and probe 3 and did not hybridize at all to probe 4. These data support the idea that hybridization to the wells after *Sma*I digestion reflects the supercoiling of a non-*Sma*I-containing segment of transferred DNA. They also indicate that the transfer event resulting in CV426 did not include the region that hybridized to probe 4 in the donor strain and would indicate that a duplication of the *pbp5*-probe 3 region had occurred. The duplication of this region may also explain the increased intensity of the well hybridization seen for CV426 in Fig. 2A.

These data indicate that *pbp5* can be transferred to an *E. faecium* recipient from a variety of *E. faecium* strains of diverse origins (human clinical isolates from both North America and Europe, as well as isolates derived from turkey feces), whether or not the donor strain also harbored a mobile vancomycin resistance operon. In this context, the rapid emergence of vancomycin- and ampicillin-resistant *E. faecium* strains in the United States is very likely the result of not only mobile vancomycin resistance genes but also mobile ampicillin resistance. It should be acknowledged that the mating pairs employed in these experiments differ from those that would be anticipated to occur in nature, where the recipients would all be expected to have their own copy of *pbp5*. Preliminary mating experiments indicate that transfer of *pbp5* from strain C68 into strain D344R occurs, with eventual replacement of the

TABLE 3. Hybridization of restriction fragments from selected strains using probes derived from regions upstream of *pbp5* in CV133 and C68

Strain	Size (kb) of hybridizing restriction fragment for probe:			
	1 (BamHI-PstI)	2 (BglIII)	3 (XbaI)	4 (XbaI)
C68	12.5	4	9	5.2
D14	Ca. 23	5.2	4.5	6
D24	12.5	4	9	5.2
D25	12.5	4	9	5.2
D279	12.5	4	9	5.2
TJ386	12.5	4	9	5.2
WB312	12.5	4	9	5.2
WC176	12.5	4	9	5.2
T420	12.5	4	4.5	7
T450	9	16	4.5	No hybridization
T471	12.5	10	Ca. 23	6.2

D344R *pbp5* region with that acquired from strain C68 (data not shown).

**Mapping the upstream *pbp5* region.** In an effort to further define the nature of the *pbp5*/Tn5382/1549 element from strain C68, we cloned and determined the sequence from restriction fragments upstream of *pbp5* from strain CV133, a transconjugant resulting from a mating between strain C68 and ampicillin-susceptible recipient strain *E. faecium* GE-1. We used sequence from these various clones to construct probes to span a region extending approximately 60 kb upstream of *pbp5*. All probes hybridized to identical-sized fragments in strains C68 and CV133 and did not hybridize to *E. faecium* strain GE-1, confirming that they were within the transferred region (data not shown). These same probes were used to hybridize the restriction digestions of several clinical ampicillin-resistant *E. faecium* (D14, D24, D25, and D29) and ampicillin- and vancomycin-resistant *E. faecium* (C68, TJ386, WB312, and WC176) strains and three *E. faecium* strains derived from turkey feces (T420, T450, and T471). Only one of these turkey strains (T471) yielded transconjugants in mating experiments. The location of the probes is detailed in Fig. 1. Results from these experiments are shown in Table 3.

All of the clinical strains that yielded transconjugants exhibited identical restriction fragment sizes hybridizing to each of the four probes. In addition, the ampicillin-resistant clinical strain that yielded transconjugants that did not grow on repeated inoculation (D25) yielded an identical pattern. In contrast, the ampicillin-resistant clinical strain that never produced any transconjugants (D14) yielded entirely different restriction fragments. The three strains isolated from turkeys for whom restriction mapping was performed (only one of which produced transconjugants) yielded fragments of different sizes than those of the clinical strains, with one strain showing two bands of similar sizes, one strain showing one band of a similar size, and the third strain showing no size similarities. *E. faecium* strain TX0016 was not included in these experiments, but restriction enzyme site identification using the available genome database indicates that at least three of the four probes will hybridize to fragments identical in size to those observed for strain C68 (data not shown).

The segments within which *pbp5* transfers are likely to be very heterogeneous, given the marked differences in restriction

maps between the clinical isolates and those isolated from turkey feces. In light of this variability, it is also interesting that the region upstream of *pbp5* in both vancomycin-resistant and -susceptible clinical isolates appears to be identical by restriction mapping. These data suggest that there may be one version of a transferable element that is primarily responsible for transfer in U.S. clinical strains and that this element was present and spreading before the emergence and spread of vancomycin resistance within the genus. If this one "super-transfer" element also encodes functions that confer virulence on the bacterium, then the increased importance of *E. faecium* as a nosocomial pathogen and its increased resistance to ampicillin could be explained by a single mechanism (3, 8). We are in the process of conducting a more detailed analysis of the transferred region from strain C68.

Some insight into the potential size of the mobile element can be derived from our recent work analyzing the mechanism by which *pbp5* was deleted from the D344R chromosome to yield D344S (11a). In that strain, a ca.-178-kb region extending from approximately 78 kb upstream of *pbp5* to 100 kb downstream has been deleted. The deletion event resulted from the interaction of heterologous Tn916-like transposons. One of these elements, likely to be nearly identical to Tn916, was not present at the same position of the C68 chromosome. The other, designated Tn5386, was not present in our clinical strains. However, it is conceivable that other Tn916-like elements present in these clinical strains, or perhaps even smaller insertion sequence elements, can prompt the excision of smaller or larger segments of the *E. faecium* genome that contain *pbp5*. Our observation that insertion occurred most commonly into the SmaI site vacated by *pbp5* suggests some consistency to the process of insertion. However, the fact that insertion into this fragment does not always occur and that transconjugant CV426 contains two copies of both *pbp5* and the region hybridized by probe 3, while at the same time lacking the region hybridized by probe 4, suggests substantial potential for variability.

In conclusion, we have shown that *pbp5*, the gene that confers high-level  $\beta$ -lactam resistance in *E. faecium*, is located within transferable elements and can be transferred from a range of *E. faecium* donor strains to an *E. faecium* recipient. These results shed new light on the molecular epidemiology of multidrug-resistant *E. faecium* strains both in the United States and around the world. It will be important to precisely characterize the transferable element that appears to be common to several different strains to better allow us to determine the specific impact of this element on the spread of high-level ampicillin resistance within the species.

#### ACKNOWLEDGMENTS

This study was supported by a Merit Review from the Department of Veterans Affairs and by a grant from the National Institute of Allergy and Infectious Diseases (AI 045626) to L.B.R.

#### REFERENCES

1. Arthur, M., C. Molinas, F. Depardieu, and P. Courvalin. 1993. Characterization of Tn1546, a Tn3-related transposon conferring glycopeptide resistance by synthesis of depsipeptide peptidoglycan precursors in *Enterococcus faecium* BM4147. *J. Bacteriol.* **175**:117–127.
2. Carias, L. L., S. D. Rudin, C. J. Donskey, and L. B. Rice. 1998. Genetic

linkage and cotransfer of a novel, *vanB*-containing transposon (Tn5382) and a low-affinity penicillin-binding protein 5 gene in a clinical vancomycin-resistant *Enterococcus faecium* isolate. *J. Bacteriol.* **180**:4426–4434.

3. Chirurgi, V. A., S. E. Oster, A. A. Goldberg, and R. E. McCabe. 1992. Nosocomial acquisition of  $\beta$ -lactamase-negative, ampicillin-resistant enterococcus. *Arch. Int. Med.* **152**:1457–1461.
4. Donobedian, S. M., J. W. Chow, J. M. Boyce, R. E. McCabe, S. M. Markowitz, P. E. Coudron, A. Kuritza, C. L. Pierson, and M. J. Zervos. 1992. Molecular typing of ampicillin-resistant, non- $\beta$ -lactamase-producing *Enterococcus faecium* isolates from diverse geographic areas. *J. Clin. Microbiol.* **30**:2757–2761.
5. Donskey, C. J., J. R. Schreiber, M. R. Jacobs, R. Shekar, F. Smith, S. Gordon, R. A. Salata, C. Whalen, and L. B. Rice. 1999. A polyclonal outbreak of predominantly VanB vancomycin-resistant enterococci in Northeast Ohio. *Clin. Infect. Dis.* **29**:573–579.
6. Garnier, F., S. Taourit, P. Glaser, P. Courvalin, and M. Galimand. 2000. Characterization of transposon Tn1549, conferring VanB-type resistance in *Enterococcus* spp. *Microbiology* **146**:1481–1489.
7. Gold, H. S. 2001. Vancomycin-resistant enterococci: mechanisms and clinical observations. *Clin. Infect. Dis.* **33**:210–219.
8. Grayson, M. L., G. M. Eliopoulos, C. B. Wennersten, K. L. Ruoff, P. C. DeGirolami, M.-J. Ferraro, and R. C. Moellering, Jr. 1991. Increasing resistance to  $\beta$ -lactam antibiotics among clinical isolates of *Enterococcus faecium*: a 22-year review at one institution. *Antimicrob. Agents Chemother.* **35**:2180–2184.
9. Hanrahan, J., C. Høyen, and L. B. Rice. 2000. Geographic distribution of a large mobile element that transfers ampicillin and vancomycin resistance between *Enterococcus faecium* strains. *Antimicrob. Agents Chemother.* **44**:1349–1351.
10. Morris, J. G., Jr., D. K. Shay, J. N. Hebden, R. J. McCarter, Jr., B. E. Perdue, W. Jarvis, J. A. Johnson, T. C. Dowling, L. B. Polish, and R. S. Schwalbe. 1995. Enterococci resistant to multiple antimicrobial agents, including vancomycin: establishment of endemicity in a university medical center. *Ann. Int. Med.* **123**:250–259.
11. Rice, L. B., S. Bellais, L. L. Carias, R. Hutton-Thomas, R. A. Bonomo, P. Caspers, M. G. Page, and L. Gutmann. 2004. Impact of specific *pbp5* mutations on expression of  $\beta$ -lactam resistance in *Enterococcus faecium*. *Antimicrob. Agents Chemother.* **48**:3028–3032.
- 11a. Rice, L. B., L. L. Carias, S. Marshall, S. B. D. Rudin, and R. Hutton-Thomas. 2005. Tn5386, and novel Tn916-like mobile element in *Enterococcus faecium* D344R that interacts with Tn916 to yield a large genomic deletion. *J. Bacteriol.* **187**:6668–6677.
12. Rice, L. B., and L. L. Carias. 1998. Transfer of Tn5385, a composite, multi-resistance element from *Enterococcus faecalis*. *J. Bacteriol.* **180**:714–721.
13. Rice, L. B., L. L. Carias, R. Hutton-Thomas, F. Sifaoui, L. Gutmann, and S. D. Rudin. 2001. Penicillin-binding protein 5 and expression of ampicillin resistance in *Enterococcus faecium*. *Antimicrob. Agents Chemother.* **45**:1480–1486.
14. Rice, L. B., S. H. Marshall, and L. L. Carias. 1992. Tn5381, a conjugative transposon identifiable as a circular form in *Enterococcus faecalis*. *J. Bacteriol.* **174**:7308–7315.
15. Rybkine, T., J.-L. Mainardi, W. Sougakoff, E. Collatz, and L. Gutmann. 1998. Penicillin-binding protein 5 sequence alterations in clinical isolates of *Enterococcus faecium* with different levels of  $\beta$ -lactam resistance. *J. Infect. Dis.* **178**:159–163.
16. Sahn, D. F., M. K. Marsilio, and G. Piazza. 1999. Antimicrobial resistance in key bloodstream bacterial isolates: electronic surveillance with the Surveillance Network Database—USA. *Clin. Infect. Dis.* **29**:259–263.
17. Schouten, M. A., R. J. Willems, W. A. Kraak, J. Top, J. A. Hoogkamp-Korstanje, and A. Voss. 2001. Molecular analysis of Tn1546-like elements in vancomycin-resistant enterococci isolated from patients in Europe shows geographic transposon type clustering. *Antimicrob. Agents Chemother.* **45**:986–989.
18. Thorisdottir, A. S., L. L. Carias, S. H. Marshall, M. Green, M. J. Zervos, C. Giorgio, L. A. Mermel, J. M. Boyce, A. A. Medeiros, H. Fraimow, and L. B. Rice. 1994. IS6770, an enterococcal insertion-like element useful for determining the clonal relationship of clinical enterococcal isolates. *J. Infect. Dis.* **170**:1539–1548.
19. Welton, L. A., L. A. Thal, M. B. Perri, S. Donabedian, J. McMahon, J. W. Chow, and M. J. Zervos. 1998. Antimicrobial resistance in enterococci isolated from turkey flocks fed virginiamycin. *Antimicrob. Agents Chemother.* **42**:705–708.
20. Williamson, R., C. LaBouguenec, L. Gutmann, and T. Horaud. 1985. One or two low affinity penicillin-binding proteins may be responsible for the range of susceptibility of *Enterococcus faecium* to penicillin. *J. Gen. Microbiol.* **131**:1933–1940.
21. Zorzi, W., X. Y. Zhou, O. Dardenne, J. Lamotte, D. Raze, J. Pierre, L. Gutmann, and J. Coyette. 1996. Structure of the low-affinity penicillin-binding protein 5 PBP5fm in wild-type and highly penicillin-resistant strains of *Enterococcus faecium*. *J. Bacteriol.* **178**:4948–4957.