

MINIREVIEW

Aminoglycoside Resistance in *Pseudomonas aeruginosa*

Keith Poole*

Department of Microbiology & Immunology, Queen's University, Kingston, Ontario, Canada

Aminoglycosides (35) are a vital component of antipseudomonal chemotherapy implicated in the treatment of a variety of infections (9, 45), particularly pulmonary infections in cystic fibrosis (CF) patients (22). These agents are bactericidal and exhibit synergy with other antimicrobials, most notably β -lactams, with which they are often administered for the treatment of *Pseudomonas aeruginosa* infections; and problems with toxicity (aminoglycosides are oto- and nephrotoxic) appear to be ameliorated by increasing the dosing intervals (157a) and, in lung infections at least, through the use of aerosolized agents (e.g., tobramycin) (22). Resistance to aminoglycosides has, however, been known for some time, with reports from the 1960s highlighting the general insusceptibilities of *P. aeruginosa* clinical isolates to, e.g., kanamycin (50, 51). Today, resistance to aminoglycosides with antipseudomonal activities, including gentamicin and tobramycin, but also amikacin, is also all too common and is present in virtually all areas of the world, but particularly in Europe and Latin America (Table 1). Such resistance is seen in respiratory isolates (96), particularly isolates from CF patients (57, 85, 115, 140), as well as bloodstream (86), urinary (12), wound (65), burn (36, 166), eye (2, 20), and aural (27, 128) isolates (Table 1). Resistance typically results from drug inactivation by plasmid- or chromosome-encoded enzymes harbored by resistant strains, although enzyme-independent resistance resultant from defects in uptake and accumulation (dubbed impermeability resistance) is also commonplace, particularly in isolates from CF patients (99–101, 114, 121, 131) and intensive care units (ICUs) (10, 48, 54).

MODIFYING ENZYMES

Inactivation of aminoglycosides, such as streptomycin, kanamycin, neomycin, and gentamicin, by resistant *P. aeruginosa* isolates has been known since the 1960s and 1970s (17, 29, 76, 157). Traditionally, aminoglycoside inactivation in resistant strains involves their modification by enzymes that phosphorylate (aminoglycoside phosphoryltransferase [APH]), acetylate (aminoglycoside acetyltransferase [AAC]), or adenylate (aminoglycoside nucleotidyltransferase [ANT]; also referred to as aminoglycoside adenytransferase) these antimicrobials (see references 6, 152, and 170 for reviews on these modifying enzymes); and such enzymes are common determinants of aminoglycoside resistance in *P. aeruginosa* (32, 37, 74, 98–101,

114, 121, 131, 146) (Table 2). Increasingly, too, individual aminoglycoside-resistant *P. aeruginosa* isolates carry multiple (i.e., two to five) modifying enzymes and exhibit broad-spectrum aminoglycoside resistance as a result (74, 98–101, 127, 131, 146).

AACs. *P. aeruginosa* resistance to aminoglycosides (i.e., gentamicin) owing to enzymatic N-acetylation has been known for some time (17, 135). Acetylation of aminoglycosides can occur at the 1, 3, 6', and 2' amino groups and involves virtually all medically useful compounds (e.g., gentamicin, tobramycin, netilmicin, and amikacin). Enzymes that modify the 3 position (3-N-aminoglycoside acetyltransferases [AAC(3)]) (11) and the 6' position (6'-N-aminoglycoside acetyltransferases [AAC(6')]) (52, 172) were discovered early in *P. aeruginosa* and remain the most common acetyltransferases and, with ANT(2'') (see below), the most common enzymes providing for aminoglycoside resistance in this organism (19, 32, 37, 74, 99–101, 114, 127, 146, 150) (Table 2). The AAC (3)-I family, of which three variants (Ia [154, 169], Ib [142], and Ic [125]) have been described in *P. aeruginosa*, is a common determinant of gentamicin resistance in this organism (3, 31, 32, 114, 142, 143, 150). AAC(3)-II (33, 74, 106, 127) and AAC(3)-III (163) are less commonly described AAC(3) enzymes that determine gentamicin resistance as well as tobramycin and netilmicin resistance [AAC(3)-II] or tobramycin and kanamycin resistance [AAC(3)-III] in *P. aeruginosa*.

The AAC(6') family of enzymes provide resistance to tobramycin, netilmicin, kanamycin, and either amikacin (I subfamily) or gentamicin (II subfamily). AAC(6')-II is not only the most common AAC(6') but also the most common AAC in *P. aeruginosa* (32, 37, 99–101, 114, 127, 145) and is thus a significant determinant of gentamicin and tobramycin resistance in this organism. While AAC(6')-I [also referred to as AAC(6')-Ia] is less common, it is significant for amikacin resistance in *P. aeruginosa* (38, 74, 75, 106, 131), although a variant of this enzyme that fails to provide for amikacin resistance, AAC(6')-Ib, has been reported in clinical isolates resistant to tobramycin (43). A variant of the latter enzyme, AAC(6')-Ib', that differs from AAC(6')-Ib by a single amino acid and that has the same activity as AAC(6')-II has been described in a few CF patient isolates resistant to tobramycin (89). A fused *aac(3)-I-aac(6')-Ib* gene encoding an enzyme active against gentamicin, tobramycin, and kanamycin has also been described (34). Novel AAC(6') enzymes similar in sequence to AAC(6')-I but significantly shorter and dubbed AAC(6')-29a and AAC(6')-29b that provide resistance to all typical AAC(6')-I substrates except netilmicin have been described (118). AAC(6')-29b displays weak acetyltransferase activity, and aminoglycoside resis-

* Mailing address: Department of Microbiology & Immunology, Rm. 737 Botterell Hall, Queen's University, Kingston, ON K7L 3N6, Canada. Phone: (613) 533-6677. Fax: (613) 533-6796. E-mail: poolek@post.queensu.ca.

TABLE 1. Summary of recent studies documenting the incidence of aminoglycoside resistance in clinical isolates of *P. aeruginosa*

Location, study ^a	Year(s)	% of isolates resistant to ^b :			Reference
		AMI	GEN	TOB	
Europe (all), SENTRY	1997-1998	5.1	18.3	17.8	139
North America (urinary), SENTRY	1998	6.7	13.3	8.8	95
Asia-Pacific (all), SENTRY	1997-1999	4.2		10.1	40
Canada (all), SENTRY	1997-1999	2.2		5.8	40
Europe (all), SENTRY	1997-1999	21.1		31.6	40
Latin America (all), SENTRY	1997-1999	30.5		35.8	40
United States (all), SENTRY	1997-1999	3.4		7.8	40
Asia-West Pacific (urinary), SENTRY	1998-1999	8.5	24.5	17.9	158
United Kingdom (all)	1999	5.6 (36) ^c	11.1 (43) ^c		57
Belgium and Luxembourg (nosocomial)	1999	10.5	23.5	19.5	161
Asia-Pacific (all), SENTRY	1997-2000	4.8	15.8	10.4	66
Europe (all), SENTRY	1997-2000	13.7	28.3	24.2	66
Latin America (all), SENTRY	1997-2000	26.8	38.2	34.5	66
North America (all), SENTRY	1997-2000	4.6	15.8	7.8	66
Europe (ICU), MYSTIC	1997-2000		38.9		44
Europe (blood), MYSTIC	1997-2001		26.8		160
Iran (burn)	1998-1999	53.3	90.7		36
United States (nosocomial)	1998-2001	4.7/3.7 ^d	14.7/16.7 ^d		67
Europe (urinary), ESGNI ^e	2000	19.4 (4) ^f	46 (23.8) ^f	35.3 (14.3) ^f	12
Latin America (urinary), SENTRY	2000	51.5	57.6	54.5	41
Latin America (skin and soft tissue), SENTRY	2000	36.2	46.6	43.1	134
Latin America (blood), SENTRY	2000	26.7			133
North America (skin and soft tissue), SENTRY	2000	2	10.5	6.6	122
North America (respiratory), SENTRY	2000	6.3	19.2	9.8	59
France (blood)	2000		27		26
United Kingdom (CF patients)	2000		47	10.1	115
Taiwan (all), TSAR	2000	10	30	25	83
North America and Europe (blood)	2000-2001	19/5.1/15.1/2.7/9.6/7.6 ^g	48.4/20.1/33.9/24.7/21.4/21.3 ^g		167
Latin America (all), SENTRY	2001	34.6	50.4		4
United States (all), MYSTIC	2001		17.8	9.1	124
United States and Europe (skin and soft tissue), TSN	2001	3.6/7.4/1.7/20.4/0.6 ^h	15.5/29.3/8.7/44.9/20.8 ^h		65
United States (blood) TSN—USA	2002		22.8		69
North America (all), MYSTIC	2002		8.4	6.9	108

^a The geographical region included in the indicated studies and the sites of infection of the isolates examined (in parentheses) are highlighted. MYSTIC, Meropenem Yearly Susceptibility Test Information Collection; SENTRY, SENTRY Antimicrobial Surveillance Program; TSAR, Taiwan Surveillance of Antimicrobial Resistance; ESGNI, European Study Group on Nosocomial Infections; TSN, The Surveillance Network.

^b Resistance rates were taken directly from the indicated studies or were calculated accordingly when the studies indicated rates of susceptibility. AMI, amikacin; GEN, gentamicin; TOB, tobramycin.

^c Rates in parentheses are for isolates from CF patients only.

^d Rates are for non-ICU isolates/ICU isolates.

^e The data shown are for a 1-day study.

^f Rates in parentheses are for isolates from European Union countries only.

^g Rates are for France/Germany/Italy/Spain/Canada/the United States.

^h Rates are for the United States/France/Germany/Italy/Spain.

tance appears to result from very tight binding (i.e., sequestering) of aminoglycosides by this enzyme (90).

APHs. Inactivation of aminoglycosides such as kanamycin (28, 135, 159), neomycin (77, 78), and streptomycin (77, 79, 80, 135) by resistant strains of *P. aeruginosa* as a result of phosphorylation has been known for >30 years. Inactivation is carried out by phosphotransferases [APH(3')] that modify the

3'-OH of these antimicrobials, and these phosphotransferases are commonly encountered in *P. aeruginosa* (74, 99, 127). Several APH(3') enzymes have been described in *P. aeruginosa*, with APH(3')-I and -II being predominant in clinical isolates resistant to kanamycin (and neomycin) (99, 123, 131, 146, 177). Indeed, a chromosomal *aphA*-encoded APH(3')-II-type enzyme, APH(3')-IIb (53), is likely responsible for the general

TABLE 2. Mechanisms of aminoglycoside resistance in clinical isolates of *P. aeruginosa*^a

Aminoglycoside resistance mechanism	Resistance to ^b :	Incidence (%) ^c	
		Miller et al. (98) ^d	Miller et al. (99)
AAC(6')-I	T, N, A	1.1–18.8 (6.2)	1.25
AAC(6')-II	G, T, N	2.1–70.3 (32.5)	18.39
AAC(3)-I	G	0.6–31.9 (8.3)	2.05
AAC(3)-II	G, T, N	1.1–55.3 (4.5) ^e	2.20
AAC(3)-III	G, T	0.1–7.1 (3.2)	0.10
AAC(3)-VI	G, T, N	0.2–6.3 (2.6)	0.15
AAC(3)-?	G, N	0.7–11.7 (4.5)	0.60
ANT(2'')-I	G, T	1.7–45.2 (16.9)	11.87
ANT(4')-II	T, A, I		0.05
APH(3')-VI	A, I		0.20
Impermeability	G, T, N, A, I	4.3–23.7 (14.0)	26.15

^a Mechanisms of aminoglycoside resistance in clinical strains and their incidence in the indicated studies are shown. The most prevalent mechanisms are indicated in boldface type.

^b Only the major antipseudomonal aminoglycosides are indicated. T, tobramycin; N, netilmicin; A, amikacin; G, gentamicin; I, isepamicin.

^c The incidence of the indicated resistance mechanisms in aminoglycoside-resistant clinical strains of *P. aeruginosa* when they occur singly. The data do not include instances in which the indicated mechanisms occur in combination with other aminoglycoside resistance mechanisms.

^d The indicated publication summarizes data from several studies, and the numbers presented reflect the range of incidences for each mechanism seen in these individual studies. The numbers in parentheses are the averages for all the studies.

^e The average incidence reported excludes the 55.3% incidence reported for a single small study in Chile.

insensitivity of *P. aeruginosa* to, e.g., kanamycin (112) that was noted in the 1960s, when this drug was first being used clinically (50, 51). Interestingly, a gene, *hpaA*, encoding an AraC-type positive regulator of *aph(3')-Iib* and genes involved in the metabolism of 4-hydroxyphenylacetic acid (4-HPA) occur immediately upstream of the *aph(3')-Iib* gene and form an operon with the *aph(3')-Iib* gene (178). HpaA activation of these genes is stimulated by 4-HPA, suggesting that the phosphotransferase may, in fact, play an intended role in metabolism and only fortuitously provides resistance to aminoglycosides. APH(3') enzymes that provide resistance to other aminoglycosides have also been described in *P. aeruginosa* and include APH(3')-VI (amikacin and isepamicin) (74, 82, 99, 156) and APH(2'') (gentamicin and tobramycin) (74).

ANTs. The adenylation of aminoglycosides such as streptomycin (80) and gentamicin (5, 31) by resistant strains of *P. aeruginosa* has been known for >20 years. The most prevalent nucleotidyltransferase is the ANT(2'')-I enzyme, which, with AAC(6') [and, to some extent, AAC(3)], represents the most common determinant of enzyme-dependent aminoglycoside resistance in *P. aeruginosa* (19, 32, 37, 99–101, 114, 146, 150). The ANT(2'')-I enzyme inactivates gentamicin and tobramycin but not netilmicin or amikacin and is thus found in gentamicin-resistant (19, 31, 114, 123) and tobramycin-resistant (89) clinical isolates. Other adenylation transferases associated with aminoglycoside resistance in *P. aeruginosa* include ANT(3'') (streptomycin resistance) (146) and ANT(4')-II (amikacin, tobramycin, and isepamicin resistance) (63, 132, 147). Two variants of ANT(4')-II, ANT(4')-IIa (63, 147) and ANT(4')-IIb (132), have been reported and are encoded by genes present in

the chromosome and/or on plasmids of amikacin-resistant clinical isolates.

Aminoglycoside-modifying enzymes and mobile genetic elements. R factors encoding aminoglycoside-modifying enzymes and other resistance determinants have been known for some time in *P. aeruginosa* (38, 135), as have transposons that carry genes for aminoglycoside resistance and resistance to other compounds (e.g., sulfonamides and chloramphenicol) (130). Indeed, aminoglycoside-modifying enzymes are often encoded by mobile elements that harbor additional resistance determinants (106). *aac(3)* genes are often associated with transposons (33) and/or integrons (142, 143, 154, 169) in *P. aeruginosa*, including integrons carrying genes for extended-spectrum β -lactamases (34, 119) or metallo- β -lactamases (118, 125, 165), as well as other aminoglycoside-modifying enzymes (118). One reported multidrug-resistant isolate, for example, harbored an integron carrying *aac(3)-Ic*, in addition to genes for the VIM-2 metallo- β -lactamase and the CmlA chloramphenicol efflux pump (125). The presence of these determinants on integrons with multiple resistance genes may explain the multidrug resistance of many aminoglycoside-resistant *P. aeruginosa* isolates. *aac(6')* genes are also often associated with integrons (105, 107, 116, 117) or transposons (43, 145) and may be associated with genes for narrow-spectrum β -lactamases (18), extended-spectrum β -lactamases (105, 107, 116, 117), or metallo- β -lactamases (155, 165) as well as with, again, genes encoding other aminoglycoside-modifying enzymes (18, 118). One *aac(6')-I*-carrying isolate harbored this determinant on a plasmid-borne integron that also contained genes for chloramphenicol (chloramphenicol acetyltransferase), β -lactam (OXA β -lactamase), and gentamicin [ANT(2'')] resistance (18). A recent report (144) of plasmid-encoded amikacin resistance in clinical strains of *P. aeruginosa* may also reflect mobilization of an *aac(6')* gene on, e.g., a transposon or integron. In one report, too, the *aph(3')-VIa* gene of the highly amikacin-resistant strain being studied was present on a transposon (82). As with other aminoglycoside-modifying enzymes, genes for ANT enzymes can be integron associated, particularly the *aadA*-encoded ANT(3'') (9) enzyme that inactivates streptomycin and spectinomycin and that is commonplace on class 1 integrons (72, 73, 109, 143, 151, 162); but they can also be encoded by *aadB*, i.e., ANT(2'')-I (18) and, possibly, ANT(4')-IIb (132). In some instances, too, these occur with other resistance genes, including those encoding β -lactamases (18, 151, 162). The *veb-1* metallo- β -lactamase gene that is typically integron associated can occur together, for example, with genes for both ANT(2'') and ANT(3'') in *P. aeruginosa* (47).

IMPERMEABILITY

Aminoglycoside resistance independent of inactivating enzymes has been known for some time in *P. aeruginosa* (13, 81, 157). Characterized by resistance to all aminoglycosides and often associated with reduced aminoglycoside accumulation (13, 91, 157), such resistance was attributed to reduced uptake owing to reduced permeability and, as such, was typically referred to as impermeability resistance. Numerous studies (74, 99–101, 114, 121, 150) have highlighted the significance of impermeability resistance in aminoglycoside-resistant clinical isolates, particularly in isolates from CF patients (in which it is

often the most common aminoglycoside resistance mechanism [62, 89, 137, 148]), including isolates originally identified as resistant to amikacin (62, 91, 131), gentamicin (13, 114), and tobramycin (89). In some instances, too, impermeability resistance occurs together with inactivating enzymes in promoting multiple aminoglycoside resistance in *P. aeruginosa* (89, 98–101, 127). Reports of an aminoglycoside-resistant variant that was isolated during treatment of experimental endocarditis in rabbits and that lacked enzymes and obvious outer membrane (OM) changes but that showed reduced levels of tobramycin uptake were also suggestive of impermeability resistance (113). Despite earlier conclusions that enzyme-independent resistance associated with reduced accumulation represented impermeability and reduced uptake, more recent studies (153, 164, 168) of such panaminoglycoside-resistant strains indicate that resistance is likely due to efflux (see below).

ADAPTIVE RESISTANCE

The ability to “train” *P. aeruginosa* to grow in the presence of elevated levels of aminoglycosides has been known for some time (61). Characterized by resistance not only to the selecting antimicrobial but to all aminoglycosides and loss of resistance in the absence of drug (46, 61), this reversible panaminoglycoside resistance came to be known as adaptive resistance (71). Shown to occur in vitro (7, 23, 46, 49, 70) and in vivo (8, 24, 171), resistance typically develops within a few hours of first exposure and disappears several hours after removal of the antibiotic. Intriguingly, resistance appears to result from reduced levels of aminoglycoside accumulation (23, 70), reminiscent of impermeability resistance. Indeed, a recent publication confirms the involvement of an aminoglycoside-inducible drug efflux system in the reduced accumulation that characterizes adaptive aminoglycoside resistance (60). Interestingly, adaptively aminoglycoside-resistant *P. aeruginosa* strains also show enhanced expression of genes associated with anaerobic respiration (68). Given the defect in aminoglycoside accumulation seen in anaerobically grown bacteria (138) and the reduced transport capability of *P. aeruginosa* grown with nitrate instead of oxygen as the terminal electron acceptor, it is also possible that reduced uptake also contributes to adaptive aminoglycoside resistance (i.e., aminoglycosides induce anaerobic respiration at the expense of aerobic respiration, and this comprises aminoglycoside uptake [68]).

EFFLUX

Recent studies have clarified the involvement of an efflux system of the resistance-nodulation-division (RND) family (for a review, see reference 120) MexXY (102) (also referred to as AmrAB [168]) in the reduced level of aminoglycoside accumulation that characterizes both impermeability resistance (153, 164, 168) and adaptive aminoglycoside resistance (60) in *P. aeruginosa*. The RND family of pumps is one of five families of drug efflux systems described to date in bacteria (120) and typically consists of three components that include an inner membrane drug-proton antiporter (the RND component), an OM channel-forming protein (the OM factor [OMF]), and a periplasmic link protein (the membrane fusion protein) that joins the other two components (120). MexX and MexY are

the periplasmic and inner membrane proteins, respectively, and are encoded by the *mexXY* operon, while the apparent OMF for this system is OprM (1, 102), the product of the third gene of an operon encoding another three-component RND-type pump, MexAB-OprM (1, 93). Still, the demonstration that mutants lacking one of the OM proteins, OpmG, OpmH, or OpmI, were aminoglycoside hypersusceptible suggests that one or more of these may also function with MexXY, perhaps as the intended OMF for this efflux system (64).

MexXY actually accommodates a range of antimicrobials, including macrolides, tetracyclines, glycolylcyclines, lincomycin, chloramphenicol, novobiocin, fluoroquinolones, and β -lactams (94, 102, 111), although it is implicated only in resistance to aminoglycosides, erythromycin, tetracyclines, and glycolylcyclines in wild-type cells (1, 25), probably because only these agents induce *mexXY* expression (93). Similar three-component RND-type aminoglycoside efflux systems have been described in *Burkholderia pseudomallei* (AmrAB-OprA [103] and BpeAB-OprB [21]) and *Escherichia coli* (AcrAD-TolC [58, 129]). A very modest contribution of MexAB-OprM and a multidrug transporter of the small multidrug resistance family, EmrE_{P.A.}, to aminoglycoside resistance has also been noted (84). Interestingly, an in vitro-isolated mutant that was selected on gentamicin and ofloxacin and that showed increased levels of MexXY expression relative to that of the wild type has been reported (94), although it is unclear if MexXY is also responsible for the fluoroquinolone resistance. Certainly, there are as yet no reports of MexXY-mediated fluoroquinolone resistance in clinical strains (and no indication that strains with impermeability resistance are correspondingly fluoroquinolone resistant). Intriguingly, the well-known divalent cation antagonism of aminoglycosides in *P. aeruginosa* is also dependent on the presence of this efflux system (92).

Expression of *mexXY* is under the control of MexZ (also referred to as AmrR [168]) a repressor of the TetR and AcrR family encoded by a gene located immediately upstream of *mexXY* (1, 168). In vitro-constructed knockout mutations in *mexZ* have been shown to increase the level of *mexXY* expression but did not provide for aminoglycoside resistance (168), and in one study (153) MexXY-expressing aminoglycoside-resistant clinical isolates lacked mutations in *mexZ*, suggesting that aminoglycoside resistance attributable to MexXY may require additional components and other means of upregulating *mexXY*. In addition, while a recent report highlighting the presence of *mexZ* mutations in aminoglycoside-resistant clinical isolates expressing *mexXY* indicates that *mexZ* mutations may play a role in *mexXY* expression in clinical strains (164), it is by no means clear that such mutations were sufficient for aminoglycoside resistance. While the most significant observation regarding the regulation of *mexXY* is its inducibility by several substrate antimicrobials, it is unclear if this is mediated by the MexZ repressor (e.g., drugs target MexZ directly, obviating repressor activity, thereby permitting *mexXY* expression, as has been seen for other drug-inducible efflux systems [141]). Alternatively, *mexXY* expression may be responding to, e.g., the interaction of drugs with their ribosome targets; and, indeed, preliminary studies indicate that for some of these, this may well be the case (K. Boisson, M. L. Sobel, K. Poole, and P. Plesiat, unpublished data).

OUTER MEMBRANE

Several early studies documented an apparent role for the OM protein OprH in aminoglycoside resistance-expression of the protein in certain mutants and in cells grown under Mg^{2+} -limiting conditions correlated with resistance (110) and delayed uptake (55) of, e.g., the aminoglycoside streptomycin. Moreover, loss of this protein in specific mutant constructs subsequently restored susceptibility to these agents (176), further highlighting its contribution to resistance. More recent studies have, however, demonstrated that OprH is actually encoded by the first gene of a three-gene operon that includes *phoP* (which encodes a response regulator) and *phoQ* (which encodes a sensor kinase) (88) and that the low Mg^{2+} concentration-dependent and mutational resistance to aminoglycosides previously attributed to OprH is, in fact, related to PhoP-PhoQ activity (87). Apparently, then, previous *oprH* knockout mutations compromised resistance to aminoglycosides because of polar effects on expression of the downstream *phoPQ* genes (87). PhoP and PhoQ are implicated in resistance to polycationic antimicrobials (e.g., polymyxins) (88) and cationic antimicrobial peptides (87), in part because of their role in promoting an aminoarabinose modification of the lipid A portion of lipopolysaccharide (LPS) (97, 104). While the details of PhoPQ involvement in aminoglycoside resistance remain to be fully elucidated and its involvement in resistance to polycations and aminoglycosides appears to differ, it is possible that PhoPQ-dependent aminoglycoside resistance in *P. aeruginosa* involves a similar modification of LPS (97). Certainly, changes to the LPS component of the OM have long been implicated in resistance to aminoglycosides (15, 39, 149, 175), including resistance in clinical isolates (56). This is, perhaps, not surprising, given that LPS appears to be a necessary target for aminoglycoside binding in the process of its uptake across the OM of *P. aeruginosa* (126, 136).

OTHERS

In addition to the mechanisms already described, there are infrequent reports of ribosomal changes (80) and defects in electron transport that adversely affect aminoglycoside uptake (14, 16), contributing to aminoglycoside resistance in *P. aeruginosa*. Recently, too, clinical isolates of *P. aeruginosa* showing high-level panaminoglycoside resistance (in contrast to impermeability resistance, which is typically low- to moderate-level panaminoglycoside resistance) were shown to carry a gene, *rmtA*, that encodes a 16S rRNA methylase (174). Such an enzyme has been reported previously in aminoglycoside-producing actinomycetes; and although it is rarely seen in clinical pathogens, high-level aminoglycoside resistance owing to a 16S rRNA methylase has been reported in both *Serratia marcescens* (30) and *Klebsiella pneumoniae* (42). Significantly, the *rmtA* gene of *P. aeruginosa* appears to be associated with mobile genetic elements (173), which has important implications vis-à-vis the dissemination of this determinant among pathogenic *Pseudomonas* strains and, possibly, other gram-negative bacteria. Finally, the well-known resistance of biofilm-grown *P. aeruginosa* isolates to multiple antimicrobials, including aminoglycosides, is slowly being elucidated, with the demonstrated anaerobic growth of such organisms (56a) likely responsible, to

some extent at least, for resistance to agents like aminoglycosides, given the apparent negative impact of anaerobic metabolism on aminoglycoside uptake (see above). A locus involved in the synthesis of periplasmic glucans, *ndvB*, has also recently been implicated in *P. aeruginosa* biofilm resistance to several agents, particularly tobramycin (90a). These glucans, which are specifically expressed in biofilm organisms, bind to tobramycin in vitro; and it has been suggested that similar binding in the periplasm of biofilm cells would restrict this agent's passage into the cytoplasm, where its targets lie.

CONCLUSIONS

While aminoglycosides remain useful agents for the treatment of severe *P. aeruginosa* infections, resistance continues to be an issue, particularly in Latin America and parts of Europe. As with any antimicrobial there are geographical variations in resistance rates that likely reflect differences in aminoglycoside prescription patterns and/or the quality of infection control practices, although geographical differences in the occurrence of individual aminoglycoside resistance determinants might also play a role. While the specificity of aminoglycoside-modifying enzymes, historically the major determinants of resistance to these agents, has in the past tended to compromise the use of only selected aminoglycosides, leaving others still effective, the increasing prevalence of strains harboring multiple aminoglycoside-modifying enzymes coupled with panaminoglycoside-exporting efflux systems threatens to compromise the use of this class of agents as a whole. The recent discovery, too, of a genetic determinant (*rmtA*) responsible for high-level panaminoglycoside resistance in *P. aeruginosa* is of great concern, given its probable mobility. Still, as with any agent, the prudent use of aminoglycosides and the use of effective infection control practices can go a long way to limiting the development and spread of aminoglycoside resistance, ensuring that these agents continue to find a place in the treatment of *P. aeruginosa* infections.

ACKNOWLEDGMENT

Work on antibiotic resistance in *P. aeruginosa* in my laboratory is supported by an operating grant from the Canadian Cystic Fibrosis Foundation.

REFERENCES

1. Aires, J. R., T. Köhler, H. Nikaido, and P. Plesiat. 1999. Involvement of an active efflux system in the natural resistance of *Pseudomonas aeruginosa* to aminoglycosides. *Antimicrob. Agents Chemother.* **43**:2624–2628.
2. Alexandrakis, G., E. C. Alfonso, and D. Miller. 2000. Shifting trends in bacterial keratitis in south Florida and emerging resistance to fluoroquinolones. *Ophthalmology* **107**:1497–1502.
3. Alvarez, M., and M. C. Mendoza. 1993. Molecular epidemiology of two genes encoding 3-*N*-aminoglycoside acetyltransferases AAC(3)I and AAC(3)II among gram-negative bacteria from a Spanish hospital. *Eur. J. Epidemiol.* **9**:650–657.
4. Andrade, S. S., R. N. Jones, A. C. Gales, and H. S. Sader. 2003. Increasing prevalence of antimicrobial resistance among *Pseudomonas aeruginosa* isolates in Latin American medical centres: 5 year report of the SENTRY Antimicrobial Surveillance Program (1997–2001). *J. Antimicrob. Chemother.* **52**:140–141.
5. Angelatou, F., S. B. Litsas, and P. Kontomichalou. 1982. Purification and properties of two gentamicin-modifying enzymes, coded by a single plasmid pPK237 originating from *Pseudomonas aeruginosa*. *J. Antibiot. (Tokyo)* **35**:235–244.
6. Azucena, E., and S. Mobashery. 2001. Aminoglycoside-modifying enzymes: mechanisms of catalytic processes and inhibition. *Drug Resist. Update* **4**:106–117.

7. **Barclay, M. L., E. J. Begg, and S. T. Chambers.** 1992. Adaptive resistance following single doses of gentamicin in a dynamic in vitro model. *Antimicrob. Agents Chemother.* **36**:1951–1957.
8. **Barclay, M. L., E. J. Begg, S. T. Chambers, P. E. Thornley, P. K. Pattemore, and K. Grimwood.** 1996. Adaptive resistance to tobramycin in *Pseudomonas aeruginosa* lung infection in cystic fibrosis. *J. Antimicrob. Chemother.* **37**:1155–1164.
9. **Bartlett, J. G.** 2003. 2003–2004 pocket book of infectious disease therapy. Lippincott Williams & Wilkins, Baltimore, Md.
10. **Bert, F., and N. Lambert-Zechovsky.** 1996. Comparative distribution of resistance patterns and serotypes in *Pseudomonas aeruginosa* isolates from intensive care units and other wards. *J. Antimicrob. Chemother.* **37**:809–813.
11. **Biddlecome, S., M. Haas, J. Davies, G. H. Miller, D. F. Rane, and P. J. Daniels.** 1976. Enzymatic modification of aminoglycoside antibiotics: a new 3-N-acetylating enzyme from a *Pseudomonas aeruginosa* isolate. *Antimicrob. Agents Chemother.* **9**:951–955.
12. **Bouza, E., R. San Juan, P. Munoz, A. Voss, J. Kluytmans, et al.** 2001. A European perspective on nosocomial urinary tract infections. I. Report on the microbiology workload, etiology and antimicrobial susceptibility (ES-GNI-003 study). *Clin. Microbiol. Infect.* **7**:523–531.
13. **Bryan, L. E., R. Haraphongse, and H. M. Van den Elzen.** 1976. Gentamicin resistance in clinical-isolates of *Pseudomonas aeruginosa* associated with diminished gentamicin accumulation and no detectable enzymatic modification. *J. Antibiot. (Tokyo)* **29**:743–753.
14. **Bryan, L. E., T. Nicas, B. W. Holloway, and C. Crowther.** 1980. Aminoglycoside-resistant mutation of *Pseudomonas aeruginosa* defective in cytochrome c552 and nitrate reductase. *Antimicrob. Agents Chemother.* **17**:71–79.
15. **Bryan, L. E., K. O'Hara, and S. Wong.** 1984. Lipopolysaccharide changes in impermeability-type aminoglycoside resistance in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **26**:250–255.
16. **Bryan, L. E., and H. M. Van den Elzen.** 1977. Effects of membrane-energy mutations and cations on streptomycin and gentamicin accumulation by bacteria: a model for entry of streptomycin and gentamicin in susceptible and resistant bacteria. *Antimicrob. Agents Chemother.* **12**:163–177.
17. **Brzezinska, M., R. Benveniste, J. Davies, P. J. Daniels, and J. Weinstein.** 1972. Gentamicin resistance in strains of *Pseudomonas aeruginosa* mediated by enzymatic N-acetylation of the deoxystreptamine moiety. *Biochemistry* **11**:761–765.
18. **Bunny, K. L., R. M. Hall, and H. W. Stokes.** 1995. New mobile gene cassettes containing an aminoglycoside resistance gene, *aacA7*, and a chloramphenicol resistance gene, *catB3*, in an integron in pBWH301. *Antimicrob. Agents Chemother.* **39**:686–693.
19. **Busch-Sorensen, C., M. Sonmezoglu, N. Frimodt-Moller, T. Højbjerg, G. H. Miller, and F. Espersen.** 1996. Aminoglycoside resistance mechanisms in Enterobacteriaceae and *Pseudomonas* spp. from two Danish hospitals: correlation with type of aminoglycoside used. *APMIS* **104**:763–768.
20. **Chalita, M. R., A. L. Hoffing-Lima, A. Paranhos, Jr., P. Schor, and R. Belfort, Jr.** 2004. Shifting trends in in vitro antibiotic susceptibilities for common ocular isolates during a period of 15 years. *Am. J. Ophthalmol.* **137**:43–51.
21. **Chan, Y. Y., T. M. Tan, Y. M. Ong, and K. L. Chua.** 2004. BpeAB-OprB, a multidrug efflux pump in *Burkholderia pseudomallei*. *Antimicrob. Agents Chemother.* **48**:1128–1135.
22. **Cheer, S. M., J. Waugh, and S. Noble.** 2003. Inhaled tobramycin (TOBI): a review of its use in the management of *Pseudomonas aeruginosa* infections in patients with cystic fibrosis. *Drugs* **63**:2501–2520.
23. **Daikos, G. L., G. G. Jackson, V. T. Lolans, and D. M. Livermore.** 1990. Adaptive resistance to aminoglycoside antibiotics from first-exposure down-regulation. *J. Infect. Dis.* **162**:414–420.
24. **Daikos, G. L., V. T. Lolans, and G. G. Jackson.** 1991. First-exposure adaptive resistance to aminoglycoside antibiotics in vivo with meaning for optimal clinical use. *Antimicrob. Agents Chemother.* **35**:117–123.
25. **Dean, C. R., M. A. Visalli, S. J. Projan, P.-E. Sum, and P. A. Bradford.** 2003. Efflux-mediated resistance to tigecycline (GAR-936) in *Pseudomonas aeruginosa* PAO1. *Antimicrob. Agents Chemother.* **47**:972–978.
26. **Decusser, J. W., P. Pina, F. Picot, C. Delalande, B. Pangon, P. Courvalin, P. Allouch, and The ColBVH Study Group.** 2003. Frequency of isolation and antimicrobial susceptibility of bacterial pathogens isolated from patients with bloodstream infections: a French prospective national survey. *J. Antimicrob. Chemother.* **51**:1213–1222.
27. **Dohar, J. E., M. A. Kenna, and R. M. Wadowsky.** 1996. In vitro susceptibility of aural isolates of *Pseudomonas aeruginosa* to commonly used ototoxic antibiotics. *Am. J. Otol.* **17**:207–209.
28. **Doi, O., S. Kondo, N. Tanaka, and H. Umezawa.** 1969. Purification and properties of kanamycin-phosphorylating enzyme from *Pseudomonas aeruginosa*. *J. Antibiot. (Tokyo)* **22**:273–282.
29. **Doi, O., M. Ogura, N. Tanaka, and H. Umezawa.** 1968. Inactivation of kanamycin, neomycin, and streptomycin by enzymes obtained in cells of *Pseudomonas aeruginosa*. *Appl. Microbiol.* **16**:1276–1281.
30. **Doi, Y., K. Yokoyama, K. Yamane, J. Wachino, N. Shibata, T. Yagi, K. Shibayama, H. Kato, and Y. Arakawa.** 2004. Plasmid-mediated 16S rRNA methylase in *Serratia marcescens* conferring high-level resistance to aminoglycosides. *Antimicrob. Agents Chemother.* **48**:491–496.
31. **Dornbusch, K., and H. O. Hallander.** 1980. Gentamicin resistance in gram-negative bacilli: occurrence of modifying enzymes and their influence on susceptibility testing. *Scand. J. Infect. Dis.* **12**:295–302.
32. **Dornbusch, K., G. H. Miller, R. S. Hare, K. J. Shaw, et al.** 1990. Resistance to aminoglycoside antibiotics in gram-negative bacilli and staphylococci isolated from blood. Report from a European collaborative study. *J. Antimicrob. Chemother.* **26**:131–144.
33. **Dubois, V., C. Arpin, P. Noury, and C. Quentin.** 2002. Clinical strain of *Pseudomonas aeruginosa* carrying a *bla*_{TEM-21} gene located on a chromosomal interrupted TnA type transposon. *Antimicrob. Agents Chemother.* **46**:3624–3626.
34. **Dubois, V., L. Poirel, C. Marie, C. Arpin, P. Nordmann, and C. Quentin.** 2002. Molecular characterization of a novel class 1 integron containing *bla*_{GES-1} and a fused product of *aac3-Ib/aac6'-Ib'* gene cassettes in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **46**:638–645.
35. **Edson, R. S., and C. L. Terrell.** 1999. The aminoglycosides. *Mayo Clin. Proc.* **74**:519–528.
36. **Estahbanati, H. K., P. P. Kashani, and F. Ghanaatpisheh.** 2002. Frequency of *Pseudomonas aeruginosa* serotypes in burn wound infections and their resistance to antibiotics. *Burns* **28**:340–348.
37. **European Study Group on Antibiotic Resistance (ESGAR).** 1987. In vitro susceptibility to aminoglycoside antibiotics in blood and urine isolates consecutively collected in twenty-nine European laboratories. *Eur. J. Clin. Microbiol.* **6**:378–385.
38. **Falkiner, F. R., G. A. Jacoby, C. T. Keane, and S. R. McCann.** 1982. Amikacin, gentamicin and tobramycin resistant *Pseudomonas aeruginosa* in a leukaemic ward. *Epidemiology and genetic studies.* *J. Hosp. Infect.* **3**:253–261.
39. **Galbraith, L., S. G. Wilkinson, N. J. Legakis, V. Genimata, T. A. Katsorichis, and E. T. Rietschel.** 1984. Structural alterations in the envelope of a gentamicin-resistant rough mutant of *Pseudomonas aeruginosa*. *Ann. Microbiol. (Paris)* **135B**:121–136.
40. **Gales, A. C., R. N. Jones, J. Turnidge, R. Rennie, and R. Ramphal.** 2001. Characterization of *Pseudomonas aeruginosa* isolates: occurrence rates, antimicrobial susceptibility patterns, and molecular typing in the global SENTRY Antimicrobial Surveillance Program, 1997–1999. *Clin. Infect. Dis.* **32**(Suppl. 2):S146–S155.
41. **Gales, A. C., H. S. Sader, and R. N. Jones.** 2002. Urinary tract infection trends in Latin American hospitals: report from the SENTRY Antimicrobial Surveillance Program (1997–2000). *Diagn. Microbiol. Infect. Dis.* **44**:289–299.
42. **Galimand, M., P. Courvalin, and T. Lambert.** 2003. Plasmid-mediated high-level resistance to aminoglycosides in *Enterobacteriaceae* due to 16S rRNA methylation. *Antimicrob. Agents Chemother.* **47**:2565–2571.
43. **Galimand, M., T. Lambert, G. Gerbaud, and P. Courvalin.** 1993. Characterization of the *aac(6')-Ib* gene encoding an aminoglycoside 6'-N-acetyltransferase in *Pseudomonas aeruginosa* BM2656. *Antimicrob. Agents Chemother.* **37**:1456–1462.
44. **Garcia-Rodriguez, J. A., and R. N. Jones.** 2002. Antimicrobial resistance in gram-negative isolates from European intensive care units: data from the Meropenem Yearly Susceptibility Test Information Collection (MYSTIC) programme. *J. Chemother.* **14**:25–32.
45. **Gilbert, D. N., Jr., R. C. Moellering, and M. A. Sande.** 2003. The Sanford guide to antimicrobial therapy 2003. Antimicrobial Therapy, Inc., Hyde Park, N.Y.
46. **Gilleland, L. B., H. E. Gilleland, J. A. Gibson, and F. R. Champlin.** 1989. Adaptive resistance to aminoglycoside antibiotics in *Pseudomonas aeruginosa*. *J. Med. Microbiol.* **29**:41–50.
47. **Girlich, D., T. Naas, A. Leelaporn, L. Poirel, M. Fennewald, and P. Nordmann.** 2002. Nosocomial spread of the integron-located *veb-1*-like cassette encoding an extended-spectrum β -lactamase in *Pseudomonas aeruginosa* in Thailand. *Clin. Infect. Dis.* **34**:603–611.
48. **Goossens, H.** 2003. Susceptibility of multi-drug-resistant *Pseudomonas aeruginosa* in intensive care units: results from the European MYSTIC study group. *Clin. Microbiol. Infect.* **9**:980–983.
49. **Gould, I. M., K. Milne, G. Harvey, and C. Jason.** 1991. Ionic binding, adaptive resistance and post-antibiotic effect of netilmicin and ciprofloxacin. *J. Antimicrob. Chemother.* **27**:741–748.
50. **Griffith, L. J.** 1966. Development of resistance to kanamycin. *Ann. N. Y. Acad. Sci.* **132**:796–799.
51. **Griffith, L. J., W. E. Ostrander, and Z. F. Smith.** 1960. A comparison of the in vitro effectiveness of kanamycin and five other antibacterial agents against common gram-negative rod pathogens. *Antibiot. Chemother.* **10**:88–92.
52. **Haas, M., S. Biddlecome, J. Davies, C. E. Luce, and P. J. Daniels.** 1976. Enzymatic modification of aminoglycoside antibiotics: a new 6'-N-acetylating enzyme from a *Pseudomonas aeruginosa* isolate. *Antimicrob. Agents Chemother.* **9**:945–950.
53. **Hachler, H., P. Santanam, and F. H. Kaysers.** 1996. Sequence and charac-

- terization of a novel chromosomal aminoglycoside phosphotransferase gene, *aph(3')-Iib*, in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **40**:1254–1256.
54. Hanberger, H., D. Diekema, A. Fluit, R. Jones, M. Struelens, R. Spencer, and M. Wolff. 2001. Surveillance of antibiotic resistance in European ICUs. *J. Hosp. Infect.* **48**:161–176.
 55. Hancock, R. E., V. J. Raffle, and T. I. Nicas. 1981. Involvement of the outer membrane in gentamicin and streptomycin uptake and killing in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **19**:777–785.
 56. Hasegawa, M., I. Kobayashi, T. Saika, and M. Nishida. 1997. Drug-resistance patterns of clinical isolates of *Pseudomonas aeruginosa* with reference to their lipopolysaccharide compositions. *Chemotherapy (Basel)* **43**:323–331.
 - 56a. Hassett, D. J., J. Cuppoletti, B. Trapnell, S. V. Lymar, J. J. Rowe, Y. S. Sun, G. M. Hilliard, K. Parvatiyar, M. C. Kamani, D. J. Wozniak, S. H. Hwang, T. R. McDermott, and U. A. Ochsner. 2002. Anaerobic metabolism and quorum sensing by *Pseudomonas aeruginosa* biofilms in chronically infected cystic fibrosis airways, rethinking antibiotic treatment strategies and drug targets. *Adv. Drug Deliv. Rev.* **54**:1425–1443.
 57. Henwood, C. J., D. M. Livermore, D. James, and M. Warner. 2001. Antimicrobial susceptibility of *Pseudomonas aeruginosa*: results of a UK survey and evaluation of the British Society for Antimicrobial Chemotherapy disc susceptibility test. *J. Antimicrob. Chemother.* **47**:789–799.
 58. Hirakawa, H., K. Nishino, T. Hirata, and A. Yamaguchi. 2003. Comprehensive studies of drug resistance mediated by overexpression of response regulators of two-component signal transduction systems in *Escherichia coli*. *J. Bacteriol.* **185**:1851–1856.
 59. Hoban, D. J., D. J. Biedenbach, A. H. Mutnick, and R. N. Jones. 2003. Pathogen of occurrence and susceptibility patterns associated with pneumonia in hospitalized patients in North America: results of the SENTRY Antimicrobial Surveillance Study (2000). *Diagn. Microbiol. Infect. Dis.* **45**:279–285.
 60. Hocquet, D., C. Vogne, F. El Garch, A. Vejux, N. Gotoh, A. Lee, O. Lomovskaya, and P. Plesiat. 2003. MexXY-OprM efflux pump is necessary for adaptive resistance of *Pseudomonas aeruginosa* to aminoglycosides. *Antimicrob. Agents Chemother.* **47**:1371–1375.
 61. Houang, E. T., and D. Greenwood. 1977. Aminoglycoside cross-resistance patterns of gentamicin-resistant bacteria. *J. Clin. Pathol.* **30**:738–744.
 62. Hurley, J. C., G. H. Miller, and A. L. Smith. 1995. Mechanism of amikacin resistance in *Pseudomonas aeruginosa* isolates from patients with cystic fibrosis. *Diagn. Microbiol. Infect. Dis.* **22**:331–336.
 63. Jacoby, G. A., M. J. Blaser, P. Santanam, H. Hachler, F. H. Kayser, R. S. Hare, and G. H. Miller. 1990. Appearance of amikacin and tobramycin resistance due to 4'-aminoglycoside nucleotidyltransferase [ANT(4')-II] in gram-negative pathogens. *Antimicrob. Agents Chemother.* **34**:2381–2386.
 64. Jo, J. T., F. S. Brinkman, and R. E. Hancock. 2003. Aminoglycoside efflux in *Pseudomonas aeruginosa*: involvement of novel outer membrane proteins. *Antimicrob. Agents Chemother.* **47**:1101–1111.
 65. Jones, M. E., J. A. Karlowsky, D. C. Draghi, C. Thornsberry, D. F. Sahn, and D. Nathwani. 2003. Epidemiology and antibiotic susceptibility of bacteria causing skin and soft tissue infections in the USA and Europe: a guide to appropriate antimicrobial therapy. *Int. J. Antimicrob. Agents* **22**:406–419.
 66. Jones, R. N., J. T. Kirby, M. L. Beach, D. J. Biedenbach, and M. A. Pfaller. 2002. Geographic variations in activity of broad-spectrum β -lactams against *Pseudomonas aeruginosa*: summary of the worldwide SENTRY Antimicrobial Surveillance Program (1997–2000). *Diagn. Microbiol. Infect. Dis.* **43**:239–243.
 67. Karlowsky, J. A., D. C. Draghi, M. E. Jones, C. Thornsberry, I. R. Friedland, and D. F. Sahn. 2003. Surveillance for antimicrobial susceptibility among clinical isolates of *Pseudomonas aeruginosa* and *Acinetobacter baumannii* from hospitalized patients in the United States, 1998 to 2001. *Antimicrob. Agents Chemother.* **47**:1681–1688.
 68. Karlowsky, J. A., D. J. Hoban, S. A. Zelenitsky, and G. G. Zhanel. 1997. Altered *denA* and *anr* gene expression in aminoglycoside adaptive resistance in *Pseudomonas aeruginosa*. *J. Antimicrob. Chemother.* **40**:371–376.
 69. Karlowsky, J. A., M. E. Jones, D. C. Draghi, C. Thornsberry, D. F. Sahn, and G. A. Volturo. 2004. Prevalence and antimicrobial susceptibilities of bacteria isolated from blood cultures of hospitalized patients in the United States in 2002. *Ann. Clin. Microbiol. Antimicrob.* **3**:7.
 70. Karlowsky, J. A., M. H. Saunders, G. A. Harding, D. J. Hoban, and G. G. Zhanel. 1996. In vitro characterization of aminoglycoside adaptive resistance in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **40**:1387–1393.
 71. Karlowsky, J. A., S. A. Zelenitsky, and G. G. Zhanel. 1997. Aminoglycoside adaptive resistance. *Pharmacotherapy* **17**:549–555.
 72. Kazama, H., K. Kizu, M. Iwasaki, H. Hamashima, M. Sasatsu, and T. Arai. 1995. Isolation and structure of a new integron that includes a streptomycin resistance gene from the R plasmid of *Pseudomonas aeruginosa*. *FEMS Microbiol. Lett.* **134**:137–141.
 73. Kazama, H., K. Kizu, M. Iwasaki, H. Hamashima, M. Sasatsu, and T. Arai. 1996. A new gene, *aadA2b*, encoding an aminoglycoside adenyltransferase, AAD(3')(9), isolated from integron InC in *Pseudomonas aeruginosa*. *Microbios* **86**:77–83.
 74. Kettner, M., P. Milosovic, M. Hletkova, and J. Kallova. 1995. Incidence and mechanisms of aminoglycoside resistance in *Pseudomonas aeruginosa* serotype O11 isolates. *Infection* **23**:380–383.
 75. Kettner, M., J. Navarova, G. Lebek, and V. Krcmery. 1984. Enzymatic modification of aminoglycoside antibiotics in gentamicin-resistant gram-negative bacteria. *Zentbl. Bakteriol. Mikrobiol. Hyg. Reihe A* **257**:372–382.
 76. Kobayashi, F., M. Yamaguchi, and S. Mitsuhashi. 1971. Inactivation of dihydrostreptomycin by *Pseudomonas aeruginosa*. *Jpn. J. Microbiol.* **15**:381–382.
 77. Kobayashi, F., M. Yamaguchi, and S. Mitsuhashi. 1971. Phosphorylated inactivation of aminoglycosidic antibiotics by *Pseudomonas aeruginosa*. *Jpn. J. Microbiol.* **15**:265–272.
 78. Kobayashi, F., M. Yamaguchi, and S. Mitsuhashi. 1972. Activity of lividomycin against *Pseudomonas aeruginosa*: its inactivation by phosphorylation induced by resistant strains. *Antimicrob. Agents Chemother.* **1**:17–21.
 79. Kobayashi, F., M. Yamaguchi, J. Sato, and S. Mitsuhashi. 1972. Purification and properties of dihydrostreptomycin-phosphorylating enzyme from *Pseudomonas aeruginosa*. *Jpn. J. Microbiol.* **16**:15–19.
 80. Kono, M., and K. O'Hara. 1976. Mechanisms of streptomycin (SM)-resistance of highly SM-resistant *Pseudomonas aeruginosa* strains. *J. Antibiot. (Tokyo)* **29**:169–175.
 81. Kono, M., and K. O'Hara. 1977. Kanamycin-resistance mechanism of *Pseudomonas aeruginosa* governed by an R-plasmid independently of inactivating enzymes. *J. Antibiot. (Tokyo)* **30**:688–690.
 82. Lambert, T., G. Gerbaud, and P. Courvalin. 1994. Characterization of transposon Tn1528, which confers amikacin resistance by synthesis of aminoglycoside 3'-O-phosphotransferase type VI. *Antimicrob. Agents Chemother.* **38**:702–706.
 83. Lauderdale, T. L., M. L. Clifford, Y. R. Shiau, P. C. Chen, H. Y. Wang, J. F. Lai, and M. Ho. 2004. The status of antimicrobial resistance in Taiwan among gram-negative pathogens: the Taiwan surveillance of antimicrobial resistance (TSAR) program, 2000. *Diagn. Microbiol. Infect. Dis.* **48**:211–219.
 84. Li, X.-Z., K. Poole, and H. Nikaido. 2003. Contributions of MexAB-OprM and an EmrE homologue to intrinsic resistance of *Pseudomonas aeruginosa* to aminoglycosides and dyes. *Antimicrob. Agents Chemother.* **47**:27–33.
 85. Livermore, D. M. 2002. Multiple mechanisms of antimicrobial resistance in *Pseudomonas aeruginosa*: our worst nightmare? *Clin. Infect. Dis.* **34**:634–640.
 86. Lyytikäinen, O., V. Golovanova, E. Kolho, P. Ruutu, A. Sivonen, L. Tiitonen, M. Hakanen, and J. Vuopio-Varkila. 2001. Outbreak caused by tobramycin-resistant *Pseudomonas aeruginosa* in a bone marrow transplantation unit. *Scand. J. Infect. Dis.* **33**:445–449.
 87. Macfarlane, E. L., A. Kwasnicka, and R. E. Hancock. 2000. Role of *Pseudomonas aeruginosa* PhoP-phoQ in resistance to antimicrobial cationic peptides and aminoglycosides. *Microbiology* **146**:2543–2554.
 88. Macfarlane, E. L., A. Kwasnicka, M. M. Ochs, and R. E. Hancock. 1999. PhoP-PhoQ homologues in *Pseudomonas aeruginosa* regulate expression of the outer-membrane protein OprH and polymyxin B resistance. *Mol. Microbiol.* **34**:305–316.
 89. MacLeod, D. L., L. E. Nelson, R. M. Shawar, B. B. Lin, L. G. Lockwood, J. E. Dirk, G. H. Miller, J. L. Burns, and R. L. Garber. 2000. Aminoglycoside-resistance mechanisms for cystic fibrosis *Pseudomonas aeruginosa* isolates are unchanged by long-term, intermittent, inhaled tobramycin treatment. *J. Infect. Dis.* **181**:1180–1184.
 90. Magnet, S., T. A. Smith, R. Zheng, P. Nordmann, and J. S. Blanchard. 2003. Aminoglycoside resistance resulting from tight drug binding to an altered aminoglycoside acetyltransferase. *Antimicrob. Agents Chemother.* **47**:1577–1583.
 - 90a. Mah, T. F., B. Pitts, B. Pellock, G. C. Walker, P. S. Stewart, and G. A. O'Toole. 2003. A genetic basis for *Pseudomonas aeruginosa* biofilm antibiotic resistance. *Nature* **426**:306–310.
 91. Maloney, J., D. Rimland, D. S. Stephens, P. Terry, and A. M. Whitney. 1989. Analysis of amikacin-resistant *Pseudomonas aeruginosa* developing in patients receiving amikacin. *Arch. Intern. Med.* **149**:630–634.
 92. Mao, W., M. S. Warren, A. Lee, A. Mistry, and O. Lomovskaya. 2001. MexXY-OprM efflux pump is required for antagonism of aminoglycosides by divalent cations in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **45**:2001–2007.
 93. Masuda, N., E. Sakagawa, S. Ohya, N. Gotoh, H. Tsujimoto, and T. Nishino. 2000. Contribution of the MexX-MexY-OprM efflux system to intrinsic resistance in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **44**:2242–2246.
 94. Masuda, N., E. Sakagawa, S. Ohya, N. Gotoh, H. Tsujimoto, and T. Nishino. 2000. Substrate specificities of MexAB-OprM, MexCD-OprJ, and MexXY-OprM efflux pumps in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **44**:3322–3327.
 95. Mathai, D., R. N. Jones, and M. A. Pfaller. 2001. Epidemiology and frequency of resistance among pathogens causing urinary tract infections in 1,510 hospitalized patients: a report from the SENTRY Antimicrobial

- Surveillance Program (North America). *Diagn. Microbiol. Infect. Dis.* **40**:129–136.
96. Mathai, D., M. T. Lewis, K. C. Kugler, M. A. Pfaller, and R. N. Jones. 2001. Antibacterial activity of 41 antimicrobials tested against over 2773 bacterial isolates from hospitalized patients with pneumonia. I. Results from the SENTRY Antimicrobial Surveillance Program (North America, 1998). *Diagn. Microbiol. Infect. Dis.* **39**:105–116.
 97. McPhee, J. B., S. Lewenza, and R. E. Hancock. 2003. Cationic antimicrobial peptides activate a two-component regulatory system, PmrA-PmrB, that regulates resistance to polymyxin B and cationic antimicrobial peptides in *Pseudomonas aeruginosa*. *Mol. Microbiol.* **50**:205–217.
 98. Miller, G. H., F. J. Sabatelli, R. S. Hare, Y. Glupczynski, P. Mackey, D. Shlaes, K. Shimizu, K. J. Shaw, et al. 1997. The most frequent aminoglycoside resistance mechanisms—changes with time and geographic area: a reflection of aminoglycoside usage patterns? *Clin. Infect. Dis.* **24**(Suppl. 1):S46–S62.
 99. Miller, G. H., F. J. Sabatelli, L. Naples, R. S. Hare, K. J. Shaw, et al. 1994. Resistance to aminoglycosides in *Pseudomonas*. *Trends Microbiol.* **2**:347–353.
 100. Miller, G. H., F. J. Sabatelli, L. Naples, R. S. Hare, K. J. Shaw, et al. 1995. The changing nature of aminoglycoside resistance mechanisms and the role of isepamicin—a new broad-spectrum aminoglycoside. *J. Chemother.* **7**(Suppl. 2):31–44.
 101. Miller, G. H., F. J. Sabatelli, L. Naples, R. S. Hare, K. J. Shaw, et al. 1995. The most frequently occurring aminoglycoside resistance mechanisms—combined results of surveys in eight regions of the world. *J. Chemother.* **7**(Suppl. 2):17–30.
 102. Mine, T., Y. Morita, A. Kataoka, T. Mitzushima, and T. Tsuchiya. 1999. Expression in *Escherichia coli* of a new multidrug efflux pump, MexXY, from *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **43**:415–417.
 103. Moore, R. A., D. DeShazer, S. Reckseidler, A. Weissman, and D. E. Woods. 1999. Efflux-mediated aminoglycoside and macrolide resistance in *Burkholderia pseudomallei*. *Antimicrob. Agents Chemother.* **43**:465–470.
 104. Moskowitz, S. M., R. K. Ernst, and S. I. Miller. 2004. PmrAB, a two-component regulatory system of *Pseudomonas aeruginosa* that modulates resistance to cationic antimicrobial peptides and addition of aminoarabino- to lipid A. *J. Bacteriol.* **186**:575–579.
 105. Mugnier, P., I. Casin, A. T. Bouthors, and E. Collatz. 1998. Novel OXA-10-derived extended-spectrum β -lactamases selected in vivo or in vitro. *Antimicrob. Agents Chemother.* **42**:3113–3116.
 106. Mugnier, P., P. Dubrous, I. Casin, G. Arlet, and E. Collatz. 1996. A TEM-derived extended-spectrum β -lactamase in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **40**:2488–2493.
 107. Mugnier, P., I. Podglajen, F. W. Goldstein, and E. Collatz. 1998. Carbapenems as inhibitors of OXA-13, a novel, integron-encoded β -lactamase in *Pseudomonas aeruginosa*. *Microbiology* **144**:1021–1031.
 108. Mutnick, A. H., P. R. Rhomberg, H. S. Sader, and R. N. Jones. 2004. Antimicrobial usage and resistance trend relationships from the MYSTIC Programme in North America (1999–2001). *J. Antimicrob. Chemother.* **53**:290–296.
 109. Naas, T., L. Poirel, and P. Nordmann. 1999. Molecular characterisation of In51, a class 1 integron containing a novel aminoglycoside adenyltransferase gene cassette, aadA6, in *Pseudomonas aeruginosa*. *Biochim. Biophys. Acta* **1489**:445–451.
 110. Nicas, T. L., and R. E. W. Hancock. 1980. Outer membrane protein H1 of *Pseudomonas aeruginosa*: involvement in adaptive and mutational resistance to ethylenediaminetetraacetate, polymyxin B, and gentamicin. *J. Bacteriol.* **143**:872–878.
 111. Okamoto, K., N. Gotoh, and T. Nishino. 2002. Alterations of susceptibility of *Pseudomonas aeruginosa* by overproduction of multidrug efflux systems, MexAB-OprM, MexCD-OprJ, and MexXY/OprM to carbapenems: substrate specificities of the efflux systems. *J. Infect. Chemother.* **8**:371–373.
 112. Okii, M., S. Iyobe, and S. Mitsuhashi. 1983. Mapping of the gene specifying aminoglycoside 3'-phosphotransferase II on the *Pseudomonas aeruginosa* chromosome. *J. Bacteriol.* **155**:643–649.
 113. Parr, T. R., Jr., and A. S. Bayer. 1988. Mechanisms of aminoglycoside resistance in variants of *Pseudomonas aeruginosa* isolated during treatment of experimental endocarditis in rabbits. *J. Infect. Dis.* **158**:1003–1010.
 114. Phillips, I., A. King, and K. Shannon. 1986. Prevalence and mechanisms of aminoglycoside resistance. A ten-year study. *Am. J. Med.* **80**:48–55.
 115. Pitt, T. L., M. Sparrow, M. Warner, and M. Stefanidou. 2003. Survey of resistance of *Pseudomonas aeruginosa* from UK patients with cystic fibrosis to six commonly prescribed antimicrobial agents. *Thorax* **58**:794–796.
 116. Poirel, L., P. Gerome, C. De Champs, J. Stephanazzi, T. Naas, and P. Nordmann. 2002. Integron-located *oxa-32* gene cassette encoding an extended-spectrum variant of OXA-2 β -lactamase from *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **46**:566–569.
 117. Poirel, L., D. Girlich, T. Naas, and P. Nordmann. 2001. OXA-28, an extended-spectrum variant of OXA-10 β -lactamase from *Pseudomonas aeruginosa* and its plasmid- and integron-located gene. *Antimicrob. Agents Chemother.* **45**:447–453.
 118. Poirel, L., T. Lambert, S. Turkoglu, E. Ronco, J. Gaillard, and P. Nordmann. 2001. Characterization of class 1 integrons from *Pseudomonas aeruginosa* that contain the *bla*_{VM-2} carbapenem-hydrolyzing β -lactamase gene and two novel aminoglycoside resistance gene cassettes. *Antimicrob. Agents Chemother.* **45**:546–552.
 119. Poirel, L., G. F. Weldhagen, C. De Champs, and P. Nordmann. 2002. A nosocomial outbreak of *Pseudomonas aeruginosa* isolates expressing the extended-spectrum β -lactamase GES-2 in South Africa. *J. Antimicrob. Chemother.* **49**:561–565.
 120. Poole, K. 2004. Efflux-mediated multidrug resistance in gram-negative bacteria. *Clin. Microbiol. Infect.* **10**:12–26.
 121. Price, K. E., P. A. Kresel, L. A. Farchione, S. B. Siskin, and S. A. Karpow. 1981. Epidemiological studies of aminoglycoside resistance in the USA. *J. Antimicrob. Chemother.* **8**(Suppl. A):89–105.
 122. Rennie, R. P., R. N. Jones, and A. H. Mutnick. 2003. Occurrence and antimicrobial susceptibility patterns of pathogens isolated from skin and soft tissue infections: report from the SENTRY Antimicrobial Surveillance Program (United States and Canada, 2000). *Diagn. Microbiol. Infect. Dis.* **45**:287–293.
 123. Reynolds, A. V., J. M. Hamilton-Miller, and W. Brumfitt. 1976. In vitro activity of amikacin and ten other aminoglycoside antibiotics against gentamicin-resistant bacterial strains. *J. Infect. Dis.* **134**(Suppl.):S291–S296.
 124. Rhomberg, P. R., R. N. Jones, and H. S. Sader. 2004. Results from the Meropenem Yearly Susceptibility Test Information Collection (MYSTIC) Programme: report of the 2001 data from 15 United States medical centres. *Int. J. Antimicrob. Agents* **23**:52–59.
 125. Riccio, M. L., J. D. Docquier, E. Dell'Amico, F. Luzzaro, G. Amicosante, and G. M. Rossolini. 2003. Novel 3-N-aminoglycoside acetyltransferase gene, *aac(3)-Ic*, from a *Pseudomonas aeruginosa* integron. *Antimicrob. Agents Chemother.* **47**:1746–1748.
 126. Rivera, M., R. E. Hancock, J. G. Sawyer, A. Haug, and E. J. McGroarty. 1988. Enhanced binding of polycationic antibiotics to lipopolysaccharide from an aminoglycoside-supersusceptible, *tolA* mutant strain of *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **32**:649–655.
 127. Rodriguez, E. F., M. M. Gonzalez, L. Z. Gonzalez, F. J. Sabatelli, and M. T. Tedjor Junco. 2000. Aminoglycoside resistance mechanisms in clinical isolates of *Pseudomonas aeruginosa* from the Canary Islands. *Zentbl. Bakteriol. Parasitenkd. Infektkrankh. Hyg. Abt. 1 Orig.* **289**:817–826.
 128. Roland, P. S., and D. W. Stroman. 2002. Microbiology of acute otitis externa. *Laryngoscope* **112**:1166–1177.
 129. Rosenberg, E. Y., D. Ma, and H. Nikaido. 2000. AcrD of *Escherichia coli* is an aminoglycoside efflux pump. *J. Bacteriol.* **182**:1754–1756.
 130. Rubens, C. E., W. F. McNeill, and W. E. Farrar, Jr. 1979. Transposable plasmid deoxyribonucleic acid sequence in *Pseudomonas aeruginosa* which mediates resistance to gentamicin and four other antimicrobial agents. *J. Bacteriol.* **139**:877–882.
 131. Saavedra, S., D. Vera, and C. H. Ramirez-Ronda. 1986. Susceptibility of aerobic gram-negative bacilli to aminoglycosides. Effects of 45 months of amikacin as first-line aminoglycoside therapy. *Am. J. Med.* **80**:65–70.
 132. Sabtcheva, S., M. Galimand, G. Gerbaud, P. Courvalin, and T. Lambert. 2003. Aminoglycoside resistance gene *ant(4')-Iib* of *Pseudomonas aeruginosa* BM4492, a clinical isolate from Bulgaria. *Antimicrob. Agents Chemother.* **47**:1584–1588.
 133. Sader, H. S., R. N. Jones, S. Andrade-Baiocchi, and D. J. Biedenbach. 2002. Four-year evaluation of frequency of occurrence and antimicrobial susceptibility patterns of bacteria from bloodstream infections in Latin American medical centers. *Diagn. Microbiol. Infect. Dis.* **44**:273–280.
 134. Sader, H. S., R. N. Jones, and J. B. Silva. 2002. Skin and soft tissue infections in Latin American medical centers: four-year assessment of the pathogen frequency and antimicrobial susceptibility patterns. *Diagn. Microbiol. Infect. Dis.* **44**:281–288.
 135. Sagai, H., V. Krcmery, K. Hasuda, S. Iyobe, and H. Knothe. 1975. R factor-mediated resistance to aminoglycoside antibiotics in *Pseudomonas aeruginosa*. *Jpn. J. Microbiol.* **19**:427–432.
 136. Saika, T., M. Hasegawa, I. Kobayashi, and M. Nishida. 1999. Ionic binding of ³H-gentamicin and short-time bactericidal activity of gentamicin against *Pseudomonas aeruginosa* isolates with different lipopolysaccharide structures. *Chemotherapy (Basel)* **45**:296–302.
 137. Saiman, L., F. Mehar, W. W. Niu, H. C. Neu, K. J. Shaw, G. Miller, and A. Prince. 1996. Antibiotic susceptibility of multiply resistant *Pseudomonas aeruginosa* isolated from patients with cystic fibrosis, including candidates for transplantation. *Clin. Infect. Dis.* **23**:532–537.
 138. Schlessinger, D. 1988. Failure of aminoglycoside antibiotics to kill anaerobic, low-pH, and resistant cultures. *Clin. Microbiol. Rev.* **1**:54–59.
 139. Schmitz, F. J., J. Verhoef, and A. C. Fluit. 1999. Prevalence of aminoglycoside resistance in 20 European university hospitals participating in the European SENTRY Antimicrobial Surveillance Programme. *Eur. J. Clin. Microbiol. Infect. Dis.* **18**:414–421.
 140. Schulin, T. 2002. In vitro activity of the aerosolized agents colistin and tobramycin and five intravenous agents against *Pseudomonas aeruginosa* isolated from cystic fibrosis patients in southwestern Germany. *J. Antimicrob. Chemother.* **49**:403–406.

141. Schumacher, M. A., and R. G. Brennan. 2002. Structural mechanisms of multidrug recognition and regulation by bacterial multidrug transcription factors. *Mol. Microbiol.* **45**:885–893.
142. Schocho, L. R., C. P. Schaffner, G. H. Miller, R. S. Hare, and K. J. Shaw. 1995. Cloning and characterization of a 3-*N*-aminoglycoside acetyltransferase gene, *aac(3)-Ib*, from *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **39**:1790–1796.
143. Severino, P., and V. D. Magalhaes. 2002. The role of integrons in the dissemination of antibiotic resistance among clinical isolates of *Pseudomonas aeruginosa* from an intensive care unit in Brazil. *Res. Microbiol.* **153**:221–226.
144. Shahid, M., A. Malik, and Sheeba. 2003. Multidrug-resistant *Pseudomonas aeruginosa* strains harbouring R-plasmids and AmpC β -lactamases isolated from hospitalised burn patients in a tertiary care hospital of North India. *FEMS Microbiol. Lett.* **228**:181–186.
145. Shaw, K. J., C. A. Cramer, M. Rizzo, R. Mierzwa, K. Gewain, G. H. Miller, and R. S. Hare. 1989. Isolation, characterization, and DNA sequence analysis of an AAC(6')-II gene from *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **33**:2052–2062.
146. Shaw, K. J., R. S. Hare, F. J. Sabatelli, M. Rizzo, C. A. Cramer, L. Naples, S. Kocsi, H. Munayyer, P. Mann, G. H. Miller, L. Verbist, H. van Landuyt, Y. Glupczynski, M. Catalano, and M. Woloj. 1991. Correlation between aminoglycoside resistance profiles and DNA hybridization of clinical isolates. *Antimicrob. Agents Chemother.* **35**:2253–2261.
147. Shaw, K. J., H. Munayyer, P. N. Rather, R. S. Hare, and G. H. Miller. 1993. Nucleotide sequence analysis and DNA hybridization studies of the *ant(4')-IIa* gene from *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **37**:708–714.
148. Shwar, R. M., D. L. MacLeod, R. L. Garber, J. L. Burns, J. R. Stapp, C. R. Clausen, and S. K. Tanaka. 1999. Activities of tobramycin and six other antibiotics against *Pseudomonas aeruginosa* isolates from patients with cystic fibrosis. *Antimicrob. Agents Chemother.* **43**:2877–2880.
149. Shearer, B. G., and N. J. Legakis. 1985. *Pseudomonas aeruginosa*: evidence for the involvement of lipopolysaccharide in determining outer membrane permeability to carbenicillin and gentamicin. *J. Infect. Dis.* **152**:351–355.
150. Shimizu, K., T. Kumada, W. C. Hsieh, H. Y. Chung, Y. Chong, R. S. Hare, G. H. Miller, F. J. Sabatelli, and J. Howard. 1985. Comparison of aminoglycoside resistance patterns in Japan, Formosa, and Korea, Chile, and the United States. *Antimicrob. Agents Chemother.* **28**:282–288.
151. Sinclair, M. I., and B. W. Holloway. 1982. A chromosomally located transposon in *Pseudomonas aeruginosa*. *J. Bacteriol.* **151**:569–579.
152. Smith, C. A., and E. N. Baker. 2002. Aminoglycoside antibiotic resistance by enzymatic deactivation. *Curr. Drug Targets Infect. Disord.* **2**:143–160.
153. Sobel, M. L., G. A. McKay, and K. Poole. 2003. Contribution of the MexXY multidrug transporter to aminoglycoside resistance in *Pseudomonas aeruginosa* clinical isolates. *Antimicrob. Agents Chemother.* **47**:3202–3207.
154. Tenover, F. C., K. L. Phillips, T. Gilbert, P. Lockhart, P. J. O'Hara, and J. J. Plorde. 1989. Development of a DNA probe from the deoxyribonucleotide sequence of a 3-*N*-aminoglycoside acetyltransferase [AAC(3)-I] resistance gene. *Antimicrob. Agents Chemother.* **33**:551–559.
155. Toleman, M. A., D. Biedenbach, D. Bennett, R. N. Jones, and T. R. Walsh. 2003. Genetic characterization of a novel metallo- β -lactamase gene, bla_{IMP-13}, harboured by a novel Tn5051-type transposon disseminating carbapenemase genes in Europe: report from the SENTRY Worldwide Antimicrobial Surveillance Programme. *J. Antimicrob. Chemother.* **52**:583–590.
156. Torres, C., M. H. Perlin, F. Baquero, D. L. Lerner, and S. A. Lerner. 2000. High-level amikacin resistance in *Pseudomonas aeruginosa* associated with a 3'-phosphotransferase with high affinity for amikacin. *Int. J. Antimicrob. Agents* **15**:257–263.
157. Tseng, J. T., L. E. Bryan, and H. M. Van den Elzen. 1973. Mechanisms and spectrum of streptomycin resistance in a natural population of *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **2**:136–141.
- 157a. Turnidge, J. 2003. Pharmacodynamics and dosing of aminoglycosides. *Infect. Dis. Clin. N. Am.* **17**:503–528.
158. Turnidge, J., J. Bell, D. J. Biedenbach, and R. N. Jones. 2002. Pathogen occurrence and antimicrobial resistance trends among urinary tract infection isolates in the Asia-Western Pacific Region: report from the SENTRY Antimicrobial Surveillance Program, 1998–1999. *Int. J. Antimicrob. Agents* **20**:10–17.
159. Umezawa, H., O. Doi, M. Ogura, S. Kondo, and N. Tanaka. 1968. Phosphorylation and inactivation of kanamycin by *Pseudomonas aeruginosa*. *J. Antibiot. (Tokyo)* **21**:154–155.
160. Unal, S., R. Masterton, and H. Goossens. 2004. Bacteraemia in Europe—antimicrobial susceptibility data from the MYSTIC Surveillance Programme. *Int. J. Antimicrob. Agents* **23**:155–163.
161. Van Eldere, J. 2003. Multicentre surveillance of *Pseudomonas aeruginosa* susceptibility patterns in nosocomial infections. *J. Antimicrob. Chemother.* **51**:347–352.
162. Vezina, G., and R. C. Levesque. 1991. Molecular characterization of the class II multiresistance transposable element Tn1403 from *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **35**:313–321.
163. Vlieghehart, J. S., P. A. Ketelaar-van Gaalen, and J. A. van de Klundert. 1991. Nucleotide sequence of the *aacC3* gene, a gentamicin resistance determinant encoding aminoglycoside-(3)-*N*-acetyltransferase III expressed in *Pseudomonas aeruginosa* but not in *Escherichia coli*. *Antimicrob. Agents Chemother.* **35**:892–897.
164. Vogne, C., J. R. Aires, C. Bailly, D. Hocquet, and P. Plesiat. 2004. Role of the multidrug efflux system MexXY in the emergence of moderate resistance to aminoglycosides among *Pseudomonas aeruginosa* isolates from patients with cystic fibrosis. *Antimicrob. Agents Chemother.* **48**:1676–1680.
165. Walsh, T. R., M. A. Toleman, W. Hryniewicz, P. M. Bennett, and R. N. Jones. 2003. Evolution of an integron carrying bla_{VIM-2} in Eastern Europe: report from the SENTRY Antimicrobial Surveillance Program. *J. Antimicrob. Chemother.* **52**:116–119.
166. Walton, M. A., C. Villarreal, D. N. Herndon, and J. P. Heggers. 1997. The use of aztreonam as an alternate therapy for multi-resistant *Pseudomonas aeruginosa*. *Burns* **23**:225–227.
167. Wenzel, R. P., D. F. Sahn, C. Thornsberry, D. C. Draghi, M. E. Jones, and J. A. Karlowsky. 2003. In vitro susceptibilities of gram-negative bacteria isolated from hospitalized patients in four European countries, Canada, and the United States in 2000–2001 to expanded-spectrum cephalosporins and comparator antimicrobials: implications for therapy. *Antimicrob. Agents Chemother.* **47**:3089–3098.
168. Westbrook-Wadman, S., D. R. Sherman, M. J. Hickey, S. N. Coulter, Y. Q. Zhu, P. Warriner, L. Y. Nguyen, R. M. Shawar, K. R. Folger, and C. K. Stover. 1999. Characterization of a *Pseudomonas aeruginosa* efflux pump contributing to aminoglycoside impermeability. *Antimicrob. Agents Chemother.* **43**:2975–2983.
169. Wohlleben, W., W. Arnold, L. Bissonnette, A. Pelletier, A. Tanguay, P. H. Roy, G. C. Gamboa, G. F. Barry, E. Aubert, and J. Davies. 1989. On the evolution of Tn21-like multiresistance transposons: sequence analysis of the gene (*aacCI*) for gentamicin acetyltransferase-3-I (AAC(3)-I), another member of the Tn21-based expression cassette. *Mol. Gen. Genet.* **217**:202–208.
170. Wright, G. D. 1999. Aminoglycoside-modifying enzymes. *Curr. Opin. Microbiol.* **2**:499–503.
171. Xiong, Y. Q., J. Caillon, M. F. Kergueris, H. Drugeon, D. Baron, G. Potel, and A. S. Bayer. 1997. Adaptive resistance of *Pseudomonas aeruginosa* induced by aminoglycosides and killing kinetics in a rabbit endocarditis model. *Antimicrob. Agents Chemother.* **41**:823–826.
172. Yagisawa, M., S. Kondo, T. Takeuchi, and H. Umezawa. 1975. Aminoglycoside 6'-*N*-acetyltransferase of *Pseudomonas aeruginosa*: structural requirements of substrate. *J. Antibiot. (Tokyo)* **28**:486–489.
173. Yamane, K., Y. Doi, K. Yokoyama, T. Yagi, H. Kurokawa, N. Shibata, K. Shibayama, H. Kato, and Y. Arakawa. 2004. Genetic environments of the *mtA* gene in *Pseudomonas aeruginosa* clinical isolates. *Antimicrob. Agents Chemother.* **48**:2069–2074.
174. Yokoyama, K., Y. Doi, K. Yamane, H. Kurokawa, N. Shibata, K. Shibayama, T. Yagi, H. Kato, and Y. Arakawa. 2003. Acquisition of 16S rRNA methylase gene in *Pseudomonas aeruginosa*. *Lancet* **362**:1888–1893.
175. Yoneyama, H., K. Sato, and T. Nakae. 1991. Aminoglycoside resistance in *Pseudomonas aeruginosa* due to outer membrane stabilization. *Chemotherapy (Basel)* **37**:239–245.
176. Young, M. L., M. Bains, A. Bell, and R. E. Hancock. 1992. Role of *Pseudomonas aeruginosa* outer membrane protein OprH in polymyxin and gentamicin resistance: isolation of an OprH-deficient mutant by gene replacement techniques. *Antimicrob. Agents Chemother.* **36**:2566–2568.
177. Young, S. A., F. C. Tenover, T. D. Gootz, K. P. Gordon, and J. J. Plorde. 1985. Development of two DNA probes for differentiating the structural genes of subclasses I and II of the aminoglycoside-modifying enzyme 3'-aminoglycoside phosphotransferase. *Antimicrob. Agents Chemother.* **27**:739–744.
178. Zeng, L., and S. Jin. 2003. *aph(3')-IIB*, a gene encoding an aminoglycoside-modifying enzyme, is under the positive control of surrogate regulator HpaA. *Antimicrob. Agents Chemother.* **47**:3867–3876.