

Increasing Prevalence and Diversity of Metallo- β -Lactamases in *Pseudomonas* spp., *Acinetobacter* spp., and *Enterobacteriaceae* from Korea

Donggeun Yong,^{1,2,3} Yeoung Seon Choi,^{2,3} Kyoung Ho Roh,^{1,2} Chang Ki Kim,¹ Youn Hee Park,¹ Jong Hwa Yum,^{2,3} Kyungwon Lee,^{1,2,3*} and Yunsop Chong^{1,2}

Department of Laboratory Medicine,¹ Research Institute of Bacterial Resistance,² and BK21 Project for Medical Sciences,³ Yonsei University College of Medicine, Seoul, Korea

Received 26 October 2005/Returned for modification 28 November 2005/Accepted 10 February 2006

Among imipenem-nonsusceptible isolates, acquired metallo- β -lactamase genes were detected in 36 of 581 (6.2%) *Pseudomonas aeruginosa* isolates, 42 of 44 (95.4%) other *Pseudomonas* species, and 136 of 513 (26.5%) *Acinetobacter* species from 2003 to 2004 at a Korean hospital. Overall, *bla*_{VIM-2}-like genes were the most prevalent and were also detected in *Enterobacteriaceae*, including *Klebsiella pneumoniae*.

Carbapenems have been the β -lactam antibiotics used most successfully to evade bacterial resistance (8), but acquired carbapenem resistance due to the production of metallo- β -lactamases (MBLs) has been increasingly reported, particularly for *Pseudomonas aeruginosa* and *Acinetobacter* spp. (13).

In recent years, carbapenem resistance found in *P. aeruginosa* and *Acinetobacter* spp. has gradually increased in Korea, and a significant proportion of these carbapenem-resistant isolates have been shown to produce VIM-2- or IMP-1-type MBL (5). Among the members of the family *Enterobacteriaceae*, imipenem resistance has been found to be virtually nonexistent.

In 2003 and 2004, an increasing trend in the rates of carbapenem resistance in *P. aeruginosa* and *Acinetobacter* spp., along with a sudden increase in imipenem-resistant *Klebsiella pneumoniae* isolates, was observed at a Korean tertiary-care hospital. In this study, we determined the prevalence of MBL-producing isolates and identified the types of MBL in imipenem-nonsusceptible *Pseudomonas* spp., *Acinetobacter* spp., and *Enterobacteriaceae* isolates from that hospital during that period.

The study included randomly selected imipenem-nonsusceptible *Pseudomonas* spp., *Acinetobacter* spp., and *Enterobacteriaceae* isolates from clinical specimens collected in 2003 and 2004. Isolates were identified by conventional testing (4) or with an ATB 32 GN system (bioMérieux, Marcy-l'Étoile, France). A disk diffusion method (2) was used for routine susceptibility testing. MBL-producing isolates were screened with the modified-imipenem-disk Hodge test using MacConkey agar plates and with the double-disk synergy test using imipenem and EDTA-sodium mercaptoacetic acid disks (6). The *bla*_{IMP-1}, *bla*_{VIM-2}, and *bla*_{SIM-1}-like genes were detected by PCR as previously described (1, 7, 11). The *bla*_{VIM-2} gene-carrying integrons from two *K. pneumoniae* isolates were amplified and sequenced as previously described (12). A template prepared from a transconjugant (the recipient was azide-resis-

tant *Escherichia coli* J53) and primers SHV-EXT-F (5'-TTCTTTACTCGCCTTTATCG-3') and SHV-EXT-R (5'-TTTATGCGTTACCTTTGAC-3') were used to detect and sequence the *bla*_{SHV} allele originally carried in an aztreonam-resistant *K. pneumoniae* isolate.

In 2004, a notable increase in imipenem resistance rates was observed for *P. aeruginosa*, other *Pseudomonas* spp., and *Acinetobacter* spp. compared to those in 2003 (Table 1). Of the relatively small number of imipenem-resistant isolates of *Enterobacteriaceae* found from 2003 to 2004, the majority were *K. pneumoniae*, while the minority consisted of *Enterobacter cloacae* and *Serratia marcescens* (Table 1).

Among the imipenem-nonsusceptible isolates tested, 36 of 581 (6%) *P. aeruginosa* isolates and 42 of 44 (95%) other *Pseudomonas* spp. produced an acquired MBL (Table 1). In Korea, VIM-2 has been the only MBL type detected in *Pseudomonas* spp. since 1995 (5; K. Lee, unpublished results). In the present study, most MBL genes detected in *P. aeruginosa* were *bla*_{VIM-2}-like, but we also detected for the first time in Korea the presence of two isolates carrying *bla*_{IMP-1}-like genes. This may suggest dissemination from Japan, where IMP-type enzymes are the most prevalent MBLs in *Pseudomonas* spp. (10). Interestingly, all 42 MBL-producing isolates of other *Pseudomonas* spp. were found to be *Pseudomonas putida* (Table 1). In this study, the number of MBL-producing isolates of *P. aeruginosa* was surpassed by the number of *P. putida* isolates, which suggests that *P. putida* plays a significant role as a reservoir of MBL genes, even though it is a species that is rarely involved in infection (4). The MBL genes detected in all 42 *P. putida* isolates were *bla*_{VIM-2}-like.

Overall, 136 of the 513 (26.5%) imipenem-nonsusceptible *Acinetobacter* sp. isolates tested carried an acquired MBL determinant (Table 1). In 1998, only *bla*_{IMP-1}-like genes were present in *Acinetobacter* spp. (15), but since 2001, the proportion of isolates with *bla*_{VIM-2}-like genes has gradually increased (J. H. Yum, unpublished results). In the present study, 64% of the MBL-producing *Acinetobacter* isolates in 2004 had *bla*_{VIM-2}-like genes, while only 29% of them had *bla*_{IMP-1}-like genes (Table 1). Two isolates of *Acinetobacter baumannii* with a novel MBL, SIM-1, in addition to seven previously

* Corresponding author. Mailing address: Department of Laboratory Medicine, Yonsei University College of Medicine, 134 Shinchon-dong, Seodaemunku, Seoul 120-752, Korea. Phone: 82-2-2228-2446. Fax: 82-2-313-0908. E-mail: leekcp@yumc.yonsei.ac.kr.

TABLE 1. Results of routine imipenem susceptibility testing and detection of MBL-producing gram-negative bacilli in 2003 and 2004 in Korea

Species	Yr	No. (%) of isolates		Tested	No. (%) of nonduplicate imipenem-nonsusceptible isolates				
		Total	Imipenem-resistant		Positive with:				
					Hodge test	Double-disk synergy test	<i>bla</i> _{VIM-2} -like gene ^a	<i>bla</i> _{IMP-1} -like gene ^a	<i>bla</i> _{SIM-1} -like gene ^a
<i>P. aeruginosa</i>	2003	2,913	816 (28)	243	27 (11)	16 (7)	16 (100)	0	0
	2004	3,057	1,009 (33)	338	38 (11)	20 (6)	18 (90)	2 (10)	0
Other <i>Pseudomonas</i> spp.	2003	144	79 (55)	14	14 (100)	14 (100)	14 (100) ^b	0	0
	2004	114	70 (61)	30	28 (93)	28 (93)	28 (100) ^b	0	0
<i>Acinetobacter</i> spp.	2003	1,765	282 (16)	194	65 (34)	51 (26)	27 (53)	21 (41)	3 (6)
	2004	1,529	459 (30)	319	222 (70)	85 (27)	54 (64)	25 (29)	6 (7)
<i>K. pneumoniae</i>	2003	1,819	36 (2)	22	2 (9)	2 (9)	2 (100)	0	0
	2004	2,006	39 (2)	22	2 (9)	0	0	0	0
<i>E. cloacae</i>	2003	821	2 (<1)	1	1 (100)	1 (100)	1 (100)	0	0
	2004	773	1 (<1)	1	1 (100)	0	0	0	0
<i>S. marcescens</i>	2003	536	1 (<1)	1	1 (100)	1 (100)	1 (100)	0	0
	2004	483	4 (<1)	4	1 (25)	0	0	0	0
Total		15,960	2,797 (18)	1189	402 (34)	217 (18)	161 (74)	48 (22)	9 (4)

^a Numbers in parentheses indicate the proportions of isolates among all MBL-producing isolates.

^b All *bla*_{VIM-2}-like-gene-positive isolates were *P. putida*.

reported isolates, were also detected (7). It is important to note that many imipenem-nonsusceptible but MBL-negative isolates of *P. aeruginosa* and *Acinetobacter* spp. were detected in this study. These findings suggest a wide dissemination of isolates with other resistance mechanisms. Carbapenem-hydrolyzing OXA-23 β -lactamase was reported for *Acinetobacter* strains from another Korean hospital (3).

Only *bla*_{VIM-2}-like genes were detected in enterobacterial isolates (Table 1), and to the best of our knowledge, this is the first detection of similar determinants in *K. pneumoniae*. The structures of class 1 integrons carrying *bla*_{VIM-2} from these two isolates (GenBank accession no. DQ153217 and DQ153218) were identical to those of *bla*_{VIM-2}-containing integrons from a *Providencia rettgeri* isolate (GenBank accession no. AY887109), suggesting the dissemination of these integrons among *Enterobacteriaceae* by horizontal transfer.

The MICs of imipenem for VIM-4-producing *K. pneumoniae* and for VIM-2-producing *Citrobacter freundii* were reported to be low, 2 μ g/ml and 1 μ g/ml, respectively (9, 14). However, the imipenem MICs for the two VIM-2-producing *K. pneumoniae* isolates in our study, determined by an agar dilution method (2), were much higher, namely, 32 μ g/ml and >128 μ g/ml, suggesting the contribution of additional resistance mechanisms. Although MBLs do not hydrolyze aztreonam, the MIC of aztreonam for one of the *bla*_{VIM-2}-positive *K. pneumoniae* isolates (YMC 03/10/U1702) was significantly higher (128 μ g/ml) than that for the other isolate (0.5 μ g/ml). The double-disk synergy test was positive for the aztreonam-resistant isolate, and the *bla*_{SHV-12} gene was detected in a transconjugant, which explains the high-level resistance to the monobactam.

In this study, 90% of the MBL-producing isolates were from patients in a general ward or an intensive care unit, while 10% were from patients in an emergency room or another outpatient department (data not shown). The majority (82%) of MBL-producing isolates were from sputum and urine specimens, as reported in a previous study (5).

In conclusion, imipenem-nonsusceptible *P. aeruginosa* isolates carrying *bla*_{VIM-2}-like genes remain highly prevalent, while the

incidence of *Acinetobacter* spp. positive for *bla*_{VIM-2}-like genes has increased significantly. The spread of *bla*_{IMP-1}-like genes to *P. aeruginosa* and *bla*_{VIM-2}-like genes to *Acinetobacter* spp. or even to *K. pneumoniae* is a cause for concern and warrants continuous surveillance to control the further spread of these resistance mechanisms.

This study was supported in part by grant 2M16330 from The Ministry of Commerce, Industry and Energy, Republic of Korea.

REFERENCES

- Chong, Y., K. Lee, R. Okamoto, and M. Inoue. 1997. Characteristics of extended-spectrum β -lactam hydrolyzing activity of *Klebsiella pneumoniae* and *Escherichia coli* strains isolated from clinical specimens. Korean J. Infect. Dis. 29:477-485.
- Clinical and Laboratory Standards Institute. 2005. Performance standards for antimicrobial susceptibility testing; 15th informational supplement. CLSI/NCCLS M100-S15. Clinical and Laboratory Standards Institute, Wayne, Pa.
- Jeon, B.-C., S. H. Jeong, I. K. Bae, S. B. Kwon, K. Lee, D. Young, J. H. Lee, J. S. Song, and S. H. Lee. 2005. Investigation of a nosocomial outbreak of imipenem-resistant *Acinetobacter baumannii* producing the OXA-23 β -lactamase in Korea. J. Clin. Microbiol. 43:2241-2245.
- Kiska, D. L., and P. H. Gilligan. 2003. *Pseudomonas*, p. 719-728. In P. R. Murray, E. J. Baron, J. H. Tenover, M. A. Tenover, and R. H. Tenover (eds.), Manual of clinical microbiology, 8th ed. American Society for Microbiology, Washington, D.C.
- Lee, K., W. G. Lee, Y. Uh, G. Y. Ha, J. Cho, Y. Chong, and the Korean Nationwide Surveillance of Antimicrobial Resistance Group. 2003. VIM- and IMP-type metallo- β -lactamase-producing *Pseudomonas* spp. and *Acinetobacter* spp. in Korean hospitals. Emerg. Infect. Dis. 9:868-871.
- Lee, K., Y. S. Lim, D. Yong, J. H. Yum, and Y. Chong. 2003. Evaluation of the Hodge test and the imipenem-EDTA double-disk synergy test for differentiation metallo- β -lactamase-producing isolates of *Pseudomonas* spp. and *Acinetobacter* spp. J. Clin. Microbiol. 41:4623-4629.
- Lee, K., J. H. Yum, D. Yong, H. M. Lee, H. D. Kim, J.-D. Docquier, G. M. Rossolini, and Y. Chong. 2005. Novel acquired metallo- β -lactamase gene, *bla*_{SIM-1}, in a class 1 integron from *Acinetobacter baumannii* clinical isolates from Korea. Antimicrob. Agents Chemother. 49:4485-4491.
- Livermore, D. M. 2002. The impact of carbapenemases on antimicrobial development and therapy. Curr. Opin. Investig. Drugs 3:218-224.
- Luzzaro, F., J.-D. Docquier, C. Colinon, A. Endimiani, G. Lombardi, G. Amicosante, G. M. Rossolini, and A. Toniolo. 2004. Emergence in *Klebsiella pneumoniae* and *Enterobacter cloacae* clinical isolates of the VIM-4 metallo- β -lactamase encoded by a conjugative plasmid. Antimicrob. Agents Chemother. 48:648-650.
- Nishio, H., M. Komatsu, N. Shibata, K. Shimakawa, N. Sueyoshi, T. Ura, K. Satoh, M. Toyokawa, T. Nakamura, Y. Wada, T. Orita, T. Kofuku, K. Yamasaki, M. Sakamoto, S. Kinoshita, M. Aihara, and Y. Arakawa. 2004.

- Metallo- β -lactamase-producing gram-negative bacilli: laboratory-based surveillance in cooperation with 13 clinical laboratories in the Kinki region of Japan. *J. Clin. Microbiol.* **42**:5256–5263.
11. **Poirel, L., T. Naas, D. Nicolas, L. Collet, S. Bellais, J.-D. Cavallo, and P. Nordmann.** 2000. Characterization of VIM-2, a carbapenem-hydrolyzing metallo- β -lactamase and its plasmid- and integron-borne gene from a *Pseudomonas aeruginosa* clinical isolate in France. *Antimicrob. Agents Chemother.* **44**:891–897.
 12. **Riccio, M. L., N. Franceschini, L. Boschi, B. Caravelli, G. Cornaglia, R. Fontana, G. Amicosante, and G. M. Rossolini.** 2000. Characterization of the metallo- β -lactamase determinant of *Acinetobacter baumannii* AC-54/97 reveals the existence of *bla*_{IMP} allelic variants carried by gene cassettes of different phylogeny. *Antimicrob. Agents Chemother.* **44**:1229–1235.
 13. **Walsh, T. R., M. A. Toleman, L. Poirel, and P. Nordmann.** 2005. Metallo- β -lactamases: the quiet before the storm? *Clin. Microbiol. Rev.* **18**:306–325.
 14. **Yan, J. J., W. C. Ko, C. L. Chuang, and J. J. Wu.** 2002. Metallo- β -lactamase-producing *Enterobacteriaceae* isolates in a university hospital in Taiwan: prevalence of IMP-8 in *Enterobacter cloacae* and first identification of VIM-2 in *Citrobacter freundii*. *J. Antimicrob. Chemother.* **50**:503–511.
 15. **Yum, J. H., D. Yong, K. Lee, H. S. Kim, and Y. Chong.** 2002. A new integron carrying VIM-2 metallo- β -lactamase gene cassette in a *Serratia marcescens* isolate. *Diagn. Microbiol. Infect. Dis.* **42**:217–219.