

Update to the Multiplex PCR Strategy for Assignment of *mec* Element Types in *Staphylococcus aureus*[∇]

Catarina Milheiro,¹ Duarte C. Oliveira,^{1*} and Hermínia de Lencastre^{1,2}

Laboratory of Molecular Genetics, Instituto de Tecnologia Química e Biológica, Universidade Nova de Lisboa, Oeiras, Portugal,¹
 and Laboratory of Microbiology, The Rockefeller University, New York, New York 10021²

Received 23 February 2007/Returned for modification 7 April 2007/Accepted 10 June 2007

Staphylococcal cassette chromosome *mec* (SCC*mec*) typing is important for the identification and definition of methicillin-resistant *Staphylococcus aureus* clones, and for routine purposes, multiplex PCR assays are the most adequate for SCC*mec* typing. Here, we describe an update to the multiplex PCR strategy for SCC*mec* typing that we described in 2002 so that SCC*mec* types IV and V may be properly identified.

Methicillin-resistant *Staphylococcus aureus* (MRSA) strains are characterized by the presence of a large heterologous mobile genetic element called the staphylococcal cassette chromosome *mec* (SCC*mec*), which includes the *mecA* gene, the central element of methicillin resistance (4). Besides the *mec* gene complex (which comprises the *mecA* gene and its regulators, *mecI* and *mecR1*), SCC*mec* contains the *ccr* gene complex, which encodes recombinases responsible for the mobility of SCC*mec* (7). Several SCC*mec* types have been defined by use of the combination of the class of the *mec* gene complex and the *ccr* allotype (1, 3–5, 9, 16). The remaining parts of SCC*mec* are called J regions (regions J1, J2, and J3), which constitute nonessential components of the cassette, although in some cases these regions carry additional antibiotic resistance determinants. J1 is the region between the chromosomal left junction and the *ccr* complex, J2 is the region between the *ccr* complex and the *mec* complex, and J3 is the region between the *mec* complex and the chromosomal right junction. Therefore, the structural organization of SCC*mec* may be summarized as J1-*ccr*-J2-*mec*-J3. Variations in the J regions within the same *mec*-*ccr* combination are used to define SCC*mec* subtypes.

In 2002, we described a multiplex PCR strategy for the rapid assignment of SCC*mec* types to MRSA strains. That strategy was able to properly identify SCC*mec* types I to III and some epidemiologically relevant variants (e.g., subtypes IA and IIIA) by probing eight loci scattered through the *mec* elements and generating specific amplification fragments of three to five bands (13). SCC*mec* type IV, which at that time was not yet recognized as an important structural type mainly due to its spread among community-acquired MRSA (CA-MRSA) strains, was not properly identified since it was positive only for the internal positive control and the *dcs* locus, which is also present in SCC*mec* types I and II. Here, we report an update to the previously described “SCC*mec* multiplex PCR strategy” in order to better characterize SCC*mec* type IV and also to include the detection of the recently described SCC*mec* type V.

Table 1 lists the characteristics of the primers used for the

updated version of the SCC*mec* multiplex PCR. In order to minimize the complexity of the multiplex PCR, the detection of linearized plasmids pUB110 and pT181 was abandoned. These loci are not critical for SCC*mec* type assignment, and its utility was to discriminate subtypes IA (positive for pUB110) and IIIA (negative for pT181). Eight new primers were added for the detection of *ccrB* allotype 2 (specific for SCC*mec* types II and IV), *ccrC* (specific for SCC*mec* type V), the SCC*mec* type III J1 region, and the SCC*mec* type V J1 region.

The conditions for the multiplex PCR assay were first optimized by using the following prototype strains: COL, SCC*mec* type I (16); N315, SCC*mec* type II (4); ANS46, SCC*mec* type III (16); MW2, SCC*mec* type IVa (1); 8/6-3P, SCC*mec* type IVb (9); Q2314, SCC*mec* type IVc (6); JCS4469, SCC*mec* type IVd; AR43/3330.1, SCC*mec* type IVE (17); M03-68, SCC*mec* type IVg (8); HAR22, SCC*mec* type IVh (11); WIS, SCC*mec* type V (5); and HDE288, SCC*mec* type VI (15). For validation purposes, a diverse collection of 60 MRSA isolates previously characterized in terms of their genetic backgrounds and SCC*mec* types was tested by use of the updated SCC*mec* multiplex PCR assay (Table 2). All assays were performed in a T1 thermocycler (Biometra, Germany). The optimal cycling conditions were the following: 94°C for 4 min; 30 cycles of 94°C for 30 s, 53°C for 30 s, and 72°C for 1 min; and a final extension at 72°C for 4 min. Each PCR mixture, in a final volume of 50 μ l, contained 5 ng of chromosomal template; 1 \times PCR buffer with 1.5 mM MgCl₂ (Applied Biosystems); 40 μ M (each) deoxynucleoside triphosphate (MBI Fermentas, Hanover, MD); 0.2 μ M primers kdp F1 and kdp R1; 0.4 μ M primers CIF2 F2, CIF2 R2, RIF5 F10, RIF5 R13, SCC*mec* III J1F, SCC*mec* III J1R, SCC*mec* V J1 F, and SCC*mec* V J1 R; 0.8 μ M primers *mecI* P2, *mecI* P3, *dcs* F2, *dcs* R1, *mecA* P4, *mecA* P7, *ccrB2* F2, *ccrB2* R2, *ccrC* F2, and *ccrC* R2; and 1.25 U of Amplitaq DNA polymerase (Applied Biosystems). The PCR products (10 μ l) were resolved in a 3% Seakem LE (Cambrex, Rockland, ME) agarose gel in 0.5% Tris-borate-EDTA buffer (Bio-Rad, Hercules, CA) at 4 V/cm for 2.5 h and were visualized with ethidium bromide.

Figure 1 illustrates the amplification patterns obtained for the prototype strains. SCC*mec* types I to III generate specific amplification patterns of three to five bands. SCC*mec* type IV generates an amplification pattern of three bands (*mecA*,

* Corresponding author. Mailing address: Instituto de Tecnologia Química e Biológica, Universidade Nova de Lisboa, Oeiras, Portugal. Phone: (351) 21 446 9862. Fax: (351) 21 442 8766. E-mail: dco@itqb.unl.pt.

[∇] Published ahead of print on 18 June 2007.

TABLE 1. Primers used in the updated version of SCCmec multiplex PCR

Primer name	Primer sequence (5'→3')	Primer specificity (SCCmec type, region)	Amplicon size (bp)	Reference
CIF2 F2 CIF2 R2	TTCGAGTTGCTGATGAAGAAGG ATTTACCACAAGGACTACCAGC	I, J1 region	495	13 13
ccrC F2 ccrC R2	GTACTIONGTTACAATGTTTGG ATAATGGCTTCATGCTTACC	V, <i>ccr</i> complex	449	This study This study
RIF5 F10 RIF5 R13	TTCTTAAGTACACGCTGAATCG ATGGAGATGAATTACAAGGG	III, J3 region	414	13 13
SCCmec V J1 F SCCmec V J1 R	TTCTCCATTCTGTTCATCC AGAGACTACTGACTTAAGTGG	V, J1 region	377	This study This study
dcs F2 dcs R1	CATCCTATGATAGCTTGGTC CTAAATCATAGCCATGACCG	I, II, IV, and VI, J3 region	342	13 13
ccrB2 F2 ccrB2 R2	AGTTTCTCAGAATTCGAACG CCGATATAGAAWGGGTTAGC	II and IV, <i>ccr</i> complex	311	This study This study
kdp F1 kdp R1	AATCATCTGCCATTGGTGATGC CGAATGAAGTGAAGAAAGTGG	II, J1 region	284	13 13
SCCmec III J1 F SCCmec III J1 R	CATTTGTGAAACACAGTACG GTTATTGAGACTCCTAAAGC	III, J1 region	243	This study This study
mecI P2 mecI P3	ATCAAGACTTGCATTCAGGC GCGGTTTCAATTCACCTTGTC	II and III, <i>mec</i> complex	209	13 13
mecA P4 mecA P7	TCCAGATTACAACCTCACCAGG CCACTTCATATCTTGTAACG	Internal positive control	162	13 13

ccrB2, and *dcs*) for subtypes a to d, g, and h; and subtypes IVE and IVF generate only two bands, since they are negative for the *dcs* locus. SCCmec type V also originates a pattern of three bands (*mecA*, *ccrC*, and J1). SCCmec type VI, which appears to be an “exotic” local variant (14), generates only two bands (*mecA* and *dcs*), and its assignment must be confirmed by *ccrAB* allotyping (*ccrAB4*). Note that SCCmec types IV and VI previously could not be discriminated since both were positive only for *mecA* and *dcs*.

As illustrated in Table 2, the updated SCCmec multiplex PCR assay performed well in assigning SCCmec types to the 60 diverse MRSA isolates. As expected, subtypes IA and IIIA, defined by the presence of pUB110 and the absence of pT181, respectively, could not be detected, whereas the sporadically occurring subtype IIIB, defined as being negative for the J3 region specific locus, was still detected. Strains PL72 and POL3 were incorrectly assigned to SCCmec type VI. These sporadically occurring strains were first tentatively classified as SCCmec type IV (13) and after *ccrB* sequencing (14) were classified as new type I variants with a partial deletion within the J1 region, which includes the J1-specific locus of SCCmec type I included in the multiplex PCR assay. This unique example of an incorrect assignment reinforces the importance of confirming multiplex PCR results by *ccr* gene complex typing, either by conventional PCR allotyping (12) or by DNA sequencing (15), at least for a subset of strains belonging to the same clonal type or whenever a new clone is detected.

In short, the updated version of the previously described SCCmec multiplex PCR strategy enables the rapid presumptive assignment of all known SCCmec types to MRSA strains.

Whereas the previous version enabled the prompt identification of only SCCmec types I to III, since SCCmec type IV was poorly identified and SCCmec type V was not detected, the means for the proper identification of SCCmec types IV and V was added to this updated version. The prompt detection of SCCmec types IV and V is particularly relevant for the characterization of the recent threat of CA-MRSA, mostly characterized by these two structural variants of the *mec* element. In practical terms, changes to the previous protocol were kept to a minimum, and so the implementation of this SCCmec typing strategy will be straightforward, particularly for those laboratories currently using the previously described SCCmec multiplex PCR strategy.

The importance of SCCmec typing is well illustrated by the fact that the proposal by Enright and colleagues (2) that MRSA clones be named according to their multilocus sequence types and SCCmec types (e.g., clone ST5-MRSA-II), which was agreed to by a subcommittee of the International Union of Microbiology Societies in Tokyo, Japan, in 2002, is currently consensual in the specialized literature. In this context, rapid and easy assays for the detection of SCCmec types, such as the multiplex PCR typing strategy described in this study, are critical tools for the proper characterization of MRSA clones. The SCCmec element, which carries the determinant for “broad-spectrum” beta-lactam resistance in staphylococci, is a critical epidemiological marker for MRSA clones. However, besides being an important tool for surveillance studies, SCCmec typing of large international collections of isolates has contributed dramatically to the elucidation of the

TABLE 2. Prototype strains and representative collection used for validation of updated SCCmec multiplex PCR assay

Strain ^a	Origin	Isolation date	MLST (ST) ^b	SCCmec type	<i>ccrAB</i> type	Multiplex PCR assignment	Reference ^c
COL	United Kingdom	1965	250	I	1	I	16
UK13136	United Kingdom	1960	250	I	1	I	16
BK793	Egypt	1961	250	I	1	I	16
E2125	Denmark	1964	247	I	1	I	16
PER34	Spain	1989	250	IA	1	I	16
HPV107	Portugal	1992	247	IA	1	I	16
BK1953	United States	1995	247	IA	1	I	16
PER184	Spain	1991	247	IA	1	I	16
PER88	Spain	1992	247	IA	1	I	16
HAR40	Belgium	1995	228	I	1	I	15
HAR21	Finland	2002	5	I	1	I	15
PL72	Poland	1991	5	I-VAR ^d	1	VI	15
POL3	Poland	1992	5	I-VAR	1	VI	15
N315	Japan	1982	5	II	2	II	4
USA100	United States	1995–2003	5	II	2	II	10
BK2464	United States	1996	5	II	2	II	16
JP1	Japan	1987	5	II	2	II	16
JP205	Japan	1998	5	II	2	II	15
USA200	United States	1995–2003	36	II	2	II	10
HAR24	Finland	2002	36	II	2	II	15
USA600	United States	1995–2003	45	II	2	II	10
ANS46	Australia	1982	239	III	3	III	16
R35	United States	1987	239	III	3	III	16
HU25	Brazil	1993	239	III	3	III	16
HUSA304	Hungary	1993	239	III	3	III	16
HU106	Hungary	1996	239	III	3	III	16
LHH1	United States	1994	239	III	3	III	16
BK2421	United States	1996	239	III	3	III	16
HSJ216	Portugal	1997	239	IIIA	3	III	16
HDG2	Portugal	1993	239	IIIB	3	IIIB	16
MW2	United States	1998	1	IVa	2	IV	1
CA05	Chicago	1999	256	IVa	2	IV	9
8/6-3P	Chicago	1996	770	IVb	2	IV	9
Q2314	Dallas, TX	1996	5	IVc	2	IV	6
JCSC4469	Japan	1982	5	IVd	2	IV	AB097677
AR43/3330.1	Ireland	1988–2002	8	IVE	2	IV	17
M03-68	Korea	2003	5	IVg	2	IV	8
HAR22	Finland	2002	22	IVh	2	IV	11
USA400	United States	1995–2003	1	IV	2	IV	10
USA800	United States	1995–2003	5	IV	2	IV	10
BM18	United States	1989	5	IV	2	IV	16
COB3	Colombia	1996	5	IV	2	IV	16
FFP311	Portugal	1996	5	IV	2	IV	14
VNG17	Portugal	1992–1993	5	IV	2	IV	14
RJP17	Portugal	1992–1993	5	IV	2	IV	14
HSA49	Portugal	1993	5	IV	2	IV	15
HSA74	Portugal	1993	5	IV	2	IV	14
ARG9	Argentina	1996	5	IV	2	IV	14
USA300	United States	1995–2003	8	IV	2	IV	10
USA500	United States	1995–2003	8	IV	2	IV	10
BK2529	United States	1996	8	IV	2	IV	16
BargII17	United States	1996	8	IV	2	IV	16
GRE120	Greece	1993	8	IV	2	IV	15
HAR36 ^e	Greece	2002	254	IV-VAR	2	IV-VAR	15
HGSA146	Portugal	2003	22	IVh	2	IV	11
HGSA163	Portugal	2003	22	IVh	2	IV	11
DEN2946	Denmark	2001	30	IV	2	IV	15
DEN2294	Denmark	2001	30	IV	2	IV	15
HAR38	Belgium	1995	45	IV	2	IV	15
USA700	United States	1995–2003	72	IV	2	IV	10
GRE14	Greece	1998	80	IV	2	IV	15
DEN2949	Denmark	2001	80	IV	2	IV	15
DEN114	Denmark	2001	80	IV	2	IV	15
DEN1451	Denmark	2001	153	IV	2	IV	15
WIS	Australia			V	<i>ccrC</i>	V	5
HT0087	Australia	2005	377	V	<i>ccrC</i>	V	18
HT0184	Greece	2005	377	V	<i>ccrC</i>	V	18
HT0350	The Netherlands	2003	377	V	<i>ccrC</i>	V	18
HT0388	Switzerland	2003	377	V	<i>ccrC</i>	V	18
HT0826	France	2003	377	V	<i>ccrC</i>	V	18
HDE288	Portugal	1996	5	VI	4	VI	14
HUC136	Portugal	1995	5	VI	4	VI	14
IPO92	Portugal	2001	5	VI	4	VI	14

^a Strains in boldface are the prototype strains for SCCmec types and subtypes.

^b MLST, multilocus sequence type; ST, sequence type.

^c Reference for SCCmec type assignment and *ccr* type.

^d VAR, variant.

^e Strain HAR36 is characterized by SCCmec type IV and is also positive for the J3 locus specific for SCCmec type III.




FIG. 1. Amplification patterns for the prototype strains obtained with the updated SCCmec multiplex PCR strategy. Lane 1, SCCmec type I (strain COL); lane 2, type II (N315); lane 3, type III (ANS46), lane 4, type IVa (MW2); lane 5, type IVb (8/6-3P); lane 6, type IVc (Q2314); lane 7, type IVd (JCSC4469); lane 8, type IVE (AR43/3330.1); lane 9, type IVg (M03-68); lane 10, type IVh (HAR22); lane 11, type V (WIS); lane 12, type VI (HDE288); lanes M, DNA molecular size marker (1-kb DNA Ladder Plus; Invitrogen Life Technologies, Carlsbad, CA).

origin(s) and evolutionary history of contemporary MRSA clones.

We thank T. Ito, D. C. Coleman, R. Daum, K. T. Park, W. B. Grubb, J. Etienne, and A. Tomasz for having kindly given some of the prototype and reference strains used in this study.

Partial support for this study was provided by projects POCI/BIA-MIC/58416/2004 from the Fundação para a Ciência e Tecnologia (FCT), Lisbon, Portugal, and FCG-55068 from the Fundação Calouste Gulbenkian, Lisbon, Portugal, both awarded to H. de Lencastre. C. Milheirico and D. C. Oliveira were supported by grants SFRH/BD/23010/2005 and SFRH/BPD/9374/2002, respectively, from FCT.

REFERENCES

- Baba, T., F. Takeuchi, M. Kuroda, H. Yuzawa, K. Aoki, A. Oguchi, Y. Nagai, N. Iwama, K. Asano, T. Naimi, H. Kuroda, L. Cui, K. Yamamoto, and K. Hiramatsu. 2002. Genome and virulence determinants of high virulence community-acquired MRSA. *Lancet* **359**:1819–1827.
- Enright, M. C., D. A. Robinson, G. Randle, E. J. Feil, H. Grundmann, and B. G. Spratt. 2002. The evolutionary history of methicillin-resistant *Staphylococcus aureus* (MRSA). *Proc. Natl. Acad. Sci. USA* **99**:7687–7692.
- Ito, T., Y. Katayama, K. Asada, N. Mori, K. Tsutsumimoto, C. Tiensasitorn, and K. Hiramatsu. 2001. Structural comparison of three types of staphylococcal cassette chromosome *mec* integrated in the chromosome in methicillin-resistant *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **45**:1323–1336.
- Ito, T., Y. Katayama, and K. Hiramatsu. 1999. Cloning and nucleotide sequence determination of the entire *mec* DNA of pre-methicillin-resistant *Staphylococcus aureus* N315. *Antimicrob. Agents Chemother.* **43**:1449–1458.
- Ito, T., X. X. Ma, F. Takeuchi, K. Okuma, H. Yuzawa, and K. Hiramatsu. 2004. Novel type V staphylococcal cassette chromosome *mec* driven by a novel cassette chromosome recombinase, *ccrC*. *Antimicrob. Agents Chemother.* **48**:2637–2651.
- Ito, T., K. Okuma, X. X. Ma, H. Yuzawa, and K. Hiramatsu. 2003. Insights on antibiotic resistance of *Staphylococcus aureus* from its whole genome: genomic island SCC. *Drug Resist. Update* **6**:41–52.
- Katayama, Y., T. Ito, and K. Hiramatsu. 2000. A new class of genetic element, staphylococcal cassette chromosome *mec*, encodes methicillin resistance in *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **44**:1549–1555.
- Kwon, N. H., K. T. Park, J. S. Moon, W. K. Jung, S. H. Kim, J. M. Kim, S. K. Hong, H. C. Koo, Y. S. Joo, and Y. H. Park. 2005. Staphylococcal cassette chromosome *mec* (SCCmec) characterization and molecular analysis for methicillin-resistant *Staphylococcus aureus* and novel SCCmec subtype IVg isolated from bovine milk in Korea. *J. Antimicrob. Chemother.* **56**:624–632.
- Ma, X. X., T. Ito, C. Tiensasitorn, M. Jamklang, P. Chongtrakool, S. Boyle-Vavra, R. S. Daum, and K. Hiramatsu. 2002. Novel type of staphylococcal cassette chromosome *mec* identified in community-acquired methicillin-resistant *Staphylococcus aureus* strains. *Antimicrob. Agents Chemother.* **46**:1147–1152.
- McDougal, L. K., C. D. Steward, G. E. Killgore, J. M. Chaitram, S. K. McAllister, and F. C. Tenover. 2003. Pulsed-field gel electrophoresis typing of oxacillin-resistant *Staphylococcus aureus* isolates from the United States: establishing a national database. *J. Clin. Microbiol.* **41**:5113–5120.
- Milheirico, C., D. C. Oliveira, and H. de Lencastre. 2007. Multiplex PCR strategy for subtyping the staphylococcal cassette chromosome *mec* type IV in methicillin-resistant *Staphylococcus aureus*: ‘SCCmec IV multiplex’. *J. Antimicrob. Chemother.* **60**:42–48.
- Okuma, K., K. Iwakawa, J. D. Turnidge, W. B. Grubb, J. M. Bell, F. G. O’Brien, G. W. Coombs, J. W. Pearman, F. C. Tenover, M. Kapi, C. Tiensasitorn, T. Ito, and K. Hiramatsu. 2002. Dissemination of new methicillin-resistant *Staphylococcus aureus* clones in the community. *J. Clin. Microbiol.* **40**:4289–4294.
- Oliveira, D. C., and H. de Lencastre. 2002. Multiplex PCR strategy for rapid identification of structural types and variants of the *mec* element in methicillin-resistant *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **46**:2155–2161.
- Oliveira, D. C., C. Milheirico, and H. De Lencastre. 2006. Redefining a structural variant of staphylococcal cassette chromosome *mec*, SCCmec type VI. *Antimicrob. Agents Chemother.* **50**:3457–3459.
- Oliveira, D. C., C. Milheirico, S. Vinga, and H. de Lencastre. 2006. Assessment of allelic variation in the *ccrAB* locus in methicillin-resistant *Staphylococcus aureus* clones. *J. Antimicrob. Chemother.* **58**:23–30.
- Oliveira, D. C., A. Tomasz, and H. de Lencastre. 2001. The evolution of pandemic clones of methicillin-resistant *Staphylococcus aureus*: identification of two ancestral genetic backgrounds and the associated *mec* elements. *Microb. Drug Resist.* **7**:349–361.
- Shore, A., A. S. Rossney, C. T. Keane, M. C. Enright, and D. C. Coleman. 2005. Seven novel variants of the staphylococcal chromosomal cassette *mec* in methicillin-resistant *Staphylococcus aureus* isolates from Ireland. *Antimicrob. Agents Chemother.* **49**:2070–2083.
- Tristan, A., M. Bes, H. Meugnier, G. Lina, B. Bozdogan, P. Courvalin, M. Reverdy, M. C. Enright, F. Vandenesch, and J. Etienne. 2007. Global distribution of Panton-Valentine leukocidin-positive methicillin-resistant *Staphylococcus aureus*, 2006. *Emerg. Infect. Dis.* **13**:594–600.



ERRATUM

Update to the Multiplex PCR Strategy for Assignment of *mec* Element Types in *Staphylococcus aureus*

Catarina Milheiro, Duarte C. Oliveira, and Hermínia de Lencastre

Laboratory of Molecular Genetics, Instituto de Tecnologia Química e Biológica, Universidade Nova de Lisboa, Oeiras, Portugal, and Laboratory of Microbiology, The Rockefeller University, New York, New York 10021

Volume 51, no. 9, p. 3374–3377, 2007. Page 3375, Table 1: The sequence for primer RIF5 R13 should read “GTC ACA GTA ATT CCA TCA ATG C.”