

ISEcp1-Mediated Transposition of *qnrB*-Like Gene in *Escherichia coli*[∇]

Vincent Cattoir,¹ Patrice Nordmann,^{1*} Jesus Silva-Sanchez,² Paula Espinal,³ and Laurent Poirel¹

Service de Bactériologie-Virologie, INSERM U914 “Emerging Resistance to Antibiotics,” Hôpital de Bicêtre, Assistance Publique/Hôpitaux de Paris, Faculté de Médecine et Université Paris-Sud, K-Bicêtre, France¹;
Departamento de Resistencia Bacteriana, Instituto Nacional de Salud Pública, Cuernavaca, Morelos, Mexico²;
and Facultad de Ciencias de la Salud, Universidad del Sinú, Montería, Colombia³

Received 12 March 2008/Returned for modification 27 April 2008/Accepted 24 May 2008

A novel QnrB-like plasmid-mediated resistance determinant, QnrB19, was identified from an *Escherichia coli* clinical isolate from Colombia. Its gene was associated with an ISEcp1-like insertion element that did not act as a promoter for its expression. Using an in vitro model of transposition, we showed that the ISEcp1-like element was able to mobilize the *qnrB19* gene.

Resistance to quinolones in *Enterobacteriaceae* most commonly arises stepwise as a result of chromosomal mutations responsible for the modification of target enzymes (DNA gyrase and topoisomerase IV) or decreased intracellular drug accumulation by the upregulation of efflux pumps and/or modified outer-membrane porins (27). Since the discovery of the first plasmid-mediated quinolone resistance (PMQR) determinant, QnrA, in 1998 (15), four other PMQR determinants have been identified to date: the QnrB (12) and QnrS (10) proteins, the aminoglycoside acetyltransferase AAC(6′)-Ib-cr (25), and the efflux pump QepA (19, 30). Resistance due to Qnr determinants is increasingly reported worldwide in enterobacterial isolates (18, 26) and has been recently identified outside *Enterobacteriaceae* in environmental *Aeromonas* isolates from France (3). The three types of Qnr determinants, QnrA, QnrB, and QnrS, belong to the pentapeptide repeat family of proteins (18, 26). By protecting DNA gyrase and topoisomerase IV from the inhibitory activity of quinolones, Qnr proteins confer resistance to quinolones (e.g., nalidixic acid) and decreased susceptibility to fluoroquinolones, therefore facilitating the recovery of chromosome-encoded target mutants with a higher level of resistance to fluoroquinolones (18, 26). Whereas *qnrA*-like genes have only been identified as part of the complex *sulI*-type integrons in association with the *orf513* transposase gene (18, 26), which is part of a region redefined as *ISCR1* (28), *qnrB*-like genes have been found to be associated with either the *orf1005* gene encoding a putative transposase (12) or the *ISCR1* element (8, 24).

The aim of this study was (i) to investigate the genetic environment of a *qnrB*-like gene from an *Escherichia coli* clinical isolate, (ii) to evaluate experimentally the mobility of that putative transposon in *E. coli*, and (iii) to determine the promoter sequences of the *qnrB*-like gene responsible for the expression of that gene.

E. coli R4525 expressing an extended-spectrum β-lactamase (ESBL) phenotype had been isolated in 2002 from a wound culture of a patient hospitalized at the Hospital San Jeronimo

in Montería, Colombia. In the course of studying the genetic support of the ESBL determinant in *E. coli* R4525, conjugation experiments followed by selection with sodium azide (100 μg/ml) and amoxicillin (50 μg/ml) as previously described (3) gave *E. coli* J53 transconjugants displaying an ESBL phenotype and decreased susceptibility to fluoroquinolones. This result prompted us to search for the presence of PMQR determinants. The screening of *qnr* genes using a multiplex strategy (5) identified a *qnrB*-like gene in both *E. coli* R4525 and its transconjugant. By contrast, PCR screening performed as described previously (16) did not detect any AAC(6′)-Ib-cr- or QepA-encoding gene. The *qnrB*-like gene was sequenced and found to encode a novel determinant, which was termed QnrB19 in accordance with the recent *qnr* gene nomenclature (11). It differed by a single amino acid substitution at position 212 from QnrB5 (GenBank accession no. DQ303919) previously identified in a non-Typhi *Salmonella* isolate from the United States (9). The MICs were determined on Mueller-Hinton solid agar plates and the results interpreted according to the Clinical and Laboratory Standards Institute guidelines (6). *E. coli* R4525 was resistant to aminoglycosides (except amikacin), chloramphenicol, tetracycline, sulfonamides, and trimethoprim (data not shown). It was resistant to nalidixic acid (MIC > 32 μg/ml) and fluoroquinolones (MICs > 32 μg/ml) (Table 1). Sequence analysis of the quinolone-resistance determining regions (QRDRs) of the *gyrA* and *parC* genes by using primers previously described (2) showed that *E. coli* R4525 possessed two amino acid substitutions both in GyrA (Ser83Leu and Asp87Tyr) and ParC (Ser80Ile and Glu84Gly), compared to wild-type QRDRs of *E. coli* (Table 1), known to confer resistance to quinolones and fluoroquinolones (27). Plasmid analysis of the *E. coli* J53 transconjugant using the Kieser technique (13) identified a single 40-kb plasmid (pR4525) shown by PCR to carry the *bla*_{CTX-M-12}, *bla*_{SHV-12}, and *qnrB19* genes (data not shown). As previously reported for QnrB-like proteins, QnrB19 expressed in the *E. coli* transconjugant conferred increased MICs of quinolones and fluoroquinolones (Table 1) (9, 12, 24).

Since preliminary experiments failed to identify genetic structures that had been associated with *qnrB*-like genes (8, 12, 24), cloning experiments were performed with EcoRI-restricted whole-cell DNA of an *E. coli* R4525 isolate, followed

* Corresponding author. Mailing address: Service de Bactériologie-Virologie-Hygiène, Hôpital de Bicêtre, 78 rue du Général Leclerc, 94275 Le Kremlin-Bicêtre cedex, France. Phone: 33-1-45-21-36-32. Fax: 33-1-45-21-63-40. E-mail: nordmann.patrice@bct.aphp.fr.

[∇] Published ahead of print on 2 June 2008.

TABLE 1. MICs of quinolones for the *E. coli* clinical isolate R4525, its transconjugant, and reference strain *E. coli* J53 Az^r

Strain ^a	Mutations in QRDRs:		MICs (μg/ml) of ^b :					
	<i>gyrA</i>	<i>parC</i>	NAL	NOR	OFL	CIP	ENR	SPX
<i>E. coli</i> R4525	S83L + D87Y	S80I + E84G	>256	>256	>32	>32	>32	>32
Tc J53/R4525	— ^c	—	32	1	1	0.25	0.5	1
<i>E. coli</i> J53 Az ^r	—	—	4	0.12	0.06	0.01	0.01	0.01

^a Az^r, azide resistance; Tc, transconjugant.

^b NAL, nalidixic acid; NOR, norfloxacin; OFL, ofloxacin; CIP, ciprofloxacin; ENR, enrofloxacin; SPX, sparfloxacin.

^c —, no mutation.

by the ligation of DNA fragments into the EcoRI site of cloning vector pBK-CMV (Stratagene, La Jolla, CA), as previously described (3). Analysis of a 3.2-kb DNA fragment carrying the *qnrB19* gene identified its location downstream of the extremity of an *ISEcp1*-like element (Fig. 1). The transposase of the *ISEcp1*-like element (termed *ISEcp1C*) differed by one and two amino acids from those of *ISEcp1* (GenBank accession no. AJ242809) and *ISEcp1B* (GenBank accession no. AF458080), respectively, whereas its imperfect inverted repeat left (IRL) and right (IRR1) sequences (2 bp mismatches) were identical to those of *ISEcp1* and *ISEcp1B* (Fig. 1). *ISEcp1*-like elements have been identified at the 5' end of several β-lactamase genes, such as the *bla*_{CTX-M}, *bla*_{CMY}, and *bla*_{ACC} genes, and associated with the 16S rRNA methyltransferase gene *rmtC*, enabling those genes to be transposed (7, 14, 17, 22, 23, 29).

The *qnrB19* gene was part of a 2,739-bp potential transposon flanked by a 5-bp duplication of the target site (ATCAA), likely evidence of a transposition event (Fig. 1). This potential transposon (named Tn2012), comprising *ISEcp1C* and *qnrB19*, was inserted inside the *orf1* gene of the transposon Tn1721 (Fig. 1). Tn1721 is an 11.1-kb nonconjugative transposon belonging to the Tn21 subgroup of the Tn3 family of bacterial transposons that confers inducible tetracycline resistance (1). The novel transposon Tn2012 was bracketed by two imperfect

14-bp IR sequences (seven mismatches), namely, the IRL of *ISEcp1C* and an IRR named IRR2 (Fig. 1). This IRR2 sequence shared 7 out of 14 bp with the original IRR1 of *ISEcp1C*. This observation is in accordance with previous studies (14, 22, 23), showing that *ISEcp1B* was able to use as IRRs sequences sharing weak identity with its original IRR1, corresponding to a one-ended transposition mechanism.

Transposition experiments were performed as previously described (14, 22) in order to determine whether *ISEcp1C* was able to mobilize the *qnrB19* gene. Briefly, the sequence of the entire Tn2012 transposon was amplified by PCR using primers *orf1*-A (5'-CGACAACGGATATTCAAAGC-3') and *orf1*-B (5'-ACTTTGCAAATTATTCTGCC-3') and then cloned into the kanamycin-resistant pCR-BluntII-TOPO plasmid (Invitrogen). This recombinant plasmid, first transferred and selected in *E. coli* TOP10 (Invitrogen), was used for the electrotransformation of *E. coli* RZ211 (pOX38-Gm). Plasmid pOX38-Gm is a self-conjugative and insertion element-free plasmid carrying a gentamicin resistance marker. The transposition of Tn2012 from the recombinant pCR-BluntII-TOPO derivative to plasmid pOX38-Gm was investigated after 24 h of growth in Trypticase soy broth by mating RZ211 with azide-resistant *E. coli* J53 and selecting for transconjugants growing on agar plates containing 8 μg of gentamicin per ml, 6 μg of




FIG. 1. Schematic map of the structure of transposon Tn2012 identified on the pR4525 natural plasmid of *E. coli* R4525. Open reading frames are shown as arrows or horizontal boxes with an arrow indicating the orientation of the coding sequence with the gene name above the corresponding boxes. The IRL, IRR1, and IRR2 motifs are indicated (blackened bp are identical, whereas whitened bp are different), and target site duplications (ATCAA) are represented by black bars.



FIG. 2. Nucleotide sequence of the upstream region of the *qnrB19* gene. The -35 and -10 promoter elements are indicated, and the transcription start site is represented by an arrow. The former start codon corresponds to the first ATG that was initially proposed for QnrB1 (226 amino acids) (11, 12).

nalidixic acid per ml, and 100 μ g of azide per ml. The plasmid DNA of several nalidixic acid-resistant transconjugants was extracted and pOX38-Gm sequencing confirmed the *ISEcp1C*-mediated transposition of *qnrB19*. The transposition frequency, calculated by dividing the number of transconjugants by the number of donors, was estimated to be 10^{-6} to 10^{-7} per donor cell. Moreover, an analysis of five insertion events of *ISEcp1C-qnrB19* sequences in plasmid pOX38-Gm showed that transposition had occurred into five different sites distantly located on the recipient plasmid. Alignment of the target sites revealed variable sequences (ATTAT, ATTAC, TCATA, TACAT, and TTCAT), exhibiting an AT-rich content as previously observed (14, 22).

It has been shown that *ISEcp1*-like elements may bring promoter sequences for the high-level expression of downstream-located β -lactamase genes (17, 23, 29). However, as opposed to what has been reported for those genes, the *qnrB19* gene was located in an opposite orientation with respect to the transposase gene of *ISEcp1C* (Fig. 1). The promoter sequences for the *qnrB19* expression were determined by using the 5' rapid amplification of cDNA ends technique (Invitrogen). Total RNA was extracted from cultures of *E. coli* R4525 by using the RNeasy Protect mini kit (Qiagen, Courtaboeuf, France). The +1 transcription start site was identified 28 bp upstream of an ATG codon located inside the presumed *qnrB19* gene. This prompted us to consider the *qnrB19* gene to be only 645 bp long and the QnrB19 protein to be 214 amino acids long, not 226 amino acids long as previously considered for QnrB1 (11, 12). This result is in accordance with other observations showing that the putative QnrB1 sequence was longer than the QnrA and QnrS sequences but also than other chromosome-encoded Qnr-like sequences from *Vibrionaceae* (11, 21). The deduced promoter region based on the +1 start site identified a -35 box (TTGACG) and a -10 box (TACCAT) separated by a 17-bp sequence (Fig. 2).

We have demonstrated here that an *ISEcp1* element was at the origin of the acquisition of a *qnrB*-like gene (*qnrB19*) and that it was not involved in the expression of that gene, as opposed to what has been reported for other antibiotic resistance genes. Although several *qnrB*-like genes have been reported to be associated with *ISCR1*, we report here a novel genetic structure responsible for *qnrB* acquisition and dissemination. It remains to be determined what could be the natural

reservoir of *ISEcp1*-like elements since it is associated with structurally unrelated resistance determinants of various origins, such as the *bla*_{CTX-M} and *bla*_{CMY} genes from *Enterobacteriaceae* and the *qnr*-like genes from *Vibrionaceae* and *Shewanellaceae* (4, 20).

Nucleotide sequence accession number. The nucleotide sequence of the *qnrB19* gene and that of Tn2012 shown in Fig. 1 were submitted to the GenBank database and can be found under accession no. EU432277 and EU523120, respectively.

This work was financed by a grant from the Ministère de l'Éducation Nationale et de la Recherche (grant UPRES, EA3539), Université Paris XI, Paris, France, and by the European Community (6th PCRD, LSHM-CT-2005-018705).

REFERENCES

- Allmeier, H., B. Cresnar, M. Greck, and R. Schmitt. 1992. Complete nucleotide sequence of Tn1721: gene organization and a novel gene product with features of a chemotaxis protein. *Gene* 111:11–20.
- Cattoir, V., P. Lesprit, C. Lasclos, E. Denamur, P. Legrand, C. J. Soussy, and E. Cambau. 2006. In vivo selection during ofloxacin therapy of *Escherichia coli* with combined topoisomerase mutations that confer high resistance to ofloxacin but susceptibility to nalidixic acid. *J. Antimicrob. Chemother.* 58:1054–1057.
- Cattoir, V., L. Poirel, C. Aubert, C. J. Soussy, and P. Nordmann. 2008. Unexpected occurrence of plasmid-mediated quinolone resistance determinants in environmental *Aeromonas* spp. *Emerg. Infect. Dis.* 14:231–237.
- Cattoir, V., L. Poirel, D. Mazel, C. J. Soussy, and P. Nordmann. 2007. *Vibrio splendidus* as the source of plasmid-mediated QnrS-like quinolone resistance determinants. *Antimicrob. Agents Chemother.* 51:2650–2651.
- Cattoir, V., L. Poirel, V. Rotimi, C. J. Soussy, and P. Nordmann. 2007. Multiplex PCR for detection of plasmid-mediated quinolone resistance *qnr* genes in ESBL-producing enterobacterial isolates. *J. Antimicrob. Chemother.* 60:394–397.
- Clinical and Laboratory Standards Institute. 2006. Performance standards for antimicrobial disk susceptibility tests. Approved standard, M2–A9, 9th ed. Clinical and Laboratory Standards Institute, Wayne, PA.
- Doloy, A., C. Verdet, V. Gautier, D. Decré, E. Ronco, A. Hammami, A. Philippon, and G. Arlet. 2006. Genetic environment of acquired *bla*_{ACC-1} β -lactamase gene in *Enterobacteriaceae* isolates. *Antimicrob. Agents Chemother.* 50:4177–4181.
- Garnier, F., N. Raked, A. Gassama, F. Denis, and M.-C. Ploy. 2006. Genetic environment of quinolone resistance gene *qnrB2* in a complex *sul1*-type integron in the newly described *Salmonella enterica* serovar Keurmassar. *Antimicrob. Agents Chemother.* 50:3200–3202.
- Gay, K., A. Robicsek, J. Strahilevitz, C. H. Park, G. Jacoby, T. J. Barrett, F. Medalla, T. M. Chiller, and D. C. Hooper. 2006. Plasmid-mediated quinolone resistance in non-Typhi serotypes of *Salmonella enterica*. *Clin. Infect. Dis.* 43:297–304.
- Hata, M., M. Suzuki, M. Matsumoto, M. Takahashi, K. Sato, S. Ibe, and K. Sakae. 2005. Cloning of a novel gene for quinolone resistance from a transferable plasmid in *Shigella flexneri* 2b. *Antimicrob. Agents Chemother.* 49:801–803.
- Jacoby, G., V. Cattoir, D. Hooper, L. Martinez-Martinez, P. Nordmann, A.

- Pascual, L. Poirel, and M. Wang. 2008. *qnr* gene nomenclature. *Antimicrob. Agents Chemother.* **52**:2297–2299.
12. Jacoby, G. A., K. E. Walsh, D. M. Mills, V. J. Walker, H. Oh, A. Robicsek, and D. C. Hooper. 2006. *qnrB*, another plasmid-mediated gene for quinolone resistance. *Antimicrob. Agents Chemother.* **50**:1178–1182.
 13. Kieser, T. 1984. Factors affecting the isolation of CCC DNA from *Streptomyces lividans* and *Escherichia coli*. *Plasmid* **12**:19–36.
 14. Lartigue, M. F., L. Poirel, D. Aubert, and P. Nordmann. 2006. In vitro analysis of *ISEcp1B*-mediated mobilization of naturally occurring β -lactamase gene *bla_{CTX-M}* of *Kluyvera ascorbata*. *Antimicrob. Agents Chemother.* **50**:1282–1286.
 15. Martinez-Martinez, L., A. Pascual, and G. A. Jacoby. 1998. Quinolone resistance from a transferable plasmid. *Lancet* **351**:797–799.
 16. Minarini, L. A., L. Poirel, V. Cattoir, A. L. Darini, and P. Nordmann. Plasmid-mediated quinolone resistance determinants among enterobacterial isolates from outpatients in Brazil. *J. Antimicrob. Chemother.*, in press.
 17. Nakano, R., R. Okamoto, N. Nagano, and M. Inoue. 2007. Resistance to gram-negative organisms due to high-level expression of plasmid-encoded *ampC* β -lactamase *bla_{CMY-4}* promoted by insertion sequence *ISEcp1*. *J. Infect. Chemother.* **13**:18–23.
 18. Nordmann, P., and L. Poirel. 2005. Emergence of plasmid-mediated resistance to quinolones in *Enterobacteriaceae*. *J. Antimicrob. Chemother.* **56**:463–469.
 19. Périchon, B., P. Courvalin, and M. Galimand. 2007. Transferable resistance to aminoglycosides by methylation of G1405 in 16S rRNA and to hydrophilic fluoroquinolones by QepA-mediated efflux in *Escherichia coli*. *Antimicrob. Agents Chemother.* **51**:2464–2469.
 20. Poirel, L., J. M. Rodriguez-Martinez, H. Mammeri, A. Liard, and P. Nordmann. 2005. Origin of plasmid-mediated quinolone resistance determinant QnrA. *Antimicrob. Agents Chemother.* **49**:3523–3525.
 21. Poirel, L., A. Liard, J. M. Rodriguez-Martinez, and P. Nordmann. 2005. *Vibrionaceae* as a possible source of Qnr-like quinolone resistance determinants. *J. Antimicrob. Chemother.* **56**:1118–1121.
 22. Poirel, L., M. F. Lartigue, J. W. Decusser, and P. Nordmann. 2005. *ISEcp1B*-mediated transposition of *bla_{CTX-M}* in *Escherichia coli*. *Antimicrob. Agents Chemother.* **49**:447–450.
 23. Poirel, L., J. W. Decusser, and P. Nordmann. 2003. Insertion sequence *ISEcp1B* is involved in the expression and mobilization of a *bla_{CTX-M}* β -lactamase gene. *Antimicrob. Agents Chemother.* **47**:2938–2945.
 24. Quiroga, M. P., P. Andres, A. Petroni, A. J. C. S. Bistué, L. Guerriero, L. J. Vargas, A. Zorreguieta, M. Tokumoto, C. Quiroga, M. E. Tolmasky, M. Galas, and D. Centron. 2007. Complex class 1 integrons with diverse variable regions, including *aac(6')-Ib-cr*, and a novel allele, *qnrB10*, associated with *ISCR1* in clinical enterobacterial isolates from Argentina. *Antimicrob. Agents Chemother.* **51**:4466–4470.
 25. Robicsek, A., J. Strahilevitz, G. A. Jacoby, M. Macielag, D. Abbanat, C. H. Park, K. Bush, and D. C. Hooper. 2006. Fluoroquinolone-modifying enzyme: a new adaptation of a common aminoglycoside acetyltransferase. *Nat. Med.* **12**:83–88.
 26. Robicsek, A., G. A. Jacoby, and D. C. Hooper. 2006. The worldwide emergence of plasmid-mediated quinolone resistance. *Lancet Infect. Dis.* **6**:629–640.
 27. Ruiz, J. 2003. Mechanisms of resistance to quinolones: target alterations, decreased accumulation and DNA gyrase protection. *J. Antimicrob. Chemother.* **51**:1109–1117.
 28. Toleman, M. A., P. M. Bennett, and T. R. Walsh. 2006. *ISCR* elements: novel gene-capturing systems of the 21st century? *Microbiol. Mol. Biol. Rev.* **70**:296–316.
 29. Wachino, J., K. Yamane, K. Kimura, N. Shibata, S. Suzuki, Y. Ike, and Y. Arakawa. 2006. Mode of transposition and expression of 16S rRNA methyltransferase gene *rmtC* accompanied by *ISEcp1*. *Antimicrob. Agents Chemother.* **50**:3212–3215.
 30. Yamane, K., J. Wachino, S. Suzuki, K. Kimura, N. Shibata, H. Kato, K. Shibayama, T. Konda, and Y. Arakawa. 2007. New plasmid-mediated fluoroquinolone efflux pump, QepA, found in an *Escherichia coli* clinical isolate. *Antimicrob. Agents Chemother.* **51**:3354–3360.