

Contribution of *dfrA* and *inhA* Mutations to the Detection of Isoniazid-Resistant *Mycobacterium tuberculosis* Isolates[∇]

Yu Min Ho,¹ Yong-Jiang Sun,² Sin-Yew Wong,² and Ann S. G. Lee^{1,3*}

Division of Medical Sciences, National Cancer Centre, Singapore 169610¹; Department of Infectious Disease, Tan Tock Seng Hospital, Singapore 308433²; and Department of Microbiology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 117597³

Received 31 March 2009/Returned for modification 28 April 2009/Accepted 25 June 2009

Screening of 127 isoniazid (INH)-resistant *Mycobacterium tuberculosis* isolates from Singapore for mutations within the *dfrA* and *inhA* genes revealed mutations in 0 and 5 (3.9%) isolates respectively, implying that mutations in *dfrA* do not contribute to the detection of INH-resistant *M. tuberculosis* and that mutations within *inhA* are rare. Thirty-seven (29%) of the 127 isolates had no mutations in any of the genes implicated in INH resistance (*katG*, *kasA*, and *ndh*; *inhA* and *ahpC* promoters), suggesting that there are new INH targets yet to be discovered.

Mycobacterium tuberculosis remains a major health concern worldwide, with approximately 2 billion people currently infected worldwide and 9.2 million new cases in 2006 (13, 27). The emergence of multidrug-resistant tuberculosis has increased global efforts to understand the molecular mechanisms of drug resistance in *M. tuberculosis*.

Isoniazid (INH) for *M. tuberculosis* chemotherapy has always been used as first-line treatment (25) since it was found to exhibit powerful bactericidal activity against the disease by effectively causing loss of acid fastness in *M. tuberculosis* through inhibition of mycolic acid synthesis (9). However, its widespread use, justified by its high specificity, low cost, and considerably low toxicity, has seen treatment failures due to increasing resistance to the drug.

Clinical resistance to INH is widely known to be caused by mutations within the *katG* and *inhA* genes and the promoter region of *inhA* (2, 13, 18, 19, 25, 29). Other genes, such as *kasA* and *ndh*, have been implicated in INH resistance, but mutations in these genes are rare and have been observed in INH-susceptible isolates and/or in association with *katG* mutations

(8, 11, 12, 14, 15, 18, 25, 28). The *ahpC* gene is involved in the cellular regulation of oxidative stress (18, 22). Mutations within the *oxyR-ahpC* intergenic region that result in increased expression of alkyl hydroperoxidase are considered to compensate for peroxide sensitivity due to loss of KatG function found in clinical resistant strains (10, 22, 28). Nevertheless, screening of INH-resistant clinical isolates of *M. tuberculosis* for mutations within *kasA*, *ndh*, and *ahpC* is often performed.

We have previously extensively characterized INH-resistant isolates for mutations within all of these regions; however, 57 (36%) of 160 INH-resistant isolates from Singapore had no detectable alterations (11, 12). This finding suggests that mutations in other genes that confer resistance to INH may exist.

Recently, dihydrofolate reductase, encoded by the *dfrA* gene, was proposed as a new target for INH (1) as it has been reported that NADP-bound INH had been shown to inhibit dihydrofolate reductase, an enzyme essential for nucleic acid synthesis. Also, the importance of mutations within the structural region of enoyl reductase (*InhA*), encoded by the *inhA* gene, have been reported where a single point mutation allele

TABLE 1. Oligonucleotide primers used for the amplification of the entire *dfrA*^a and *inhA*^b genes

Gene	Primer sequence	Annealing temp (°C)	Amplicon size (bp)	Nucleotide positions
<i>dfrA</i> (F)	GAC GAA GCG ATG AGG AGA AG	55	632	300,058–300,072
<i>dfrA</i> (R)	TCG TTG TGA AGA ACT ACG ATC C			300,699–300,678
<i>inhA</i> (F)1	CTA CAT CGA CAC CGA TAT GAC	55	700	290,948–290,968
<i>inhA</i> (R)1	GAC CGT CAT CCA GTT GTA G			291,647–291,629
<i>inhA</i> (F)2	GCA TCA ACC CGT TCT TCG AC	55	677	291,469–291,488
<i>inhA</i> (R)2	TAA TGC CAT TGA TCG GTG ATA C			292,145–292,124

^a The *M. tuberculosis* sequence used to design the primers was obtained from GenBank (accession no. BX842580.1).

^b The *M. tuberculosis* sequence used to design the primers was obtained from GenBank (accession no. BX842576.1).

* Corresponding author. Mailing address: Division of Medical Sciences, National Cancer Centre, Singapore, 11 Hospital Drive, Singapore 169610, Republic of Singapore. Phone: 65 6436-8313. Fax: 65 6372-0161. E-mail: dmslsg@nccs.com.sg.

[∇] Published ahead of print on 6 July 2009.

TABLE 2. *M. tuberculosis* isolates with mutations in the *inhA* gene

Isolate	Mutation	Amino acid substitution	No. (%) of isolates	Mutation(s) at:				
				<i>katG</i>	<i>ahpC</i>	<i>inhA</i> promoter	<i>kasA</i>	<i>ndh</i>
I59	GGA→GGC	G3G	1 (0.8)					
I82	ATC→ACC	I194T	1 (0.8)					
IR27	GGA→GGC	G3G	1 (0.8)					
IRS7	ATC→ACC	I21T	1 (0.8)		-6 (G→A) ^a 27 (G→T) ^b	-15 (C→T)		
MDR2	ATC→ACC	I21T	1 (0.8)			-15 (C→T)		

^a Position relative to the mRNA start site.

^b Nucleotide substitution within the defective *oxyR* gene.

(S94A) transferred by using specialized linkage transduction was able to confer clinically relevant levels of resistance and inhibit mycolic acid synthesis (18, 26). Most studies investigating mutations in the *inhA* gene analyzed the promoter region of *inhA* and not the region within the gene (6–8, 16, 17). Thus, we aimed to screen for mutations within both the *inhA* and *dfrA* genes of INH-resistant *M. tuberculosis* isolates to determine whether mutations within these genes contribute to INH resistance.

A total of 127 INH-resistant clinical isolates and 15 INH-sensitive isolates from Singapore were included in this study in order to gain molecular insight into INH resistance. DNA extracted from the isolates was analyzed by amplifying two overlapping fragments for the entire *inhA* gene and one fragment for the entire *dfrA* gene (Table 1). Amplification of the genes through PCR was done by using specific oligonucleotide primers for the respective gene fragments (Table 1). PCR products were purified (NucleoSpin Extract Column II; Macherey-Nagel) and directly sequenced with the BigDye Terminator sequencing kit V3.1 (Applied Biosystems), followed by analysis on the 3130XL Genetic Analyzer (Applied Biosystems). Confirmation of mutations was done by reamplification and resequencing. The nucleotide sequences obtained were aligned against the reference sequences of the respective genes of *M. tuberculosis* reference strain H37Rv by using the DNASTAR SeqMan II software (Lasergene).

Screening of the entire *inhA* gene revealed mutations in five (3.9%) of 127 INH-resistant isolates (synonymous mutation GGA→GGC at nucleotide position 9 or G3G [$n = 2$], I21T [$n = 2$], or I194T [$n = 1$]). Of the five isolates with mutations, four also had mutations in the *ahpC* promoter, the *inhA* promoter, and the *kasA* gene, which were detected in our previous study (Table 2) (11). Such an occurrence might suggest that some of the observed substitutions are not unique mutations that actually contribute to INH resistance. Also, the low mutational frequency within the *inhA* gene among INH-resistant isolates suggests this (11). Furthermore, a previous study reported that a previously known mutation, at position 491 of the *rrs* gene, was in fact a polymorphism within the F11 family and not actually involved in streptomycin resistance (24). Hence, mutations found, especially rarely occurring ones, should not be hastily perceived to confer resistance.

Screening of the entire *dfrA* gene did not reveal any mutations in any of the 127 INH-resistant isolates, suggesting that *dfrA* mutations may not contribute to INH resistance and that screening for *dfrA* mutations in INH-resistant isolates is unnecessary.

To determine if the mutations detected within the *inhA* gene are mutations and not polymorphisms, we screened 15 INH-susceptible isolates and found no mutations in *inhA*. These isolates also did not have mutations in the *dfrA* gene.

Mutations at both I21T (3, 5, 18) and I194T (5, 13) were previously reported to confer resistance to INH. Ile21 and Ile194 were shown to be 2 of the 10 most important amino acid residues making conserved hydrogen bonds with NADH cofactor in the wild-type InhA protein, which encourages INH sensitivity (21). Amino acid replacements will result in lowered NADH affinity and therefore confer resistance to INH since InhA inactivation is dependent on the presence of bound NADH (4, 13, 20, 23, 28).

Of the 127 INH-resistant samples we have screened, there remain 37 samples (29%) with no mutations in any of the genes reported to be involved in INH resistance. This possibly implies that there are new INH targets yet to be discovered. Our findings suggest that mutations in *dfrA* do not contribute to the detection of INH-resistant *M. tuberculosis* and that mutations within the *inhA* gene are rare.

We acknowledge the Central Tuberculosis Laboratory, Department of Pathology, Singapore General Hospital, for providing isolates.

This work was funded by a Singhealth Cluster Research Grant (SHF/FG346P/2006).

REFERENCES

- Argyrou, A., M. W. Vetting, B. Aladegbami, and J. S. Blanchard. 2006. *Mycobacterium tuberculosis* dihydrofolate reductase is a target for isoniazid. *Nat. Struct. Mol. Biol.* **13**:408–413.
- Banerjee, A., E. Dubnau, A. Quemard, V. Balasubramanian, K. S. Um, T. Wilson, D. Collins, G. de Lisle, and W. R. Jacobs, Jr. 1994. *inhA*, a gene encoding a target for isoniazid and ethionamide in *Mycobacterium tuberculosis*. *Science* **263**:227–230.
- Basso, L. A., and J. S. Blanchard. 1998. Resistance to antitubercular drugs. *Adv. Exp. Med. Biol.* **456**:115–144.
- Basso, L. A., R. Zheng, J. M. Musser, W. R. Jacobs, Jr., and J. S. Blanchard. 1998. Mechanisms of isoniazid resistance in *Mycobacterium tuberculosis*: enzymatic characterization of enoyl reductase mutants identified in isoniazid-resistant clinical isolates. *J. Infect. Dis.* **178**:769–775.
- Brossier, F., N. Veziris, C. Truffot-Pernot, V. Jarlier, and W. Sougakoff. 2006. Performance of the genotype MTBDR line probe assay for detection of resistance to rifampin and isoniazid in strains of *Mycobacterium tuberculosis* with low- and high-level resistance. *J. Clin. Microbiol.* **44**:3659–3664.
- Caws, M., P. M. Duy, D. Q. Tho, N. T. Lan, D. V. Hoa, and J. Farrar. 2006. Mutations prevalent among rifampin- and isoniazid-resistant *Mycobacterium tuberculosis* isolates from a hospital in Vietnam. *J. Clin. Microbiol.* **44**:2333–2337.
- Guo, H., Q. Seet, S. Denkin, L. Parsons, and Y. Zhang. 2006. Molecular characterization of isoniazid-resistant clinical isolates of *Mycobacterium tuberculosis* from the USA. *J. Med. Microbiol.* **55**:1527–1531.
- Hazbón, M. H., M. Brimacombe, M. Bobadilla del Valle, M. Cavatore, M. I. Guerrero, M. Varma-Basil, H. Billman-Jacobe, C. Lavender, J. Fyfe, L. Garcia-Garcia, C. I. Leon, M. Bose, F. Chaves, M. Murray, K. D. Eisenach, J. Sifuentes-Osornio, M. D. Cave, A. Ponce de Leon, and D. Alland. 2006. Population genetics study of isoniazid resistance mutations and evolution of

- multidrug-resistant *Mycobacterium tuberculosis*. Antimicrob. Agents Chemother. **50**:2640–2649.
9. Johnson, R., E. M. Streicher, G. E. Louw, R. M. Warren, P. D. van Helden, and T. C. Victor. 2006. Drug resistance in *Mycobacterium tuberculosis*. Curr. Issues Mol. Biol. **8**:97–111.
 10. Kelley, C. L., D. A. Rouse, and S. L. Morris. 1997. Analysis of *ahpC* gene mutations in isoniazid-resistant clinical isolates of *Mycobacterium tuberculosis*. Antimicrob. Agents Chemother. **41**:2057–2058.
 11. Lee, A. S., I. H. Lim, L. L. Tang, A. Telenti, and S. Y. Wong. 1999. Contribution of *kasA* analysis to detection of isoniazid-resistant *Mycobacterium tuberculosis* in Singapore. Antimicrob. Agents Chemother. **43**:2087–2089.
 12. Lee, A. S., A. S. Teo, and S. Y. Wong. 2001. Novel mutations in *ndh* in isoniazid-resistant *Mycobacterium tuberculosis* isolates. Antimicrob. Agents Chemother. **45**:2157–2159.
 13. Leung, E. T., P. L. Ho, K. Y. Yuen, W. L. Woo, T. H. Lam, R. Y. Kao, W. H. Seto, and W. C. Yam. 2006. Molecular characterization of isoniazid resistance in *Mycobacterium tuberculosis*: identification of a novel mutation in *inhA*. Antimicrob. Agents Chemother. **50**:1075–1078.
 14. Mdluli, K., R. A. Slayden, Y. Zhu, S. Ramaswamy, X. Pan, D. Mead, D. D. Crane, J. M. Musser, and C. E. Barry III. 1998. Inhibition of a *Mycobacterium tuberculosis* β -ketoacyl ACP synthase by isoniazid. Science **280**:1607–1610.
 15. Miesel, L., T. R. Weisbrod, J. A. Marcinkeviciene, R. Bittman, and W. R. Jacobs, Jr. 1998. NADH dehydrogenase defects confer isoniazid resistance and conditional lethality in *Mycobacterium smegmatis*. J. Bacteriol. **180**:2459–2467.
 16. Morris, S., G. H. Bai, P. Suffys, L. Portillo-Gomez, M. Fairchok, and D. Rouse. 1995. Molecular mechanisms of multiple drug resistance in clinical isolates of *Mycobacterium tuberculosis*. J. Infect. Dis. **171**:954–960.
 17. Park, H., E. J. Song, E. S. Song, E. Y. Lee, C. M. Kim, S. H. Jeong, J. H. Shin, J. Jeong, S. Kim, Y. K. Park, G. H. Bai, and C. L. Chang. 2006. Comparison of a conventional antimicrobial susceptibility assay to an oligonucleotide chip system for detection of drug resistance in *Mycobacterium tuberculosis* isolates. J. Clin. Microbiol. **44**:1619–1624.
 18. Ramaswamy, S., and J. M. Musser. 1998. Molecular genetic basis of antimicrobial agent resistance in *Mycobacterium tuberculosis*: 1998 update. Tuberc. Lung Dis. **79**:3–29.
 19. Rouse, D. A., Z. Li, G. H. Bai, and S. L. Morris. 1995. Characterization of the *katG* and *inhA* genes of isoniazid-resistant clinical isolates of *Mycobacterium tuberculosis*. Antimicrob. Agents Chemother. **39**:2472–2477.
 20. Rozwarski, D. A., G. A. Grant, D. H. Barton, W. R. Jacobs, Jr., and J. C. Sacchettini. 1998. Modification of the NADH of the isoniazid target (InhA) from *Mycobacterium tuberculosis*. Science **279**:98–102.
 21. Schroeder, E. K., L. A. Basso, D. S. Santos, and O. N. de Souza. 2005. Molecular dynamics simulation studies of the wild-type, I21V, and I16T mutants of isoniazid-resistant *Mycobacterium tuberculosis* enoyl reductase (InhA) in complex with NADH: toward the understanding of NADH-InhA different affinities. Biophys. J. **89**:876–884.
 22. Sherman, D. R., K. Mdluli, M. J. Hickey, T. M. Arain, S. L. Morris, C. E. Barry III, and C. K. Stover. 1996. Compensatory *ahpC* gene expression in isoniazid-resistant *Mycobacterium tuberculosis*. Science **272**:1641–1643.
 23. Telenti, A., N. Honore, C. Bernasconi, J. March, A. Ortega, B. Heym, H. E. Takiff, and S. T. Cole. 1997. Genotypic assessment of isoniazid and rifampin resistance in *Mycobacterium tuberculosis*: a blind study at reference laboratory level. J. Clin. Microbiol. **35**:719–723.
 24. Victor, T. C., A. van Rie, A. M. Jordaan, M. Richardson, G. D. van Der Spuy, N. Beyers, P. D. van Helden, and R. Warren. 2001. Sequence polymorphism in the *rrs* gene of *Mycobacterium tuberculosis* is deeply rooted within an evolutionary clade and is not associated with streptomycin resistance. J. Clin. Microbiol. **39**:4184–4186.
 25. Vilchèze, C., and W. R. Jacobs, Jr. 2007. The mechanism of isoniazid killing: clarity through the scope of genetics. Annu. Rev. Microbiol. **61**:35–50.
 26. Vilchèze, C., F. Wang, M. Arai, M. H. Hazbon, R. Colangeli, L. Kremer, T. R. Weisbrod, D. Alland, J. C. Sacchettini, and W. R. Jacobs, Jr. 2006. Transfer of a point mutation in *Mycobacterium tuberculosis inhA* resolves the target of isoniazid. Nat. Med. **12**:1027–1029.
 27. World Health Organization. 2008, posting date. Global tuberculosis control: surveillance, planning, financing. World Health Organization, Geneva, Switzerland. http://www.who.int/tb/publications/global_report/2008/pdf/full_report.pdf.
 28. Zhang, M., J. Yue, Y. P. Yang, H. M. Zhang, J. Q. Lei, R. L. Jin, X. L. Zhang, and H. H. Wang. 2005. Detection of mutations associated with isoniazid resistance in *Mycobacterium tuberculosis* isolates from China. J. Clin. Microbiol. **43**:5477–5482.
 29. Zhang, Y., B. Heym, B. Allen, D. Young, and S. Cole. 1992. The catalase-peroxidase gene and isoniazid resistance of *Mycobacterium tuberculosis*. Nature **358**:591–593.