

## AUTHOR'S CORRECTION

### Origin and Molecular Evolution of the Determinant of Methicillin Resistance in Staphylococci

Sae Tsubakishita, Kyoko Kuwahara-Arai, Takashi Sasaki, and Keiichi Hiramatsu

*Department of Infection Control Science, Department of Microbiology, Faculty of Medicine,  
Juntendo University, 2-1-1 Hongo, Bunkyo-ku, Tokyo 113-8421, Japan*

Volume 54, no. 10, p. 4352–4359, 2010. Page 4353, column 2: The following paragraph should appear before Results and Discussion.

**Nucleotide sequence accession numbers.** The nucleotide sequences of *S. fleurettii* strains CCUG 43834<sup>T</sup>, SFMP08, SFMN01, and SFMN02, *S. sciuri* strains ATCC 29062<sup>T</sup>, ATCC 700061<sup>T</sup>, and ATCC 700058<sup>T</sup>, *S. vitulinus* strain SVMP01, *S. lentus* strain ATCC 29070<sup>T</sup>, *S. xylosus* strain ATCC 29971<sup>T</sup>, and *S. kloosii* strain DSM 20676<sup>T</sup> have been deposited in the DDBJ/EMBL/GenBank databases under accession no. AB546266, AB546267, AB546268, AB546269, AB547234, AB547235, AB547236, AB546780, AB546859, AB545964, and AB545965, respectively.