

New Small Plasmid Harboring *bla*_{KPC-2} in *Pseudomonas aeruginosa*

Renata Galetti,^a Leonardo Neves Andrade,^a Michael Chandler,^c Alessandro de Mello Varani,^b Ana Lúcia Costa Darini^a

Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, Brazil^a; Faculdade de Ciências Agrárias e Veterinárias, Universidade Estadual Paulista Júlio de Mesquita Filho, Jaboticabal, Brazil^b; Laboratoire de Microbiologie et Génétique Moléculaires, Centre National de la Recherche Scientifique, Toulouse, France^c

The aim of this study was to characterize the genetic context of *bla*_{KPC-2} in *Pseudomonas aeruginosa* sequence type 244 from Brazil. The *bla*_{KPC-2} gene was detected in a new small plasmid, pBH6. Complete sequencing revealed that pBH6 was 3,652 bp long and included the Tn3 resolvase and Tn3 inverted repeat (IR), a partial copy of *ISKpn6*, and a putative *ori* region but no *rep* genes. pBH6 replicated stably into *Escherichia coli* strain DH10B and *P. aeruginosa* strain PAO.

Carbapenem resistance mediated by the production of *Klebsiella pneumoniae* carbapenemase (KPC) enzymes has been reported worldwide in *Enterobacteriaceae*, and the *bla*_{KPC-2} gene was found to be associated with different transposons and plasmids (1–3). However, *bla*_{KPC} has also emerged in clinical isolates of *Pseudomonas aeruginosa*. First reported in Colombia in 2006 (4), subsequent worldwide reports described plasmid and chromosomal genetic contexts for this gene. In Brazil, KPC-producing *P. aeruginosa* was first described in 2012 (5), followed by a second report (6) that highlighted a worrying increase in KPC-producing *P. aeruginosa*. However, the genetic context of *bla*_{KPC} in the Brazilian strains remained unknown. At present, two complete sequences of *P. aeruginosa bla*_{KPC-2}-carrying plasmids, pCol-1 (GenBank accession number [KC609323](#)) from *P. aeruginosa* sequence type 308 (ST308) and pPA-2 (GenBank accession number [KC609322](#)) from *P. aeruginosa* ST1006 (7), are available in public databases. The aim of this study was to characterize the genetic context of *bla*_{KPC-2} in a *P. aeruginosa* isolate from Brazil. Carbapenem-resistant *P. aeruginosa* (referred to here as BH6) was isolated in Belo Horizonte, Brazil, in 2011. BH6 was recovered from tracheal secretions of a female patient in the intensive care unit of a tertiary medical care center. This isolate was characterized as multidrug resistant, KPC-2 producing, and of ST244 (8). MIC values were ≥ 32 $\mu\text{g/ml}$ for imipenem, ≥ 32 $\mu\text{g/ml}$ for meropenem, ≥ 256 $\mu\text{g/ml}$ for ceftazidime, ≥ 256 $\mu\text{g/ml}$ for aztreonam, ≥ 256 $\mu\text{g/ml}$ for cefepime, and 1.0 $\mu\text{g/ml}$ for polymyxin B (9, 10). The *bla*_{KPC-2} gene was isolated by PCR amplification, as previously described (11), followed by Sanger sequencing. The ST was characterized according to the *P. aeruginosa* MLST database guideline (see <http://pubmlst.org/paeruginosa/>).

The genomic location of *bla*_{KPC-2} was determined, as previously described, by hybridization on I-CeuI pulsed-field gel electrophoresis (PFGE) using *bla*_{KPC-2} and 16S rRNA genes as probes (to determine whether it was located on the chromosome) and S1 PFGE using *bla*_{KPC-2} as a probe (to determine whether it was plasmid associated) (12). We observed three bands in the gel after S1 PFGE, and all three of them hybridized with the *bla*_{KPC-2} probe. Since it seemed possible that they represented the linear, open-circular, and closed-circular forms of a single plasmid, plasmid DNA was extracted and purified using a PureLink HiPure plasmid filter midprep kit (Invitrogen) and digested with BamHI (Thermo Fisher Scientific). This treatment generated a single plasmid band that indicated the presence of a single plasmid species. To determine whether this plasmid, pBH6, belonged to a recog-




FIG 1 Circular plasmid carrying the *bla*_{KPC-2} and origin of replication (*ori*) genes. The partial copy of *ISKpn6* carrying the IRR (black arrow) is shown. A Tn3 IR is located between the *bla*_{KPC-2} and the Tn3 resolvase (longer black arrow).

nized incompatibility (Inc) group, we used a PCR-based replicon typing (PBRT) scheme (13); however, we were unable to identify a particular replicon by using this procedure. Finally, plasmid DNA was sequenced using the Ion PGM system (Life Technologies). *De novo* assembly was carried out using the CLC Genomics work-

Received 29 January 2016 Returned for modification 29 January 2016

Accepted 2 March 2016

Accepted manuscript posted online 7 March 2016

Citation Galetti R, Andrade LN, Chandler M, Varani ADM, Darini ALC. 2016. New small plasmid harboring *bla*_{KPC-2} in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 60:3211–3214. doi:10.1128/AAC.00247-16.

Address correspondence to Renata Galetti, regaletti@hotmail.com.

Copyright © 2016, American Society for Microbiology. All Rights Reserved.




FIG 2 Similarities among pBH6 (3,642 bp), pCol-1 (31,529 bp, coordinates 17066 to 19135), pPA2 (7,995 bp, coordinates 1133 to 3283), pFOS18 (23,939 bp, coordinates 4910 to 11956), and pKP13d (45,574 bp, coordinates 4910 to 11956). Red lines indicate the region with 100% similarity among the 5 plasmids. Uncolored lines indicate different regions among the plasmids.

bench 8.5.0 (CLCbio, Aarhus, Denmark), which generated a single contig that represented the entire circular pBH6 plasmid. This sequence includes a single BamHI site. Gene prediction was carried out with the Prokka pipeline (14). Data files were compiled using Sequin (see <http://www.ncbi.nlm.nih.gov/Sequin/>), and the SnapGene viewer was used for visualization and analysis. Pairwise alignment was performed by BLASTN and BLASTP homology searches (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) (pBH6 GenBank accession number [LGVH01000782.1](https://www.ncbi.nlm.nih.gov/nuccore/LGVH01000782.1)).

There were no chromosomal *bla*_{KPC-2} copies. Sequencing revealed a circular genetic element of 3,652 bp and included the Tn3 transposon resolvase (15), a Tn3 inverted repeat (IR), a carbapenemase-encoding gene (*bla*_{KPC-2}), a partial transposase of *ISKpn6*, and the right end of *ISKpn6*, IRR (Fig. 1). Positions 3 to 3,027 displayed homology with plasmids pKP13d (82% query coverage,

99% identity [GenBank accession number [CP003997](https://www.ncbi.nlm.nih.gov/nuccore/CP003997)]) from a KPC-2-producing *Klebsiella pneumoniae* Kp13 isolate from Brazil (16) and pFOS18 (89% query coverage, 99% identity [[KJ653815](https://www.ncbi.nlm.nih.gov/nuccore/KJ653815)]) from a KPC-2-producing *K. pneumoniae* isolate from China (17) (Fig. 2). Furthermore, *P. aeruginosa* ST1006 plasmids pCol-1 (56% query coverage, 100% identity [[KC609323](https://www.ncbi.nlm.nih.gov/nuccore/KC609323)]) from *P. aeruginosa* ST308 and pPA-2 (58% query coverage, 99% identity [[KC609322](https://www.ncbi.nlm.nih.gov/nuccore/KC609322)]) also show homology covering the *bla*_{KPC-2} gene and the partial *ISKpn6* copy.

Further analysis found that the region between ~3200 and 3600 (400 bp) of pBH6 (which includes the DNA outside the known genes and transposon features and may contain the replication origin) is similar to that of several KPC-2-producing plasmids deposited in GenBank. Interestingly, almost all related plasmids bear a *rep* gene upstream of this region and, as observed in

TABLE 1 Plasmids related to pBH6 that show similar *ori* regions (more than 97% identity)

Plasmid	Size (bp)	Coordinates	% identity	Genomic context		
				Upstream to <i>ori</i>	Downstream to <i>ori</i>	GenBank accession No.
<i>A. hydrophila</i> pKPC2	44,451	14,647–14,978	100	<i>repA</i>	Tn3 <i>tnpR</i>	KR014106
<i>E. cloacae</i> WCHECI-14653 pKPC2-EC14653	88,214	10,429–10,612	100	<i>repA</i>	Hypothetical	KP868646
<i>K. pneumoniae</i> 565 pKPCAPSS	127,970	62,229–62,565	99	<i>repA</i>	Tn <i>Swi1 tnpR</i>	KP008371
<i>A. hydrophila</i> AH1 pN6	11,467	437–106	99	<i>repA</i>	Tn <i>Swi1 tnpA</i>	KC355363
<i>Enterobacter cloacae</i> ECN49 pKPC-ECN49	41,317	40,271–40,469	99	<i>repB</i>	Tn3 <i>tnpR</i>	KP726894
<i>Citrobacter freundii</i> p112298-KPC	117,288	10,979–11,177	98	<i>repA</i>	IS26 <i>tnpA</i>	KP987215
<i>K. pneumoniae</i> pFOS18	23,939	5,108–4,910	98	<i>repA</i>	Tn <i>Shes11 tnpR</i>	KJ653815
<i>Enterobacter aerogenes</i> pHS112625	18,184	7,459–7,261	98	<i>repA</i>	Tn3 <i>tnpR</i>	KJ210592
<i>E. coli</i> pHS102707	69,453	14,234–14,036	98	<i>repA</i>	Tn3 <i>tnpR</i>	KF701335
<i>K. pneumoniae</i> pHS092839	15,499	4,142–3,944	98	<i>repA</i>	Tn3 <i>tnpR</i>	KF724506
<i>K. pneumoniae</i> pCT-KPC LJ04	151,466	81,583–81,385	97	<i>repA</i>	Tn3 <i>tnpR</i>	KT185451
<i>K. pneumoniae</i> p628-KPC	105,008	15,073–14,875	97	<i>repA</i>	Tn3 <i>tnpR</i>	KP987218
<i>K. pneumoniae</i> pKP1034	136,848	121,075–120,877	97	<i>repA</i>	Tn3 <i>tnpR</i>	KP893385
<i>E. coli</i> pHS10842	13,915	4,585–4,387	97	<i>repA</i>	Tn3 <i>tnpR</i>	KP125892
<i>K. pneumoniae</i> pHS092753	12,966	4,167–3,969	97	<i>repA</i>	Tn3 <i>tnpR</i>	KF826293
<i>K. pneumoniae</i> pHS10505	15,464	4,716–4,518	97	<i>repA</i>	Tn3 <i>tnpR</i>	KF826292
<i>E. coli</i> pHS10842	14,757	4,330–4,132	97	<i>repA</i>	Tn3 <i>tnpR</i>	KF826291
<i>K. pneumoniae</i> pHS082416	12,915	4,116–3,918	97	<i>repA</i>	Tn3 <i>tnpR</i>	KF724507
<i>K. pneumoniae</i> pHS062105	42,848	32,964–32,766	97	<i>repA</i>	Tn3 <i>tnpR</i>	KF623109
<i>K. pneumoniae</i> JM45 p1	317,154	120,918–121,116	97	<i>repA</i>	Hypothetical	CP006657
<i>K. pneumoniae</i> pKPC-LK30	86,518	61,401–61,203	97	Tn3 <i>tnpR</i>	Tn3 <i>tnpA</i>	KC405622
<i>K. pneumoniae</i> pKPHS2	111,195	25,304–25,502	97	<i>repA</i>	Tn3 <i>tnpR</i>	CP003224
<i>K. pneumoniae</i> pK048	151,188	15,712–15,514	97	<i>repA</i>	Tn3 <i>tnpR</i>	FJ628167

pBH6, a recombinase of the Tn3 family element downstream. A list of related plasmids is shown in Table 1. Only *Aeromonas hydrophila* pKPC2, *K. pneumoniae* 565 pKPCAPSS, and *Aeromonas hydrophila* AH1-pN6 show high similarity to each other over the entire 331-bp region (82% query coverage, 100% identity). The other plasmids, while maintaining high levels of similarity, only span between 100 and 300 bp, which suggests that this region may be related to plasmid replication and recognized as an origin of replication (*ori*).

It is known that small plasmids, such as pUC19, do not require a plasmid-specific *rep* gene but can replicate from a plasmid origin using bacterial host enzymes. The genetic environment of *bla*_{KPC-2} here is different from that of those already described. In most *Enterobacteriaceae* and *Pseudomonas* spp., *bla*_{KPC-2} appears associated with Tn4401-like transposons (1) carried by different plasmids (13).

To verify whether pBH6 could be established and replicate in other bacteria, purified plasmid DNA was transformed (18, 19) into *Escherichia coli* DH10B and *P. aeruginosa* PAO recipient strains followed by selection on MacConkey agar supplemented with ceftazidime (8 µg/ml). Transformants were easily obtained in both recipient strains. To evaluate the stability of pBH6, transformants from each strain isolated on LB solid medium were subcultured for 10 consecutive days in Mueller-Hinton (MH) and LB liquid media with or without 8 µg/ml ceftazidime. After the tenth subculture, both transformants still carried *bla*_{KPC-2} as judged by PCR, suggesting that this small plasmid can replicate and remain stable for several subcultures.

In summary, we have identified and obtained the sequence of a small 3,652-bp plasmid, pBH6, from KPC-producing *P. aeruginosa* ST244. Plasmid pBH6 carries the *bla*_{KPC-2} gene and remnants of a Tn3 family transposon and of *ISKpn6* and may have been derived from the Tn4401 transposon in which *ISKpn6* and *bla*_{KPC} are juxtaposed (20). When known sequences are taken into account, only 675 bp remain for providing replication functions, which implies that this region carries the origin of plasmid replication. Plasmid pBH6 does not include an identifiable *rep* gene and can replicate autonomously in *E. coli* and *P. aeruginosa*.

ACKNOWLEDGMENTS

We thank Ilana B. C. Camargo and Hyllo B. Marcelo, Jr., for providing the isolate studied. We also thank Joseane Cristina Ferreira, Ludmilla Tonani Carvalho, and Rubens Eduardo da Silva for their support in some experiments. We are grateful to Alessandra Carattoli for her precious advice on the plasmid experiments.

Research grant 2014/14494-8 from the São Paulo Research Foundation (FAPESP) is appreciated. Renata Galetti was supported by a postdoctoral fellowship from FAPESP (grant 2015/11728-0). Leonardo Neves Andrade was supported by a postdoctoral fellowship (CAPES/PNPD). There are no conflicts of interest to declare.

FUNDING INFORMATION

This work was funded by Sao Paulo Research Foundation (FAPESP) (2014/14494-8).

REFERENCES

- Andrade LN, Curiao T, Ferreira JC, Longo JM, Climaco EC, Martinez R, Bellissimo-Rodrigues F, Basile-Filho A, Evaristo MA, Del Peloso PF, Ribeiro VB, Barth AL, Paula MC, Baquero F, Canton R, Darini AL, Coque TM. 2011. Dissemination of *bla*_{KPC-2} by the spread of *Klebsiella pneumoniae* clonal complex 258 clones (ST258, ST11, ST437) and plasmids (IncFII, IncN, IncL/M) among *Enterobacteriaceae* species in Brazil.

- Antimicrob Agents Chemother 55:3579–3583. <http://dx.doi.org/10.1128/AAC.01783-10>.
- Munoz-Price LS, Poirel L, Bonomo RA, Schwaber MJ, Daikos GL, Cormican M, Cornaglia G, Garau J, Gniadkowski M, Hayden MK, Kumarasamy K, Livermore DM, Maya JJ, Nordmann P, Patel JB, Paterson DL, Pitout J, Villegas MV, Wang H, Woodford N, Quinn JP. 2013. Clinical epidemiology of the global expansion of *Klebsiella pneumoniae* carbapenemases. *Lancet Infect Dis* 13:785–796. [http://dx.doi.org/10.1016/S1473-3099\(13\)70190-7](http://dx.doi.org/10.1016/S1473-3099(13)70190-7).
- Pereira PS, de Araujo CF, Seki LM, Zahner V, Carvalho-Assef AP, Asensi MD. 2013. Update of the molecular epidemiology of KPC-2-producing *Klebsiella pneumoniae* in Brazil: spread of clonal complex 11 (ST11, ST437 and ST340). *J Antimicrob Chemother* 68:312–316. <http://dx.doi.org/10.1093/jac/dks396>.
- Villegas MV, Lolans K, Correa A, Kattan JN, Lopez JA, Quinn JP, Colombian Nosocomial Resistance Study Group. 2007. First identification of *Pseudomonas aeruginosa* isolates producing a KPC-type carbapenem-hydrolyzing beta-lactamase. *Antimicrob Agents Chemother* 51:1553–1555. <http://dx.doi.org/10.1128/AAC.01405-06>.
- Jacome PR, Alves LR, Cabral AB, Lopes AC, Maciel MA. 2012. First report of KPC-producing *Pseudomonas aeruginosa* in Brazil. *Antimicrob Agents Chemother* 56:4990. <http://dx.doi.org/10.1128/AAC.00699-12>.
- Rizek C, Fu L, Dos Santos LC, Leite G, Ramos J, Rossi F, Guimaraes T, Levin AS, Costa SF. 2014. Characterization of carbapenem-resistant *Pseudomonas aeruginosa* clinical isolates, carrying multiple genes coding for this antibiotic resistance. *Ann Clin Microbiol Antimicrob* 13:43. <http://dx.doi.org/10.1186/s12941-014-0043-3>.
- Naas T, Bonnin RA, Cuzon G, Villegas MV, Nordmann P. 2013. Complete sequence of two KPC-harboring plasmids from *Pseudomonas aeruginosa*. *J Antimicrob Chemother* 68:1757–1762. <http://dx.doi.org/10.1093/jac/dkt094>.
- Curran B, Jonas D, Grundmann H, Pitt T, Dowson CG. 2004. Development of a multilocus sequence typing scheme for the opportunistic pathogen *Pseudomonas aeruginosa*. *J Clin Microbiol* 42:5644–5649. <http://dx.doi.org/10.1128/JCM.42.12.5644-5649.2004>.
- Clinical and Laboratory Standards Institute. 2013. Performance standards for antimicrobial susceptibility testing; 23rd informational supplement. CLSI M100-S23. Clinical and Laboratory Standards Institute, Wayne, PA.
- Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, Harbarth S, Hindler JF, Kahlmeter G, Olsson-Liljequist B, Paterson DL, Rice LB, Stelling J, Struelens MJ, Vatopoulos A, Weber JT, Monnet DL. 2012. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 18:268–281. <http://dx.doi.org/10.1111/j.1469-0691.2011.03570.x>.
- Curiao T, Morosini MI, Ruiz-Garbajosa P, Robustillo A, Baquero F, Coque TM, Canton R. 2010. Emergence of *bla*_{KPC-3}-Tn4401a associated with a pKPN3/4-like plasmid within ST384 and ST388 *Klebsiella pneumoniae* clones in Spain. *J Antimicrob Chemother* 65:1608–1614. <http://dx.doi.org/10.1093/jac/dkq174>.
- Ferreira JC, Penha Filho RA, Andrade LN, Berchieri A Jr, Darini AL. 2014. Detection of chromosomal *bla* in diverse *Escherichia coli* isolates from healthy broiler chickens. *Clin Microbiol Infect* 20:O623–O626. <http://dx.doi.org/10.1111/1469-0691.12531>.
- Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. 2005. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 63:219–228. <http://dx.doi.org/10.1016/j.mimet.2005.03.018>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <http://dx.doi.org/10.1093/bioinformatics/btu153>.
- Partridge SR. 2015. What's in a name? *ISSwiI* corresponds to transposons related to Tn2 and Tn3. *mBio* 6(5):e01344-15. <http://dx.doi.org/10.1128/mBio.01344-15>.
- Ramos PI, Picao RC, Almeida LG, Lima NC, Girardello R, Vivan AC, Xavier DE, Barcellos FG, Pelisson M, Vespero EC, Medigue C, Vasconcelos AT, Gales AC, Nicolas MF. 2014. Comparative analysis of the complete genome of KPC-2-producing *Klebsiella pneumoniae* Kp13 reveals remarkable genome plasticity and a wide repertoire of virulence and resistance mechanisms. *BMC Genomics* 15:54. <http://dx.doi.org/10.1186/1471-2164-15-54>.

17. Jiang Y, Shen P, Wei Z, Liu L, He F, Shi K, Wang Y, Wang H, Yu Y. 2015. Dissemination of a clone carrying a fosA3-harboring plasmid mediates high fosfomycin resistance rate of KPC-producing *Klebsiella pneumoniae* in China. *Int J Antimicrob Agents* 45:66–70. <http://dx.doi.org/10.1016/j.ijantimicag.2014.08.010>.
18. Itoh N, Kouzai T, Koide Y. 1994. Efficient transformation of *Pseudomonas* strains with pNI vectors by electroporation. *Biosci Biotechnol Biochem* 58:1306–1308. <http://dx.doi.org/10.1271/bbb.58.1306>.
19. Choi KH, Kumar A, Schweizer HP. 2006. A 10-min method for preparation of highly electrocompetent *Pseudomonas aeruginosa* cells: application for DNA fragment transfer between chromosomes and plasmid transformation. *J Microbiol Methods* 64:391–397. <http://dx.doi.org/10.1016/j.mimet.2005.06.001>.
20. Nordmann P, Dortet L, Poirel L. 2012. Carbapenem resistance in *Enterobacteriaceae*: here is the storm! *Trends Mol Med* 18:263–272. <http://dx.doi.org/10.1016/j.molmed.2012.03.003>.