



Genomic Characterization of Carbapenemase-Producing *Klebsiella pneumoniae* with Chromosomally Carried *bla*_{NDM-1}

Noriko Sakamoto,^a Yukihiro Akeda,^{a,b,c} Yo Sugawara,^a Dan Takeuchi,^a Daisuke Motooka,^d Norihisa Yamamoto,^{a,b,c} Warawut Laolerd,^e Pitak Santanirand,^e Shigeyuki Hamada^a

^aJapan-Thailand Research Collaboration Center for Infectious Diseases, Research Institute for Microbial Diseases, Osaka University, Osaka, Japan

^bDivision of Infection Control and Prevention, Osaka University Hospital, Osaka University, Osaka, Japan

^cDepartment of Infection Control and Prevention, Graduate School of Medicine, Osaka University, Osaka, Japan

^dDepartment of Infection Metagenomics, Genome Information Center, Research Institute for Microbial Diseases, Osaka University, Osaka, Japan

^eFaculty of Medicine, Ramathibodi Hospital, Mahidol University, Bangkok, Thailand

ABSTRACT We report here *Klebsiella pneumoniae* strains carrying chromosomal *bla*_{NDM-1} in Thailand. The genomes of these two isolates include a 160-kbp insertion containing *bla*_{NDM-1}, which is almost identical to that in the IncHI1B-like plasmid. Further analysis indicated that IS5-mediated intermolecular transposition and Tn3 transposase-mediated homologous recombination resulted in the integration of *bla*_{NDM-1} into the chromosome from an IncHI1B-like plasmid. The spread of this type of carbapenem-resistant *Enterobacteriaceae* may threaten public health and warrants further monitoring.

KEYWORDS IS5, *Klebsiella pneumoniae*, Tn3, beta-lactamases, *bla*_{NDM-1}, carbapenemase-producing *Enterobacteriaceae*, carbapenems

Carbapenems are the antibiotics of last resort for combating multidrug-resistant organisms. However, their efficacy is increasingly compromised by carbapenem-resistant *Enterobacteriaceae* (CRE), which produce a variety of carbapenemases, including New Delhi metallo-β-lactamase (NDM). NDM-1 was initially reported in 2009 (1), and NDM-1-producing CRE originating on the Indian subcontinent have rapidly disseminated into different parts of the world (2). Carbapenemase genes, including *bla*_{NDM-1}, are typically plasmid borne and are responsible for the spread of CRE by horizontal gene transfer. Plasmids act as the scaffolds upon which arrays of antibiotic resistance genes are assembled, generating multiple-drug-resistant phenotypes in *Enterobacteriaceae*. However, only a limited number of CRE have been found to carry carbapenemase genes on their chromosomes (3–6). Notably, only two strains of *Escherichia coli* have been reported to carry chromosomally integrated *bla*_{NDM-1} in China and India (7, 8). However, no such organisms have been found among *Klebsiella pneumoniae* strains.

We have previously used long- and short-read whole-genome sequencing (WGS) by the PacBio RSII and Illumina MiSeq systems to examine clinical isolates of *K. pneumoniae* in a molecular epidemiology study of CRE in Bangkok, Thailand (Y. Akeda, W. Laolerd, Y. Sugawara, N. Sakamoto, D. Motooka, N. Yamamoto, D. Takeuchi, R. K. Shanmugakani, I. Nishi, M. Suzuki, K. Shibayama, P. Santanirand, K. Tomono, S. Hamada, submitted for publication) (see the supplemental material). The long reads acquired from PacBio RSII were assembled using the PacBio hierarchical genome assembly process (version 3) through the SMRT portal. To further improve the quality of the assembled sequences, short-read mapping was performed using Illumina MiSeq read data from CLC Workbench version 7.5 (CLC bio). During the course of that study, we noted that two *K. pneumoniae* isolates (strains KP64 and KP67) collected from different patients did not

Received 23 July 2018 Returned for modification 30 July 2018 Accepted 4 October 2018

Accepted manuscript posted online 15 October 2018

Citation Sakamoto N, Akeda Y, Sugawara Y, Takeuchi D, Motooka D, Yamamoto N, Laolerd W, Santanirand P, Hamada S. 2018. Genomic characterization of carbapenemase-producing *Klebsiella pneumoniae* with chromosomally carried *bla*_{NDM-1}. *Antimicrob Agents Chemother* 62:e01520-18. <https://doi.org/10.1128/AAC.01520-18>.

Copyright © 2018 American Society for Microbiology. All Rights Reserved.

Address correspondence to Yukihiro Akeda, akeda@biken.osaka-u.ac.jp.




FIG 1 Comparison of chromosomal 160-kbp regions containing *bla*_{NDM-1} in strains KP64 and KP67 and *bla*_{NDM-1}-carrying plasmid pKP3301 from strain KP33. The gray-shaded area indicates the regions with high identity (>99% similarity) between the two sequences. The 160-kbp region containing *bla*_{NDM-1} is highlighted with a red box. Nucleotide letters below the sequences of the KP64 and KP67 chromosomes represent target site duplications.

carry the *bla*_{NDM-1} gene on their plasmids. To determine the location of *bla*_{NDM-1} in these strains, we performed Southern blotting using probes for *bla*_{NDM-1} and 16S rRNA genes after pulsed-field gel electrophoresis (PFGE) (see Fig. S1A and B in the supplemental material). A bacterial conjugation assay with strains KP64, KP67, and KP33 (a control strain carrying the *bla*_{NDM-1} gene on its plasmid), with the sodium azide-resistant *E. coli* strain TUM3456, was used to detect transconjugants. However, no *bla*_{NDM-1}-carrying transconjugants were obtained from KP64 and KP67, whereas KP33 produced transconjugants. These results suggest that *bla*_{NDM-1} in strains KP64 and KP67 is not carried on a plasmid, but strains KP64 and KP67 carry the *bla*_{NDM-1} gene on their chromosomes.

The genomic characteristics of strains KP64, KP67, and KP33, all of which belong to multilocus sequence type (MLST) ST14, are summarized in Table 1. The chromosomal genomic sequences for KP64 and KP67 are closely related to that of KP33, but three large insertions were observed (Fig. S2). Two insertion sequences (67-kbp and 105-kbp insertion regions) harbor conjugal transfer systems and ABC transporter genes, respectively. However, no significant similarities with plasmids, mobile elements, or target site duplications were observed. A database search with the WGS data for strains KP64 and KP67 revealed that these isolates carry *bla*_{NDM-1} on their chromosomes, in addition to diverse antimicrobial resistance genes (160-kbp insertion region). Both the KP64 and KP67 strains contain two plasmids, plasmids pKP6401 and pKP6402 and plasmids pKP6701 and pKP6702, respectively. pKP6401 and pKP6701 contain the genes associated with plasmid partitioning and type I restriction-modification systems but no antimicrobial resistance genes. They share more than 99% sequence identity with the 104-kbp plasmid pPKPN2 from NDM-1-producing *K. pneumoniae* isolate PittNDM01 (9). pKP6402 and pKP6702 each contain five antimicrobial resistance genes directed against β -lactams and aminoglycosides. Approximately 45% of the pPKPN3 sequence shares \sim 99% identity with pKP6402 and pKP6702. pKP6402 and pKP6702 lack several antimicrobial resistance genes directed against β -lactams, aminoglycosides, sulfonamide, and rifampin that are carried by pPKPN3. An antimicrobial susceptibility test showed that strains KP64 and KP67 are multidrug resistant, including to carbapenems.

WGS revealed that the chromosomes of strains KP64 and KP67 include a 160-kbp region that contains *bla*_{NDM-1} and seven other antimicrobial resistance genes (Fig. 1 and

TABLE 1 Genomic features of clinical isolates of *K. pneumoniae* containing the *bla*_{NDM-1} gene

Isolate	Genomic structure	Length (bp)	GC content (%)	No. of CDs ^a	Resistant genes	Replicon type	DDBJ accession no.
KP64	Chromosome	5,587,050	57.06	5,405	<i>bla</i> _{NDM-1} , <i>aadA2</i> , <i>armA</i> , <i>aph(3')-VIa</i> , <i>msr(E)</i> , <i>mph(E)</i> , <i>sul1</i> , <i>dfrA12</i> , <i>bla</i> _{SHV-28b}		AP018750
	Plasmid pKP6401	102,903	50.99	137	<i>bla</i> _{CTX-M-15s} , <i>bla</i> _{OXA-1} , <i>cat</i> , <i>aac(6')Ib-cr</i> , <i>fosA</i> , <i>oqxAB</i> , <i>dfrA1</i>	FIB	AP018751
	Plasmid pKP6402	31,835	52.66	49	Not detected <i>bla</i> _{CTX-M-15s} , <i>bla</i> _{OXA-1} , <i>bla</i> _{TEM-1A'} , <i>aac(6')Ib</i> , <i>aadA2</i>	R	AP018752
KP67	Chromosome	5,587,048	57.06	5,400	<i>bla</i> _{NDM-1} , <i>aadA2</i> , <i>armA</i> , <i>aph(3')-VIa</i> , <i>msr(E)</i> , <i>mph(E)</i> , <i>sul1</i> , <i>dfrA12</i> , <i>bla</i> _{SHV-28b}		AP018753
	Plasmid pKP6701	102,917	50.99	131	<i>bla</i> _{CTX-M-15s} , <i>bla</i> _{OXA-1} , <i>cat</i> , <i>aac(6')Ib-cr</i> , <i>fosA</i> , <i>oqxAB</i> , <i>dfrA1</i>	FIB	AP018754
	Plasmid pKP6702	31,833	52.65	48	Not detected <i>bla</i> _{CTX-M-15s} , <i>bla</i> _{OXA-1} , <i>bla</i> _{TEM-1A'} , <i>aac(6')Ib</i> , <i>aadA2</i>	R	AP018755
KP33	Chromosome	5,256,831	57.39	5,137	<i>bla</i> _{SHV-28b} , <i>bla</i> _{CTX-M-15s} , <i>bla</i> _{OXA-1} , <i>fosA</i> , <i>oqxAB</i> , <i>cat</i> , <i>aac(6')Ib-cr</i> , <i>dfrA1</i>		AP018747
	Plasmid pKP3301	296,750	46.54	404	<i>bla</i> _{NDM-1} , <i>aadA2</i> , <i>armA</i> , <i>aph(3')-VIa</i> , <i>msr(E)</i> , <i>mph(E)</i> , <i>sul1</i> , <i>dfrA12</i> , <i>bla</i> _{CTX-M-15s} , <i>aac(6')Ib</i> , <i>aac(6')Ib-cr</i> , <i>cat</i> , <i>dfrA1</i> , <i>dfrA14</i> , <i>qnrB1</i> , <i>tet(D)</i>	H11B-like	AP018748
	Plasmid pKP3302	102,910	50.98	132	Not detected	FIB	AP018749

^aCDs, coding DNA sequences.




FIG 2 Hypothetical mechanism of integration of the 160-kbp region of plasmid pKP3301 into the KP64 and KP67 chromosomes. (A) Schematic view of the chromosome of strain KP33 carrying an IS5 copy at the putative insertion site in the 160-kbp region. (B) Plasmid pKP3301 containing *bla*_{NDM-1}, *orf9*, and a 4-bp target site (AATG) in the middle of the *orf9* gene. (C) IS5 on the chromosome of strain KP33 recognizes the 4-bp target site in *orf9*, intermolecular transposition generates a cointegrate molecule, and the IncHI1B-like plasmid moiety is truncated (dashed line) by Tn3 transposase-mediated homologous recombination. (D) Generation of the 160-kbp region bracketed by IS5 copies. Purple boxes represent homologous regions among different strains.

Fig. S2). This region shares >99% sequence identity with the 297-kbp IncHI1B-like *bla*_{NDM-1}-carrying plasmid pKP3301 from strain KP33, which belongs to ST14 (Table 1 and Fig. S3). pKP3301 is closely related to the sequences of the plasmids from the NDM-1-producing *K. pneumoniae* isolates PittNDM01 and KP617, which originated in India and South Korea, respectively (9, 10), suggesting that ST14 carrying an IncHI1B-like plasmid is the putative ancestor of strains KP64 and KP67.

Further WGS analysis revealed that IS5 copies were located at both ends of this 160-kbp region and that these two copies share >99% similarity with the IS*Kpn26* family of transposases. We also found duplicated TTAG target sites for IS5 and truncated *orf9* genes adjacent to the IS5 copies (Fig. 1). Based on this information, we hypothesized that the following mechanism is responsible for the insertion of the genomic locus containing *bla*_{NDM-1} into isolates KP64 and KP67 from a plasmid carrying *bla*_{NDM-1} (Fig. 2). An IS5 copy is also situated at the insertion site of the 160-kbp region on the chromosome of strain KP33 (Fig. 2A), whereas pKP3301 carries the 160-kbp region bracketed by Tn3 transposase copies (Fig. 2B). It is noteworthy that both the KP64 and KP67 strains carry a copy of the Tn3 transposase in the middle of the 160-kbp insertion region. IS5 on the chromosome may then have recognized the 4-bp target site in the *orf9* gene in pKP3301, causing the intermolecular transposition that generated the cointegrand (11) (Fig. 2A to C), followed by homologous recombination between the two Tn3 transposase copies (Fig. 2C and D).

There have been several reports of *Enterobacteriaceae* carrying *bla*_{NDM-1} on their chromosomes, although only two cases have been reported for *E. coli*, in China (7) and India (8), but none have been reported for *K. pneumoniae*. A WGS analysis of Chinese *E. coli* strain Y5 showed that *bla*_{NDM-1} was embedded in its chromosome, and an IS*CR1* element probably mediated the transposition of *bla*_{NDM-1} from a plasmid possibly derived from *Proteus mirabilis* PM58. Therefore, the integration

mechanism differs completely from that involved in the two strains analyzed in this study. Poirel et al. previously suggested the chromosomal location of *bla*_{NDM-1} in *E. coli*, based on the results of a conjugation assay. However, the integration mechanism and genetic context surrounding this chromosomal *bla*_{NDM-1} gene are unclear (8). The *bla*_{OXA-48}-type and *bla*_{KPC} genes are carried on certain transposons, such as Tn1999 and Tn4401, and are sometimes integrated into chromosomes by these transposons; however, *bla*_{NDM} was originally carried on transposon Tn125 in an *Acinetobacter* species. The Tn125 structure and its flanking sequences in the *Enterobacteriaceae* are frequently inserted by various mobile elements, generating diverse genetic contexts for *bla*_{NDM} (12, 13). The plasticity of the genetic contexts for *bla*_{NDM} indicates that this gene can be transferred by various mechanisms from plasmids to chromosomes, making tracing the origins of chromosomal *bla*_{NDM} genes very complicated.

In conclusion, this is the first report of *K. pneumoniae* strains carrying the *bla*_{NDM-1} gene on their chromosomes. We attribute the genetic transmission of *bla*_{NDM-1} from the IncHI1B-like plasmid to the chromosomes of these strains to IS5 and the Tn3 transposase. The *bla*_{NDM} genes are potentially transferable between plasmid and chromosome via multiple mobile elements; hence, monitoring the chromosomal *bla*_{NDM} genes should be epidemiologically important.

Accession number(s). The WGS data are available from the DDBJ (DNA Data Bank of Japan) database under accession numbers AP018747, AP018748, AP018749, AP018750, AP018751, AP018752, AP018753, AP018754, and AP018755.

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/AAC.01520-18>.

SUPPLEMENTAL FILE 1, PDF file, 0.9 MB.

ACKNOWLEDGMENTS

This work was supported by the Japan Initiative for Global Research Network on Infectious Diseases (J-GRID) of the Ministry of Education, Culture, Sport, Science, and Technology of Japan and the Japan Agency for Medical Research and Development (AMED) under grant number JP18fm0108003.

We are also grateful to Yoshikazu Ishii of Toho University Graduate School of Medicine for providing *E. coli* TUM3456. We thank Janine Miller from the Edanz Group for editing a draft of the manuscript.

We declare no conflict of interest.

REFERENCES

1. Yong D, Toleman MA, Giske CG, Cho HS, Sundman K, Lee K, Walsh TR. 2009. Characterization of a new metallo-beta-lactamase gene, *bla*_{NDM-1}, and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimicrob Agents Chemother* 53:5046–5054. <https://doi.org/10.1128/AAC.00774-09>.
2. Lee CR, Lee JH, Park KS, Kim YB, Jeong BC, Lee SH. 2016. Global dissemination of carbapenemase-producing *Klebsiella pneumoniae*: epidemiology, genetic context, treatment options, and detection methods. *Front Microbiol* 7:895. <https://doi.org/10.3389/fmicb.2016.00895>.
3. Beyrouthy R, Robin F, Delmas J, Gibold L, Dalmasso G, Dabboussi F, Hamze M, Bonnet R. 2014. IS1R-mediated plasticity of IncL/M plasmids leads to the insertion of *bla*_{OXA-48} into the *Escherichia coli* chromosome. *Antimicrob Agents Chemother* 58:3785–3790. <https://doi.org/10.1128/AAC.02669-14>.
4. Huang W, Wang G, Sebra R, Zhuge J, Yin C, Aguero-Rosenfeld ME, Schuetz AN, Dimitrova N, Fallon JT. 2017. Emergence and evolution of multidrug-resistant *Klebsiella pneumoniae* with both *bla*_{KPC} and *bla*_{CTX-M} integrated in the chromosome. *Antimicrob Agents Chemother* 61:e00076-17. <https://doi.org/10.1128/AAC.00076-17>.
5. Mathers AJ, Stoesser N, Chai W, Carroll J, Barry K, Cherunvanky A, Sebra R, Kasarskis A, Peto TE, Walker AS, Sifri CD, Crook DW, Sheppard AE. 2017. Chromosomal integration of the *Klebsiella pneumoniae* carbapenemase gene, *bla*_{KPC}, in *Klebsiella* species is elusive but not rare. *Antimicrob Agents Chemother* 61:e01823-16. <https://doi.org/10.1128/AAC.01823-16>.
6. Turton JF, Doumith M, Hopkins KL, Perry C, Meunier D, Woodford N. 2016. Clonal expansion of *Escherichia coli* ST38 carrying a chromosomally integrated OXA-48 carbapenemase gene. *J Med Microbiol* 65:538–546. <https://doi.org/10.1099/jmm.0.000248>.
7. Shen P, Yi M, Fu Y, Ruan Z, Du X, Yu Y, Xie X. 2017. Detection of an *Escherichia coli* sequence type 167 strain with two tandem copies of *bla*_{NDM-1} in the chromosome. *J Clin Microbiol* 55:199–205. <https://doi.org/10.1128/JCM.01581-16>.
8. Poirel L, Dortet L, Bernabeu S, Nordmann P. 2011. Genetic features of *bla*_{NDM-1}-positive *Enterobacteriaceae*. *Antimicrob Agents Chemother* 55:5403–5407. <https://doi.org/10.1128/AAC.00585-11>.
9. Doi Y, Hazen TH, Boitano M, Tsai YC, Clark TA, Korlach J, Rasko DA. 2014. Whole-genome assembly of *Klebsiella pneumoniae* coproducing NDM-1 and OXA-232 carbapenemases using single-molecule, real-time sequencing. *Antimicrob Agents Chemother* 58:5947–5953. <https://doi.org/10.1128/AAC.03180-14>.
10. Kwon T, Jung YH, Lee S, Yun MR, Kim W, Kim DW. 2016. Comparative genomic analysis of *Klebsiella pneumoniae* subsp. *pneumoniae* KP617 and PittNDM01,

- NUHL24835, and ATCC BAA-2146 reveals unique evolutionary history of this strain. *Gut Pathog* 8:34. <https://doi.org/10.1186/s13099-016-0117-1>.
11. He S, Hickman AB, Varani AM, Siguier P, Chandler M, Dekker JP, Dyda F. 2015. Insertion sequence IS26 reorganizes plasmids in clinically isolated multidrug-resistant bacteria by replicative transposition. *mBio* 6:e00762-15. <https://doi.org/10.1128/mBio.00762-15>.
 12. Wailan AM, Paterson DL, Kennedy K, Ingram PR, Bursle E, Sidjabat HE. 2016. Genomic characteristics of NDM-producing *Enterobacteriaceae* isolates in Australia and their *bla_{NDM}* genetic contexts. *Antimicrob Agents Chemother* 60:136–141. <https://doi.org/10.1128/AAC.01243-15>.
 13. Partridge SR, Iredell JR. 2012. Genetic contexts of blaNDM-1. *Antimicrob Agents Chemother* 56:6065–6067. <https://doi.org/10.1128/AAC.00117-12>.