



Evaluation of the *In Vitro* Activity of Eravacycline against a Broad Spectrum of Recent Clinical Anaerobic Isolates

David R. Snyderman,^a Laura A. McDermott,^a Nilda V. Jacobus,^a Kathryn Kerstein,^b Trudy H. Grossman,^{b*} Joyce A. Sutcliffe^{b*}

^aDepartment of Medicine, Tufts Medical Center, Boston, Massachusetts, USA

^bTetraphase Pharmaceuticals, Watertown, Massachusetts, USA

ABSTRACT The novel fluorocycline antibiotic eravacycline is in development for use in the treatment of serious infections caused by susceptible and multidrug-resistant (MDR) aerobic and anaerobic Gram-negative and Gram-positive pathogens. Eravacycline and 11 comparator antibiotics were tested against recent anaerobic clinical isolates, including MDR *Bacteroides* spp. and *Clostridium difficile*. Eravacycline was potent *in vitro* against all the isolates tested, including strains with tetracycline-specific resistance determinants and MDR anaerobic pathogens resistant to carbapenems and/or β -lactam- β -lactamase inhibitor combinations.

KEYWORDS eravacycline, anaerobes, *Bacteroides*, antimicrobial resistance

Increased resistance of anaerobic bacteria to standard antibiotics requires the development of new antibiotics for use in mixed aerobic-anaerobic organism infections (1–3). Eravacycline is a novel fluorocycline antibiotic in phase 3 clinical development for the treatment of serious infections due to Gram-negative and Gram-positive aerobic and anaerobic pathogens (4, 5). Eravacycline retains activity against commonly described tetracycline resistance mechanisms, such as tetracycline efflux pumps and ribosomal protection (6, 7), and is active *in vitro* against Gram-negative aerobic pathogens resistant to other classes of antibiotics, including *Acinetobacter baumannii* and *Enterobacteriaceae* expressing extended-spectrum β -lactamases, carbapenemases, and colistin/polymyxin resistance (8–14). The spectrum of eravacycline also includes potency against Gram-positive pathogens, such as methicillin-susceptible and -resistant staphylococci, vancomycin-susceptible and -resistant enterococci, and penicillin-susceptible and -resistant *Streptococcus pneumoniae* (10, 15). Eravacycline, however, has reduced activity against *Pseudomonas aeruginosa* and *Burkholderia cenocepacia* (10).

(These data were presented at the 55th Interscience Conference on Antimicrobial Agents and Chemotherapy, poster C-547, 17 to 21 September 2015, San Diego, CA [16].)

Earlier evaluations showed that eravacycline was potent against a wide variety of anaerobic pathogens *in vitro* (10, 17); however, the number of isolates in each species was limited and not as representative of antibiotic resistance as is currently seen. To expand and define the *in vitro* spectrum of eravacycline against anaerobic pathogens, particularly *Bacteroides fragilis* and *Clostridium difficile*, eravacycline and comparator antibiotics (tigecycline, minocycline, imipenem, meropenem, piperacillin-tazobactam, ampicillin-sulbactam, moxifloxacin, metronidazole, linezolid, clindamycin, and vancomycin [with Gram-positive isolates only]) were tested against 540 recent anaerobic clinical isolates, including MDR isolates, collected in the United States from 2012 to 2015 at Tufts Medical Center from hospitalized patient cultures and medical centers participating in *B. fragilis* and *C. difficile* surveillance studies (1, 3). Prior to testing, identification of the isolates was confirmed using API 20A methodology (bioMérieux, Inc., Durham, NC) and methods outlined in the Wadsworth-KTL anaerobic bacteriology manual (18), including plating on selective media and susceptibility to special potency

Received 25 October 2017 Returned for modification 13 November 2017 Accepted 18 February 2018

Accepted manuscript posted online 26 February 2018

Citation Snyderman DR, McDermott LA, Jacobus NV, Kerstein K, Grossman TH, Sutcliffe JA. 2018. Evaluation of the *in vitro* activity of eravacycline against a broad spectrum of recent clinical anaerobic isolates. Antimicrob Agents Chemother 62:e02206-17. <https://doi.org/10.1128/AAC.02206-17>.

Copyright © 2018 American Society for Microbiology. All Rights Reserved.

Address correspondence to David R. Snyderman, dsnyderman@tuftsmedicalcenter.org.

* Present address: Trudy H. Grossman, Melinta Therapeutics, New Haven, Connecticut, USA; Joyce A. Sutcliffe, Consultant, Westbrook, Connecticut, USA.

antimicrobial disks. MIC assays were performed by an agar dilution method according to Clinical and Laboratory Standards Institute (CLSI) recommendations (19) using American Type Culture Collection strains *Bacteroides fragilis* ATCC 25285, *Bacteroides thetaiotaomicron* ATCC 29741, *Eggerthella lenta* ATCC 43055, and *Clostridium difficile* ATCC 700057 as controls. Percent resistance was calculated using applicable CLSI breakpoints (20), except for tigecycline, for which FDA breakpoints (21) were used, or EUCAST cutoff values (ECOFF) (22), which were used for *C. difficile*.

The activities of eravacycline and comparators against Gram-negative anaerobic isolates are illustrated in Table 1. Versus comparator antibiotics, eravacycline showed the lowest MIC values against isolates of the *B. fragilis* group, including those resistant to tigecycline, minocycline, meropenem, piperacillin-tazobactam, ampicillin-sulbactam, moxifloxacin, and clindamycin. All isolates within the *B. fragilis* group ($n = 286$) were inhibited by ≤ 4 $\mu\text{g/ml}$ eravacycline. The MIC values that inhibited 90% of the isolates (MIC_{90}) in a panel for eravacycline (MIC_{90} , 1 $\mu\text{g/ml}$) were 8- and 16-fold lower than those of tigecycline (MIC_{90} , 8 $\mu\text{g/ml}$) and minocycline (MIC_{90} , 16 $\mu\text{g/ml}$), respectively. Eravacycline was potent against *Prevotella* spp. ($n = 29$) and *Fusobacterium* spp. ($n = 20$); all isolates were inhibited by ≤ 0.5 $\mu\text{g/ml}$ eravacycline.

The susceptibilities of the Gram-positive anaerobic isolates to eravacycline and comparator antibiotics are shown in Table 2. Eravacycline was potent against all isolates at concentrations of ≤ 1 $\mu\text{g/ml}$, including *C. difficile* strains resistant to moxifloxacin and clindamycin and with elevated MICs (4 $\mu\text{g/ml}$) to vancomycin and metronidazole. The eravacycline MIC_{90} values against *Clostridium perfringens* ($n = 15$), *C. difficile* ($n = 76$), other *Clostridium* spp. ($n = 22$), *Peptostreptococcus* spp. ($n = 53$), *Propionibacterium* species (including *P. acnes*, which was recently reclassified as *Cutibacterium acnes* spp.) ($n = 13$), and *Bifidobacterium* spp. ($n = 15$) were 1 $\mu\text{g/ml}$, 0.12 $\mu\text{g/ml}$, 0.12 $\mu\text{g/ml}$, 0.25 $\mu\text{g/ml}$, 0.25 $\mu\text{g/ml}$, and 0.5 $\mu\text{g/ml}$, respectively. The MIC ranges for *E. lenta* ($n = 6$) and *Lactobacillus* spp. ($n = 5$) were 0.03 to 0.12 $\mu\text{g/ml}$ and 0.06 to 0.5 $\mu\text{g/ml}$, respectively.

The presence of the following tetracycline resistance genes previously reported in *B. fragilis* (<http://faculty.washington.edu/marilynr/>) was detected by standard PCR methodology for a set of 27 *B. fragilis* isolates covering the full range of minocycline MIC values (≤ 0.25 to 32 $\mu\text{g/ml}$): *tet(Q)* and *tet(M)*, encoding ribosomal protection mechanisms (23); and *tet(X)*, *tet(X1)*, and *tet(X2)*, encoding tetracycline-modifying flavin-dependent monooxygenases (24). The following primer sets were used in PCRs: for *tet(Q)*, forward 5'-GTGCGTTTCGACAATGCATCTATTGTAG and reverse 5'-TGATGACATTGATTTTGGAAACATG primers (derived from GenBank accession no. Z21523) or forward 5'-ATCGGTATCAATGAGTTGTT and reverse 5'-GACTGATTCTGGAGGAAGTA primers (25); for *tet(X)*, *tet(X1)*, and *tet(X2)*, forward 5'-CAGGAAGCAATGAAAAAGCGG and reverse 5'-TAGCTTTTCTAAAGGAAATATCCG primers (derived from GenBank accession no. M37699); for *tet(X)* and *tet(X2)* only, forward 5'-TTAGCCTTACCAATGGGTGT and reverse 5'-CAAATCTGCTGTTTCACTCG primers (25); for *tet(X1)* only, forward 5'-TCAGGACAAGAAGCAATGAA and reverse 5'-TATTTCGGGGTTGTCAAAC primers (25); and for *tet(M)*, forward 5'-AACTCGAACAAGAGGAAAGC and reverse 5'-ATGGAAGCCCAGAAA GGAT primers (26). Plasmids carrying the *tet(X)*, *tet(Q)*, or *tet(M)* gene were used as positive PCR control templates, and sequencing reactions were performed by Genewiz (Cambridge, MA) to verify *tet(X)* alleles as well as to determine the *tet(Q)* sequences of two isolates with the lowest minocycline MIC values (≤ 2 $\mu\text{g/ml}$).

As shown in Table 3, 23 of 27 isolates were positive for *tet(Q)*, with minocycline, tigecycline, and eravacycline MIC values ranging from 0.5 to 32 $\mu\text{g/ml}$, 0.25 to 16 $\mu\text{g/ml}$, and 0.06 to 4 $\mu\text{g/ml}$, respectively, while the 4 isolates negative for *tet(Q)* had minocycline, tigecycline, and eravacycline MIC values ranging from ≤ 0.25 $\mu\text{g/ml}$, 0.5 to 2 $\mu\text{g/ml}$, and 0.12 to 0.5 $\mu\text{g/ml}$, respectively. The *tet(Q)* genes from two positive isolates with relatively lower minocycline MIC values of 0.5 and 2 $\mu\text{g/ml}$ were sequenced; each encoded amino acid sequences with 100% identity to 74 Tet(Q) proteins in the UniProtKB database (<http://www.uniprot.org/help/uniprotkb>, released 15 March 2017), indicating that the Tet(Q) protein in these two isolates was a common variant. In addition to being positive for *tet(Q)*, two isolates were positive for *tet(X)*, and three

TABLE 1 In vitro activities of eravacycline and comparator antibiotics against 335 Gram-negative anaerobic clinical isolates

Organism (n)	Antimicrobial agent	MIC (µg/ml)			% resistant by guideline ^a	
		Range	MIC ₅₀	MIC ₉₀	CLSI/FDA	EUCAST
<i>Bacteroides fragilis</i> group (286)	Eravacycline	0.03 to 4	0.25	1	NA	NA
	Tigecycline	0.12 to 16	1	8	3.1	NA
	Minocycline	≤0.25 to 64	8	16	NA	NA
	Imipenem	≤0.12 to 4	0.25	1	0	0
	Meropenem	≤0.12 to 16	0.25	2	1.0	1.0
	Piperacillin-tazobactam	≤0.5 to 256	4	32	0.7	10.8
	Ampicillin-sulbactam	≤0.5 to >128	2	16	4.9	22.7
	Moxifloxacin	≤0.5 to 64	2	32	38.8	NA
	Linezolid	≤1 to >16	2	4	NA	NA
	Clindamycin	≤0.5 to >128	4	>128	43.7	43.7
	Metronidazole	≤1 to 2	≤1	2	0	0
<i>Bacteroides fragilis</i> (110)	Eravacycline	0.03 to 4	0.25	1	NA	NA
	Tigecycline	0.12 to 16	1	8	4.5	NA
	Minocycline	≤0.25 to 64	8	16	NA	NA
	Imipenem	≤0.12 to 4	0.25	1	0	0
	Meropenem	≤0.12 to 16	0.25	2	1.8	1.8
	Piperacillin-tazobactam	≤0.5 to 128	1	8	0.9	1.8
	Ampicillin-sulbactam	1 to 32	2	16	1.8	17.3
	Moxifloxacin	≤0.5 to 32	2	16	36.4	NA
	Linezolid	2 to >16	2	4	NA	NA
	Clindamycin	≤0.5 to >128	1	>128	30.0	30.0
	Metronidazole	≤1 to 2	≤1	2	0	0
Non- <i>B. fragilis</i> (176)	Eravacycline	0.06 to 4	0.25	1	NA	NA
	Tigecycline	0.12 to 16	1	8	2.3	NA
	Minocycline	≤0.25 to 32	4	16	NA	NA
	Imipenem	≤0.12 to 4	0.5	1	0	0
	Meropenem	≤0.12 to 16	0.25	2	0.6	0.6
	Piperacillin-tazobactam	≤0.5 to 256	8	32	0.6	16.5
	Ampicillin-sulbactam	≤0.5 to >128	4	16	6.8	26.1
	Moxifloxacin	≤0.5 to 64	4	32	40.3	NA
	Linezolid	≤1 to >16	2	4	NA	NA
	Clindamycin	≤0.5 to >128	8	>128	52.3	52.3
	Metronidazole	≤1 to 2	≤1	2	0	0
<i>Bacteroides caccae</i> (10)	Eravacycline	0.12 to 0.5	0.5	0.5	NA	NA
	Tigecycline	0.5 to 4	1	4	0	NA
	Minocycline	4 to 16	8	16	NA	NA
	Imipenem	≤0.12 to 1	0.25	0.5	0	0
	Meropenem	≤0.12 to 4	0.25	1	0	0
	Piperacillin-tazobactam	≤0.5 to 4	2	4	0	0
	Ampicillin-sulbactam	1 to 8	2	8	0	0
	Moxifloxacin	≤0.5 to 64	4	32	50.0	NA
	Linezolid	All 2	2	2	NA	NA
	Clindamycin	4 to >128	>128	>128	90.0	90.0
	Metronidazole	≤1 to 2	≤1	2	0	0
<i>Bacteroides ovatus</i> (30)	Eravacycline	0.06 to 1	0.25	1	NA	NA
	Tigecycline	0.25 to 8	1	8	0	NA
	Minocycline	≤0.25 to 16	4	8	NA	NA
	Imipenem	≤0.12 to 2	0.25	0.5	0	0
	Meropenem	≤0.12 to 8	0.25	2	0	0
	Piperacillin-tazobactam	≤0.5 to 32	4	32	0	13.3
	Ampicillin-sulbactam	≤0.5 to 32	2	16	3.3	20.0
	Moxifloxacin	≤0.5 to 32	4	8	20.0	NA
	Linezolid	≤1 to 8	2	4	NA	NA
	Clindamycin	≤0.5 to >128	8	>128	53.3	53.3
	Metronidazole	≤1 to 2	≤1	2	0	0
<i>Bacteroides thetaiotaomicron</i> (70)	Eravacycline	0.06 to 4	0.25	1	NA	NA
	Tigecycline	0.12 to 16	1	8	4.3	NA
	Minocycline	≤0.25 to 16	4	8	NA	NA
	Imipenem	≤0.12 to 1	0.5	1	0	0
	Meropenem	≤0.12 to 2	0.25	1	0	0
	Piperacillin-tazobactam	1 to 64	16	32	0	20.0
	Ampicillin-sulbactam	1 to 64	2	16	4.3	18.6
	Moxifloxacin	≤0.5 to 64	2	32	35.7	NA
	Linezolid	2 to 4	4	4	NA	NA
	Clindamycin	≤0.5 to >128	8	>128	51.4	51.4
	Metronidazole	≤1 to 2	≤1	2	0	0

(Continued on next page)

Downloaded from <http://aac.asm.org/> on February 28, 2021 by guest

TABLE 1 (Continued)

Organism (n)	Antimicrobial agent	MIC ($\mu\text{g/ml}$)			% resistant by guideline ^a	
		Range	MIC ₅₀	MIC ₉₀	CLSI/FDA	EUCAST
<i>Parabacteroides distasonis</i> (26)	Eravacycline	0.12 to 4	0.25	1	NA	NA
	Tigecycline	0.5 to 16	1	8	3.8	NA
	Minocycline	≤ 0.25 to 32	4	16	NA	NA
	Imipenem	0.25 to 2	0.5	2	0	0
	Meropenem	≤ 0.12 to 4	0.5	4	0	0
	Piperacillin-tazobactam	1 to 64	8	32	0	26.9
	Ampicillin-sulbactam	2 to 64	8	32	15.4	38.5
	Moxifloxacin	≤ 0.5 to 64	2	32	46.2	NA
	Linezolid	2 to 4	2	4	NA	NA
	Clindamycin	≤ 0.5 to >128	4	>128	38.5	38.5
	Metronidazole	≤ 1 to 2	≤ 1	2	0	0
<i>Bacteroides uniformis</i> (15)	Eravacycline	0.06 to 1	0.25	1	NA	NA
	Tigecycline	0.25 to 8	1	4	0	NA
	Minocycline	≤ 0.25 to 32	8	16	NA	NA
	Imipenem	≤ 0.12 to 2	0.25	1	0	0
	Meropenem	≤ 0.12 to 4	0.25	2	0	0
	Piperacillin-tazobactam	≤ 0.5 to 32	4	32	0	13.3
	Ampicillin-sulbactam	≤ 0.5 to 16	2	16	0	20.0
	Moxifloxacin	≤ 0.5 to 64	8	32	53.3	NA
	Linezolid	2 to 4	2	4	NA	NA
	Clindamycin	≤ 0.5 to >128	>128	>128	60.0	60.0
	Metronidazole	≤ 1 to 2	≤ 1	2	0	0
<i>Bacteroides vulgatus</i> (18)	Eravacycline	0.06 to 0.5	0.12	0.5	NA	NA
	Tigecycline	0.12 to 8	0.5	8	0	NA
	Minocycline	≤ 0.25 to 16	8	16	NA	NA
	Imipenem	0.25 to 4	0.5	2	0	0
	Meropenem	0.25 to 16	0.5	4	5.6	5.6
	Piperacillin-tazobactam	2 to 256	4	32	5.6	11.1
	Ampicillin-sulbactam	2 to 64	16	64	16.7	61.1
	Moxifloxacin	1 to 64	32	64	72.2	NA
	Linezolid	2 to >16	2	4	NA	NA
	Clindamycin	≤ 0.5 to >128	>128	>128	55.6	55.6
	Metronidazole	≤ 1 to 2	≤ 1	≤ 1	0	0
Other <i>Parabacteroides/Bacteroides</i> spp. (7) ^b	Eravacycline	0.06 to 1				
	Tigecycline	0.5 to 8				
	Minocycline	≤ 0.25 to 32				
	Imipenem	≤ 0.12 to 1				
	Meropenem	0.25 to 1				
	Piperacillin-tazobactam	≤ 0.5 to 16				
	Ampicillin-sulbactam	2 to >128				
	Moxifloxacin	≤ 0.5 to 8				
	Linezolid	2 to 4				
	Clindamycin	≤ 0.5 to >128				
Metronidazole	≤ 1 to 2					
<i>Prevotella</i> spp. (29)	Eravacycline	≤ 0.015 to 0.5	0.25	0.5	NA	NA
	Tigecycline	≤ 0.06 to 1	0.5	1	0	NA
	Minocycline	≤ 0.25 to 32	4	32	NA	NA
	Imipenem	≤ 0.12 to 2	≤ 0.12	2	0	0
	Meropenem	≤ 0.12 to 8	≤ 0.12	2	0	0
	Piperacillin-tazobactam	≤ 0.5 to 128	≤ 0.5	8	3.4	3.4
	Ampicillin-sulbactam	≤ 0.5 to 32	1	4	3.4	3.4
	Moxifloxacin	≤ 0.5 to >64	1	16	10.3	NA
	Linezolid	≤ 1 to 4	2	4	NA	NA
	Clindamycin	≤ 0.5 to >128	≤ 0.5	>128	27.6	27.6
	Metronidazole	≤ 1 to >16	≤ 1	2	3.4	3.4
<i>Fusobacterium</i> spp. (20)	Eravacycline	≤ 0.015 to 0.5	0.03	0.12	NA	NA
	Tigecycline	≤ 0.06 to 1	0.12	0.5	0	NA
	Minocycline	≤ 0.25 to 0.5	≤ 0.25	0.5	NA	NA
	Imipenem	≤ 0.12 to 0.5	≤ 0.12	0.5	0	0
	Meropenem	≤ 0.12 to 0.5	0.25	0.25	0	0
	Piperacillin-tazobactam	≤ 0.5 to 2	1	2	0	0
	Ampicillin-sulbactam	≤ 0.5 to 2	1	2	0	0
	Moxifloxacin	≤ 0.5 to 2	≤ 0.5	1	0	NA
	Linezolid	≤ 1 to 2	≤ 1	≤ 1	NA	NA
	Clindamycin	All ≤ 0.5	≤ 0.5	≤ 0.5	0	0
	Metronidazole	All ≤ 1	≤ 1	≤ 1	0	0

^aThe FDA breakpoint applicable only to tigecycline. EUCAST epidemiological cutoff (ECOFF) breakpoints were used for *C. difficile*. NA, not applicable.

^bOnly the MIC range is indicated when the total number of isolates was <10 .

TABLE 2 *In vitro* activities of eravacycline and comparator antibiotics against 205 Gram-positive anaerobic clinical isolates

Species (n)	Antimicrobial agent	MIC ($\mu\text{g/ml}$)			% resistant by guideline ^a	
		Range	MIC ₅₀	MIC ₉₀	CLSI/FDA	EUCAST
<i>Clostridium perfringens</i> (15)	Eravacycline	0.03 to 1	0.12	1	NA	NA
	Tigecycline	0.12 to 2	0.5	2	0	NA
	Minocycline	≤ 0.25 to 8	0.5	8	NA	NA
	Imipenem	≤ 0.12 to 1	0.25	0.5	0	0
	Meropenem	≤ 0.12 to 1	≤ 0.12	0.25	0	0
	Piperacillin-tazobactam	≤ 0.5 to 1	≤ 0.5	1	0	0
	Ampicillin-sulbactam	≤ 0.5 to 2	≤ 0.5	1	0	0
	Moxifloxacin	≤ 0.5 to 4	1	4	0	NA
	Linezolid	≤ 1 to 2	2	2	NA	NA
	Clindamycin	≤ 0.5 to 128	1	64	13.3	13.3
	Metronidazole	≤ 1 to 2	≤ 1	2	0	0
	Vancomycin	0.5 to 64	1	32	NA	20.0
<i>Clostridium difficile</i> (76)	Eravacycline	≤ 0.015 to 0.25	0.06	0.12	NA	NA
	Tigecycline	≤ 0.06 to 1	0.12	0.25	0	6.6
	Minocycline	≤ 0.25 to 16	≤ 0.25	4	NA	NA
	Imipenem	≤ 0.12 to 16	4	8	6.6	NA
	Meropenem	≤ 0.12 to 8	2	4	0	NA
	Piperacillin-tazobactam	≤ 0.5 to 64	8	16	0	NA
	Ampicillin-sulbactam	≤ 0.5 to 16	2	4	0	NA
	Moxifloxacin	≤ 0.5 to 32	2	16	18.4	18.4
	Linezolid	≤ 1 to 8	2	2	NA	NA
	Clindamycin	≤ 0.5 to >128	4	128	28.9	NA
	Metronidazole	≤ 1 to 4	≤ 1	≤ 1	0	1.3
	Vancomycin	≤ 0.5 to 4	1	2	NA	2.6
Other <i>Clostridium</i> spp. (22)	Eravacycline	≤ 0.015 to 0.5	0.03	0.12	NA	NA
	Tigecycline	≤ 0.06 to 1	0.12	0.5	0	NA
	Minocycline	≤ 0.25 to 16	≤ 0.25	4	NA	NA
	Imipenem	≤ 0.12 to 4	1	4	0	0
	Meropenem	≤ 0.12 to 2	1	2	0	0
	Piperacillin-tazobactam	≤ 0.5 to 32	2	16	0	9.1
	Ampicillin-sulbactam	≤ 0.5 to 2	≤ 0.5	2	0	0
	Moxifloxacin	≤ 0.5 to >64	2	4	9.1	NA
	Linezolid	≤ 1 to 8	2	4	NA	NA
	Clindamycin	≤ 0.5 to >128	4	128	40.9	40.9
	Metronidazole	≤ 1 to >16	≤ 1	≤ 1	4.5	4.5
	Vancomycin	≤ 0.25 to 16	2	16	NA	31.8
<i>Peptostreptococcus</i> spp. (53)	Eravacycline	0.03 to 0.25	0.12	0.25	NA	NA
	Tigecycline	≤ 0.06 to 1	0.25	0.25	0	NA
	Minocycline	≤ 0.25 to 16	0.5	8	NA	NA
	Imipenem	≤ 0.12 to 0.5	≤ 0.12	≤ 0.12	0	0
	Meropenem	≤ 0.12 to 0.5	≤ 0.12	≤ 0.12	0	0
	Piperacillin-tazobactam	≤ 0.5 to 4	≤ 0.5	≤ 0.5	0	0
	Ampicillin-sulbactam	≤ 0.5 to 2	≤ 0.5	1	0	0
	Moxifloxacin	≤ 0.5 to 16	≤ 0.5	4	9.4	NA
	Linezolid	≤ 1 to 2	≤ 1	2	NA	NA
	Clindamycin	≤ 0.5 to >128	≤ 0.5	64	11.3	11.3
	Metronidazole	≤ 1 to >16	≤ 1	≤ 1	5.7	5.7
	Vancomycin	≤ 0.25 to 4	0.5	1	NA	5.7
<i>Propionibacterium</i> spp./ <i>Cutibacterium acnes</i> (13)	Eravacycline	≤ 0.015 to 1	0.06	0.25	NA	NA
	Tigecycline	0.12 to 2	0.12	0.5	0	NA
	Minocycline	≤ 0.25 to 8	0.5	2	NA	NA
	Imipenem	≤ 0.12 to 0.5	≤ 0.12	≤ 0.12	0	0
	Meropenem	≤ 0.12 to 0.25	≤ 0.12	0.25	0	0
	Piperacillin-tazobactam	≤ 0.5 to 1	≤ 0.5	≤ 0.5	0	0
	Ampicillin-sulbactam	≤ 0.5 to 4	≤ 0.5	1	0	0
	Moxifloxacin	≤ 0.5 to 4	≤ 0.5	1	0	NA
	Linezolid	≤ 1 to 2	≤ 1	≤ 1	NA	NA
	Clindamycin	≤ 0.5 to 64	1	64	23.1	23.1
	Metronidazole	≤ 1 to >16	>16	>16	92.3	92.3
	Vancomycin	0.5 to 32	1	8	NA	23.1

(Continued on next page)

Downloaded from <http://aac.asm.org/> on February 28, 2021 by guest

TABLE 2 (Continued)

Species (n)	Antimicrobial agent	MIC ($\mu\text{g/ml}$)			% resistant by guideline ^a	
		Range	MIC ₅₀	MIC ₉₀	CLSI/FDA	EUCAST
<i>Bifidobacterium</i> spp. (15)	Eravacycline	0.12 to 0.5	0.25	0.5	NA	NA
	Tigecycline	0.12 to 1	0.5	1	0	NA
	Minocycline	0.5 to 64	1	32	NA	NA
	Imipenem	≤ 0.12 to 2	0.5	2	0	0
	Meropenem	≤ 0.12 to 8	0.5	4	0	0
	Piperacillin-tazobactam	1 to 16	2	8	0	0
	Ampicillin-sulbactam	1 to 16	1	2	0	6.7
	Moxifloxacin	1 to 64	1	32	13.3	NA
	Linezolid	≤ 1 to 2	≤ 1	2	NA	NA
	Clindamycin	≤ 0.5 to >128	≤ 0.5	>128	13.3	13.3
	Metronidazole	4 to >16	>16	>16	93.3	93.3
	Vancomycin	0.5 to 32	1	4	NA	13.3
	<i>Eggerthella lenta</i> (6) ^b	Eravacycline	0.03 to 0.12			
Tigecycline		≤ 0.06 to 0.5				
Minocycline		1 to 32				
Imipenem		≤ 0.12 to 0.5				
Meropenem		≤ 0.12 to 0.5				
Piperacillin-tazobactam		≤ 0.5 to 64				
Ampicillin-sulbactam		≤ 0.5 to 8				
Moxifloxacin		≤ 0.5 to >64				
Linezolid		≤ 1 to 4				
Clindamycin		≤ 0.5 to 16				
Metronidazole		All ≤ 1				
Vancomycin	1 to 4					
<i>Lactobacillus</i> spp. (5) ^b	Eravacycline	0.06 to 0.5				
	Tigecycline	0.25 to 1				
	Minocycline	≤ 0.25 to 4				
	Imipenem	≤ 0.12 to 2				
	Meropenem	≤ 0.12 to 8				
	Piperacillin-tazobactam	≤ 0.5 to 4				
	Ampicillin-sulbactam	≤ 0.5 to 2				
	Moxifloxacin	≤ 0.5 to 2				
	Linezolid	≤ 1 to 4				
	Clindamycin	All ≤ 0.5				
	Metronidazole	All >16				
Vancomycin	0.5 to >64					

^aThe FDA breakpoint is applicable only to tigecycline. EUCAST epidemiological cutoff (ECOFF) breakpoints used for *C. difficile*. NA, not applicable.

^bOnly the MIC range is indicated when the total number of isolates is <10 .

isolates were positive for both *tet*(X1) and *tet*(X2); this set of isolates showed eravacycline, tigecycline, and minocycline MIC values ranging from 0.25 to 1 $\mu\text{g/ml}$, 2 to 8 $\mu\text{g/ml}$, and 2 to 32 $\mu\text{g/ml}$, respectively. No isolates were positive for *tet*(M).

In conclusion, this evaluation showed that eravacycline exhibited potent activity in

TABLE 3 *In vitro* activities of eravacycline and comparator tetracyclines against 27 *Bacteroides fragilis* clinical isolates characterized for tetracycline resistance determinants

No. of strains	Presence of tetracycline resistance determinant ^a					MIC range ($\mu\text{g/ml}$) ^b		
	<i>tet</i> (Q)	<i>tet</i> (X) ²	<i>tet</i> (X2) ²	<i>tet</i> (X1) ³	<i>tet</i> (M)	ERV	TIG	MIN
4	–	–	–	–	–	0.12–0.5	0.5–2	≤ 0.25
18	+ ⁴	–	–	–	–	0.06–4	0.25–16	0.5–32
3	+ ⁴	–	+	+	–	0.25–1	2–8	2–16
2	+	+	–	–	–	0.5–1	2–8	4–32

^a*tet*(X) was distinguished from *tet*(X2) by generating PCR products with universal *tet*(X) primers, sequencing with primers specific to both *tet*(X) and *tet*(X2), and identifying sequences specific to either *tet*(X) or *tet*(X2). *tet*(X1) was identified by generating PCR products with primers specific to *tet*(X1) and sequencing with the same primer set to verify *tet*(X1)-specific sequences. The *tet*(Q) gene in special studies laboratory strain numbers 28441 and 27741, with MIN MICs of 0.5 and 2 $\mu\text{g/ml}$, respectively, were sequenced and shown to encode amino acid sequences identical to each other and to 74 *Tet*(Q) proteins in the UniProtKB database (<http://www.uniprot.org/help/uniprotkb>, released on 15 March 2017), indicating that the genes encoded a common variant of *Tet*(Q).

^bMIC determined by agar dilution method; ERV, eravacycline; TIG, tigecycline; MIN, minocycline.

in vitro against MDR *B. fragilis* and other Gram-negative species, as well as Gram-positive anaerobic species, including isolates containing tetracycline-specific resistance determinants and isolates resistant to commonly used antibiotics, including carbapenems, fluoroquinolones, clindamycin, and β -lactam- β -lactamase inhibitor combinations. Based on its activity *in vitro*, eravacycline shows promise for the treatment of mixed aerobic-anaerobic infections, such as intra-abdominal infections.

ACKNOWLEDGMENTS

We thank Barbara Rapino and Carolina Baez Giangreco for technical assistant and Roselia Martinez for aid in the editing and graphical presentation of the manuscript.

Financial support for this study was provided by Tetrphase Pharmaceuticals, Inc.

At the time of the study, Kathryn Kerstein, Trudy H. Grossman, and Joyce A. Sutcliffe were employees of Tetrphase Pharmaceuticals, Inc.

REFERENCES

- Snydman DR, Jacobus NV, McDermott LA, Golan Y, Hecht DW, Goldstein EJ, Harrell L, Jenkins S, Newton D, Pierson C, Rihs JD, Yu VL, Venezia R, Finegold SM, Rosenblatt JE, Gorbach SL. 2010. Lessons learned from the anaerobe survey: historical perspective and review of the most recent data (2005–2007). *Clin Infect Dis* 50(Suppl 1):S26–S33. <https://doi.org/10.1086/647940>.
- Snydman DR, Jacobus NV, McDermott LA, Goldstein EJ, Harrell L, Jenkins SG, Newton D, Patel R, Hecht DW. 2017. Trends in antimicrobial resistance among *Bacteroides* species and *Parabacteroides* species in the United States from 2010–2012 with comparison to 2008–2009. *Anaerobe* 43:21–26. <https://doi.org/10.1016/j.anaerobe.2016.11.003>.
- Snydman DR, McDermott LA, Jacobus NV, Thorpe C, Stone S, Jenkins SG, Goldstein EJ, Patel R, Forbes BA, Mirrett S, Johnson S, Gerding DN. 2015. U.S.-based national sentinel surveillance study for the epidemiology of *Clostridium difficile*-associated diarrheal isolates and their susceptibility to fidaxomicin. *Antimicrob Agents Chemother* 59:6437–6443. <https://doi.org/10.1128/AAC.00845-15>.
- Solomkin JS, Ramesh MK, Cesnauskas G, Novikovs N, Stefanova P, Sutcliffe JA, Walpole SM, Horn PT. 2014. Phase 2, randomized, double-blind study of the efficacy and safety of two dose regimens of eravacycline versus ertapenem for adult community-acquired complicated intra-abdominal infections. *Antimicrob Agents Chemother* 58:1847–1854. <https://doi.org/10.1128/AAC.01614-13>.
- Solomkin J, Evans D, Slepavicius A, Lee P, Marsh A, Tsai L, Sutcliffe JA, Horn P. 2017. Assessing the efficacy and safety of eravacycline vs ertapenem in complicated intra-abdominal infections in the Investigating Gram-Negative Infections Treated With Eravacycline (IGNITE 1) trial: a randomized clinical trial. *JAMA Surg* 152:224–232. <https://doi.org/10.1001/jamasurg.2016.4237>.
- Grossman TH, Starosta AL, Fyfe C, O'Brien W, Rothstein DM, Mikolajka A, Wilson DN, Sutcliffe JA. 2012. Target- and resistance-based mechanistic studies with TP-434, a novel fluorocycline antibiotic. *Antimicrob Agents Chemother* 56:2559–2564. <https://doi.org/10.1128/AAC.06187-11>.
- Xiao XY, Hunt DK, Zhou J, Clark RB, Dunwoody N, Fyfe C, Grossman TH, O'Brien WJ, Plamondon L, Ronn M, Sun C, Zhang WY, Sutcliffe JA. 2012. Fluorocyclines. 1. 7-fluoro-9-pyrrolidinoacetamido-6-demethyl-6-deoxytetracycline: a potent, broad spectrum antibacterial agent. *J Med Chem* 55:597–605. <https://doi.org/10.1021/jm201465w>.
- Abdallah M, Olafisoye O, Cortes C, Urban C, Landman D, Quale J. 2015. Activity of eravacycline against *Enterobacteriaceae* and *Acinetobacter baumannii*, including multidrug-resistant isolates, from New York City. *Antimicrob Agents Chemother* 59:1802–1805. <https://doi.org/10.1128/AAC.04809-14>.
- Grossman T, O'Brien W, Fyfe C, Sutcliffe J. 2014. Eravacycline is potent against third generation cephalosporin- and carbapenem-resistant *Enterobacteriaceae*, carbapenem-resistant *Acinetobacter baumannii* and has isolate-specific bactericidal activity, abstr C-1374. Abstr 54th Intersci Conf Antimicrob Agents Chemother, 5 to 9 September 2014, Washington, DC.
- Sutcliffe JA, O'Brien W, Fyfe C, Grossman TH. 2013. Antibacterial activity of eravacycline (TP-434), a novel fluorocycline, against hospital and community pathogens. *Antimicrob Agents Chemother* 57:5548–5558. <https://doi.org/10.1128/AAC.01288-13>.
- Fyfe C, LeBlanc G, Close B, Nordmann P, Dumas J, Grossman TH. 2016. Eravacycline is active against bacterial isolates expressing the polymyxin resistance gene *mcr-1*. *Antimicrob Agents Chemother* 60:6989–6990. <https://doi.org/10.1128/AAC.01646-16>.
- Zhang Y, Lin X, Bush K. 2016. *In vitro* susceptibility of beta-lactamase-producing carbapenem-resistant *Enterobacteriaceae* (CRE) to eravacycline. *J Antibiot* 69:600–604. <https://doi.org/10.1038/ja.2016.73>.
- Livermore DM, Mushtaq S, Warner M, Woodford N. 2016. *In-vitro* activity of eravacycline against carbapenem-resistant *Enterobacteriaceae* and *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 60:3840–3844. <https://doi.org/10.1128/AAC.00436-16>.
- Bassetti M, Corey R, Doi Y, Morrissey I, Grossman T, Olesky M, Sutcliffe J. 2016. *In vitro* global surveillance of eravacycline and comparators against *Enterobacteriaceae*, *Acinetobacter baumannii*, *Stenotrophomonas maltophilia*, including multidrug-resistant (MDR) isolates, over a three-year period (2013–15), abstr 1825. *Infect Dis Soc Am, ID Week 2016*, 26 to 30 October 2016, New Orleans, LA.
- Bassetti M, Corey R, Doi Y, Morrissey I, Grossman T, Olesky M, Sutcliffe J. 2016. *In vitro* global surveillance of eravacycline and comparators against *Staphylococcus spp.* and *Enterococcus spp.* over a three-year period (2013–15), abstr 1820. *Infect Dis Soc Am, ID Week 2016*, 26 to 30 October 2016, New Orleans, LA.
- McDermott L, Jacobus NV, Snydman DR, Kerstein K, Grossman TH, Sutcliffe JA. 2015. Evaluation of the *in vitro* activity of eravacycline against a broad spectrum of recent clinical anaerobic isolates, abstr C-547. 55th Intersci Conf Antimicrob Agents Chemother/Int Cong Chemother Infect, 17 to 21 September 2015, San Diego, CA.
- Morrissey I, Sutcliffe J, Hackel M, Hawser S. 2015. Activity of eravacycline against anaerobic bacteria from Europe collected in 2013–14, abstr 784. *Infect Dis Soc Am, ID Week 2015*, 7 to 11 October 2015, San Diego, CA.
- Jousimies-Somer HR, Summanen P, Citron DM, Baron EJ, Wexler H, Finegold SM. 2002. *Wadsworth-KTL anaerobic bacteriology manual*, 6th ed. Star Publishing Co., Belmont, CA.
- CLSI. 2012. *Methods for antimicrobial susceptibility testing of anaerobic bacteria*; approved standard, 8th ed. CLSI document M11-A8. Clinical and Laboratory Standards Institute, Wayne, PA.
- CLSI. 2016. *Performance standards for antimicrobial susceptibility testing*, 26th ed. CLSI Document M100S. Clinical and Laboratory Standards Institute, Wayne, PA.
- T. M. Wyeth. 2014. *Product monograph. Tygacil (tigecycline for injection); sterile, lyophilized powder for intravenous use, 50 mg/vial*. T. M. Wyeth, Kirkland, Quebec City, Canada. http://www.pfizer.ca/sites/g/files/g10017036/f/201410/TYGACIL_PM_E_168980_17Jan2014.pdf.
- European Committee on Antimicrobial Susceptibility Testing (EUCAST). 2016. *Breakpoint tables for interpretation of MICs and zone diameters, version 6.0*. European Committee on Antimicrobial Susceptibility Testing, Växjö, Sweden. http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_6.0_Breakpoint_table.pdf.
- Connell SR, Tracz DM, Nierhaus KH, Taylor DE. 2003. *Ribosomal protection*.

- tion proteins and their mechanism of tetracycline resistance. *Antimicrob Agents Chemother* 47:3675–3681. <https://doi.org/10.1128/AAC.47.12.3675-3681.2003>.
24. Yang W, Moore IF, Koteva KP, Bareich DC, Hughes DW, Wright GD. 2004. TetX is a flavin-dependent monooxygenase conferring resistance to tetracycline antibiotics. *J Biol Chem* 279:52346–52352. <https://doi.org/10.1074/jbc.M409573200>.
25. Bartha NA, Soki J, Urban E, Nagy E. 2011. Investigation of the prevalence of *tetQ*, *tetX* and *tetX1* genes in *Bacteroides* strains with elevated tetracycline minimum inhibitory concentrations. *Int J Antimicrob Agents* 38:522–525. <https://doi.org/10.1016/j.ijantimicag.2011.07.010>.
26. Bradford PA, Petersen PJ, Tuckman M, Jones CH. 2008. *In vitro* activity of tigecycline and occurrence of tetracycline resistance determinants in isolates from patients enrolled in phase 3 clinical trials for community-acquired pneumonia. *Clin Microbiol Infect* 14:882–886. <https://doi.org/10.1111/j.1469-0691.2008.02063.x>.