



Characterization of β -Lactamase Content of Ceftazidime-Resistant Pathogens Recovered during the Pathogen-Directed Phase 3 REPRISÉ Trial for Ceftazidime-Avibactam: Correlation of Efficacy against β -Lactamase Producers

Rodrigo E. Mendes,^a Mariana Castanheira,^a Leah N. Woosley,^a Gregory G. Stone,^{b*}  Patricia A. Bradford,^{b*} Robert K. Flamm^a

^aJMI Laboratories, North Liberty, Iowa, USA

^bAstraZeneca Pharmaceuticals, Waltham, Massachusetts, USA

ABSTRACT REPRISÉ was a pathogen-directed (ceftazidime-resistant) phase 3 prospective, open-label, randomized, multicenter trial that evaluated the efficacy, safety, and tolerability of ceftazidime-avibactam (CAZ-AVI) and best available therapy (BAT) in the treatment of hospitalized adults with complicated intra-abdominal infections (cIAI) and complicated urinary tract infections (cUTI). This study characterized the β -lactamase content of ceftazidime-resistant *Enterobacteriaceae* and *Pseudomonas aeruginosa* recovered during the baseline visits of patients enrolled in REPRISÉ. Ceftazidime had MIC₉₀ results of >64 μ g/ml against baseline *Enterobacteriaceae* and *P. aeruginosa*. *bla*_{CTX-M} variants were the most common β -lactamases found in *Escherichia coli* (detected in 94.3% of all *E. coli* isolates) and *Klebsiella pneumoniae* (91.2%), whereas *Proteus mirabilis* often carried plasmid AmpC (pAmpC) (66.7%). *bla*_{KPC} (6 isolates), *bla*_{NDM-1} (3), *bla*_{OXA-48} (3), and *bla*_{VIM} (2) were detected in 4.9% (14/284) of *Enterobacteriaceae*. Overall, clinical cure rates against the *Enterobacteriaceae* were 91.2% and 90.8% for the CAZ-AVI and BAT groups, respectively, or 92.5% and 92.9% in the subset of patients infected with isolates harboring *bla*_{CTX-M}. Patients with baseline isolates carrying AmpC genes (pAmpC and/or overexpression of intrinsic AmpC) showed clinical cure rates of 80.0% and 89.5% for CAZ-AVI and BAT arms, respectively. Favorable microbiological responses were generally lower than clinical cure rates in both arms, but CAZ-AVI (80.0 to 85.0%) showed microbiological response rates consistently higher than those for BAT (57.9 to 64.3%) among patients with non-carbapenemase-producing *Enterobacteriaceae*. Lower microbiological response rates (50.0%) were found in patients with carbapenemase producers from both arms. This study expands on efficacy data analysis of CAZ-AVI among patients infected with ceftazidime-resistant pathogens, especially *bla*_{CTX-M}-carrying isolates, and although clinical cure rates for CAZ-AVI and BAT were similar, eradication rates for CAZ-AVI were higher than those for BAT. (This study has been registered at ClinicalTrials.gov under identifier NCT01644643.)

KEYWORDS CTX-M-15, ESBL, carbapenemase, clinical efficacy

Antimicrobial resistance is recognized as one of the most serious public health threats worldwide, and failure to address it could compromise modern medical advances (1). Currently, great concern is focused on emerging multidrug resistance in Gram-negative pathogens (2, 3). This concern relates to the increased prevalence of multidrug resistance among Gram-negative pathogens, including those producing extended-spectrum β -lactamase (ESBL) (4, 5), which led to a concomitant increase in carbapenem agent use and, consequently, increased selective pressure (6). The latter has led to the global emergence and dissemination of Gram-negative organisms

Citation Mendes RE, Castanheira M, Woosley LN, Stone GG, Bradford PA, Flamm RK. 2019. Characterization of β -lactamase content of ceftazidime-resistant pathogens recovered during the pathogen-directed phase 3 REPRISÉ trial for ceftazidime-avibactam: correlation of efficacy against β -lactamase producers. *Antimicrob Agents Chemother* 63:e02655-18. <https://doi.org/10.1128/AAC.02655-18>.

Copyright © 2019 American Society for Microbiology. All Rights Reserved.

Address correspondence to Rodrigo E. Mendes, rodrigo-mendes@jmilabs.com.

* Present address: Gregory G. Stone, Pfizer, Inc., Groton, Connecticut, USA; Patricia A. Bradford, Antimicrobial Development Specialists, LLC, Nyack, New York, USA.

Received 19 December 2018

Returned for modification 22 January 2019

Accepted 17 March 2019

Accepted manuscript posted online 25 March 2019

Published 24 May 2019

producing class A *Klebsiella pneumoniae* carbapenemase (KPC), class B metallo- β -lactamase (MBL), and/or class D carbapenemases (OXA-48-like), challenging antimicrobial therapy and increasing mortality (7–10).

Ceftazidime-avibactam (CAZ-AVI) was approved for complicated urinary tract infections (cUTI), including acute pyelonephritis; complicated intra-abdominal infections (cIAI); and hospital-acquired pneumonia, including ventilator-associated pneumonia (11, 12). Several phase 3 trials investigated the safety and efficacy of CAZ-AVI against standard clinical comparator agents (13–15). The molecular characterization of CAZ-AVI isolates from various clinical trials was previously reported (16, 17). The study presented here describes the characterization of the β -lactamase content of baseline pathogens recovered from a phase 3 prospective, open-label, randomized, multicenter trial to evaluate the efficacy, safety, and tolerability of CAZ-AVI and the best available therapy (BAT) in the treatment of hospitalized adults with cIAI and cUTI caused by ceftazidime-resistant Gram-negative pathogens, described here as those with MIC results of $\geq 8 \mu\text{g/ml}$ (15). Moreover, the efficacy results for CAZ-AVI and BAT were evaluated against subsets of molecularly characterized pathogens.

RESULTS

Bacterial pathogens and β -lactamase profiles. Isolates included in this study resulted in elevated ceftazidime MIC₅₀ and MIC₉₀ values (MIC_{50/90} of 64/>64 $\mu\text{g/ml}$ for *Enterobacteriaceae* and MIC_{50/90} of >64/>64 $\mu\text{g/ml}$ for *Pseudomonas aeruginosa*) (15). Overall, CTX-M-encoding genes alone or combined with other β -lactamase genes were the most prevalent resistance determinants (89.4%; 254/284) among *Enterobacteriaceae* (Table 1). *bla*_{CTX-M} variants were most commonly detected among *Escherichia coli* (94.3%) and *K. pneumoniae* (91.2%) isolates, followed by *Enterobacter* species (76.5%) and *Citrobacter freundii* (71.4%) isolates. In contrast, *Proteus mirabilis* often (66.7%) carried plasmid AmpC (pAmpC) (Table 1). Other species, such as *Klebsiella oxytoca*, *Providencia* spp., and *Serratia marcescens*, were isolated in numbers too small to provide any valuable analysis. *bla*_{OXA} noncarbapenemase genes (2 *bla*_{OXA-2}, 4 *bla*_{OXA-9}, 4 *bla*_{OXA-10}, and 175 *bla*_{OXA-1}) were also commonly observed (65.1%; 185/284) in *Enterobacteriaceae* isolates, whereas *bla*_{CTX-M} was associated with *bla*_{OXA-1} in 70.9% of isolates (180/254). Other ESBL genes, such as *bla*_{VEB-6} (1 isolate), *bla*_{SHV-2} (1), *bla*_{SHV-5} (3), *bla*_{SHV-12} (4), *bla*_{SHV-18} (2), *bla*_{SHV-38} (2), and *bla*_{TEM-15} (1), were less prevalent (4.9%; 14/284). A total of 4.9% (14/284) of *Enterobacteriaceae* isolates carried carbapenemase genes, including *bla*_{KPC} (6 isolates), *bla*_{NDM-1} (3), *bla*_{VIM} (2), and *bla*_{OXA-48} (3). These genes were mostly found in *K. pneumoniae*, except for 2 *bla*_{NDM-1} and 1 *bla*_{VIM-1} harboring isolate, which were detected in *Enterobacter cloacae*, *P. mirabilis*, and *Providencia rettgeri*. In addition, all carbapenemase-producing *Enterobacteriaceae* isolates were cultured from urine, except for 1 KPC-3-producing *K. pneumoniae* isolate. These isolates were collected from patients in Argentina (2 KPC-2-producing isolates), Bulgaria (1 NDM-1, 1 VIM-1, and 1 VIM-4), Israel (3 KPC-3), Romania (1 NDM-1 and 1 OXA-48), Russia (1 NDM-1), and Spain (1 KPC-3 and 2 OXA-48). *P. aeruginosa* exhibited mostly overexpression of intrinsic AmpC (44.4%; 8/18), with or without a variety of *bla*_{OXA}, *bla*_{PER}, and *bla*_{VEB} genes. One isolate carried *bla*_{VIM-2} (Table 1).

Efficacy analysis of CAZ-AVI and BAT. Similar clinical cure rates at the test-of-cure (TOC) visit were obtained for the CAZ-AVI (91.2 to 92.5%) and BAT (90.8 to 92.9%) groups that had patients infected with *Enterobacteriaceae* or patients with baseline isolates carrying only *bla*_{CTX-M} genes (β -lactamase genes other than AmpC and/or carbapenemase could be present with *bla*_{CTX-M}) (Table 2). Patients with baseline *Enterobacteriaceae* isolates carrying AmpC genes (pAmpC and/or overexpression of intrinsic AmpC) without noncarbapenemase β -lactamase genes showed clinical cure rates of 80.0% and 89.5% for CAZ-AVI and BAT arms, respectively. Patients with baseline *Enterobacteriaceae* isolates carrying carbapenemase genes were enrolled in a small number, but all patients in the CAZ-AVI arm showed clinical cure (all cUTI cases). These patients were infected with pathogens carrying *bla*_{KPC} ($n = 3$), *bla*_{VIM} ($n = 1$), *bla*_{NDM} ($n = 2$), and *bla*_{OXA-48} ($n = 2$), and the CAZ-AVI MIC results obtained against these

TABLE 1 β -Lactamase-encoding genes detected among ceftazidime-resistant baseline pathogens recovered from patients enrolled in both arms of the ceftazidime-avibactam phase 3 trial^e

Pathogen (no.; % of total)	β -Lactam class(es) ^a	Result(s) ^a	No. of isolates
<i>E. coli</i> (123; 40.7)	A, D	CTX-M-15; OXA-1	44
	A	CTX-M-15; TEM-1	25
	A, D	CTX-M-15; OXA-1; TEM-1	12
	A	CTX-M-15	10
	A	CTX-M-27	5
	A, C	CMY-2; TEM-1	5
	A, C, D	CTX-M-15; CMY-2; OXA-1; TEM-1	4
	A	CTX-M-27; TEM-1	2
	A	CTX-M-55	2
	A, C, D	CTX-M-15; CMY-42; OXA-1	2
	A	CTX-M-1	1
	A	CTX-M-14	1
	A	CTX-M-15; CTX-M-3; TEM-1	1
	A	CTX-M-2	1
	A	CTX-M-32	1
	A, C	DHA-1; SHV-12; TEM-1	1
	A, D	CTX-M-14; CTX-M-15; OXA-1	1
	A, D	CTX-M-15; CTX-M-27; OXA-1; TEM-1	1
	A, D	CTX-M-15; CTX-M-3; OXA-1	1
	A, D	CTX-M-15; OXA-1; TEM-33	1
A, D	CTX-M-3; OXA-1	1	
C	ACT-24	1	
<i>K. pneumoniae</i> (125; 41.4)	A, D	CTX-M-15; OXA-1; SHV-1; TEM-1	37
	A, D	CTX-M-15; OXA-1; SHV-11; TEM-1	27
	A, D	CTX-M-15; OXA-1; SHV-1	7
	A	CTX-M-15; SHV-1; TEM-1	6
	A, D	CTX-M-15; OXA-1; SHV-11	5
	A, D	CTX-M-15; OXA-10; SHV-1; TEM-1	4
	A	CTX-M-15; SHV-1	3
	A	CTX-M-15; CTX-M-3; SHV-1; TEM-1	3
	A	CTX-M-15; SHV-11; TEM-1	2
	A	CTX-M-3; SHV-11	2
	A	KPC-3; SHV-11	2
	A	KPC-3; SHV-11; TEM-1	2
	A, D	CTX-M-15; OXA-1; OXA-48; SHV-11	2
	A, D	OXA-2; SHV-18	2
	A	CTX-M-15; CTX-M-3; TEM-1	1
	A	CTX-M-15; SHV-11; SHV-12; TEM-1	1
	A	CTX-M-15; SHV-38; TEM-1	1
	A	CTX-M-2; SHV-11; TEM-1	1
	A	CTX-M-3; SHV-11; SHV-2; TEM-1	1
	A	CTX-M-3; SHV-11; TEM-1	1
	A	KPC-2; SHV-11; TEM-1	1
	A	SHV-5	1
	A, B	NDM-1; SHV-11	1
	A, B	SHV-11; TEM-1; VIM-4	1
	A, C	CMY-4; CTX-M-15; SHV-1	1
	A, C, D	CMY-4; CTX-M-15; OXA-1; SHV-1	1
	A, C, D	CTX-M-15; DHA-1; OXA-1; SHV-11; TEM-1	1
	A, D	CTX-M-15; CTX-M-3; OXA-1; SHV-1; TEM-1	1
	A, D	CTX-M-15; KPC-2; OXA-1; SHV-1; TEM-1	1
	A, D	CTX-M-15; OXA-1; OXA-48; SHV-1; TEM-1	1
A, D	CTX-M-15; OXA-1; SHV-1; SHV-11; TEM-1	1	
A, D	CTX-M-15; OXA-1; SHV-1; SHV-5; TEM-1	1	
A, D	CTX-M-15; OXA-1; SHV-38; TEM-1	1	
A, D	CTX-M-15; OXA-9; SHV-1; TEM-1	1	
A, D	OXA-9; SHV-11; TEM-1	1	
<i>Enterobacter</i> spp. (17; 5.6) ^b	A, C, D	CTX-M-15; cAmpC; OXA-1; TEM-1	5
	C	cAmpC	3
	A	CTX-M-15	1
	A, B, D	CTX-M-15; NDM-1; OXA-1; TEM-1	1
	A, C	CTX-M-15; cAmpC	1
	A, C	CTX-M-3; cAmpC; TEM-1	1

(Continued on next page)

TABLE 1 (Continued)

Pathogen (no.; % of total)	β -Lactam class(es) ^a	Result(s) ^a	No. of isolates
	A, C, D	CTX-M-15; CTX-M-3; cAmpC; OXA-1; TEM-1	1
	A, C, D	CTX-M-15; cAmpC; OXA-1	1
	A, C, D	cAmpC; OXA-1; SHV-12; TEM-1	1
	A, D	CTX-M-15; CTX-M-3; OXA-1; TEM-1	1
	A, D	CTX-M-15; OXA-1	1
<i>C. freundii</i> (7; 2.3)	A, C	cAmpC; TEM-1	1
	A, C, D	CMY-86; CTX-M-15; OXA-1; TEM-1	1
	A, C, D	CTX-M-15; CTX-M-3; cAmpC; OXA-1	1
	A, C, D	CTX-M-15; cAmpC; OXA-1; TEM-1	1
	A, D	CTX-M-15; OXA-1; TEM-1	1
	A, C	CTX-M-15; cAmpC; DHA-4; TEM-1	1
	C	cAmpC	1
<i>P. mirabilis</i> (6; 2.0)	A	TEM-1; VEB-6	1
	A, B, C	CMY-16; SHV-12; TEM-1; VIM-1	1
	A, C	ACC-4; TEM-1	1
	A, C	CMY-16; TEM-1	1
	A, D	CTX-M-3; OXA-9; SHV-5; TEM-1	1
	C	CMY-16	1
<i>K. oxytoca</i> (2; 0.7)	A, D	CTX-M-15; OXA-1; TEM-1	1
	A, D	CTX-M-15; OXA-9; TEM-15	1
<i>Providencia</i> spp. (2; 0.7) ^c	A, B	CTX-M-3; NDM-1; TEM-1	1
	A, C	ACC-4; TEM-1	1
<i>S. marcescens</i> (2; 0.7)	A, C, D	CMY-4; CTX-M-15; OXA-1; TEM-1	1
	A, D	CTX-M-15; CTX-M-3; OXA-1; TEM-1	1
<i>P. aeruginosa</i> (18; 6.0)	C	cAmpC	5
	A, D	OXA-10; VEB-9	3
	A, D	OXA-2; OXA-74 ^d ; PER-1	2
	A, D	OXA-2; PER-1	2
	A, C, D	cAmpC; OXA-10; VEB-9	1
	A, D	OXA-10; VEB-1	1
	A, D	OXA-74 ^d ; PER-1	1
	B	VIM-2	1
	C, D	cAmpC; OXA-17 ^d	1
	C, D	cAmpC; OXA-2	1

^aMolecular class according to Bush and Jacoby (26). cAmpC represents overexpression of the intrinsic chromosomal *ampC* gene according to reverse transcription-quantitative PCR (qRT-PCR) experiments.

^bIncludes 16 *E. cloacae* and 1 *Enterobacter aerogenes* isolate.

^cIncludes 1 *P. rettgeri* and 1 *P. stuartii* isolate.

^dOXA-10-like enzymes.

^eNote that five patients had 2 pathogens at the baseline visit.

isolates were 0.5 to 4 μ g/ml, except against *bla*_{NDM-1}-harboring strains (CAZ-AVI MIC, >256 μ g/ml). In the BAT arm, carbapenemase-producing *Enterobacteriaceae* isolates carried *bla*_{KPC-3} ($n = 3$), *bla*_{VIM} ($n = 1$), *bla*_{NDM-1} ($n = 1$), and *bla*_{OXA-48} ($n = 1$) (Table 2). Patients with unfavorable clinical and microbiological outcomes in both study arms are described in Tables 3 and 4. Correlations were not apparent between β -lactamases present in the clinical isolates or between the CAZ-AVI MIC and the result of an unfavorable response in patients treated with CAZ-AVI; however, 4/8 patients infected with carbapenemase-producing *Enterobacteriaceae* in the CAZ-AVI arm had an unfavorable microbiological response (Tables 2 and 4). Patients infected with *P. aeruginosa* showed clinical cure rates of 84.6% and 100.0% for CAZ-AVI and BAT arms, respectively (Table 2). One patient in the CAZ-AVI arm had a VIM-2-harboring *P. aeruginosa* isolate with a CAZ-AVI MIC of 32 μ g/ml, and a clinical cure as well as a favorable microbiological response were documented.

For patients enrolled in the CAZ-AVI arm (82.6%), favorable microbiological responses tended to be lower than clinical cure rates (90.6%). A similar trend was

TABLE 2 Clinical and microbiological responses at the test-of-cure visit by β -lactamase status (mMITT population)

Patient subgroup	Cure rate at test of cure						Comparison between groups		
	CAZ-AVI (n = 149)			BAT (n = 146)			No. (%) of patients with microbiological cure ^b	% difference of clinical cure rate ^c	% difference of microbiological cure rate ^d
	No. of patients	No. (%) of patients with clinical cure ^a	No. (%) of patients with microbiological cure ^b	No. of patients	No. (%) of patients with clinical cure ^a	No. (%) of patients with microbiological cure ^b			
All	149	135 (90.6)	123 (82.6)	146	133 (91.1)	92 (63.0)	-0.5	19.6	
<i>Enterobacteriaceae</i> ^e	136	124 (91.2)	112 (82.4)	141	128 (90.8)	89 (63.1)	+0.4	+19.3	
CTX-M ^f	107	99 (92.5)	91 (85.0)	112	104 (92.9)	72 (64.3)	-0.4	+20.7	
AmpC/ESBL ^g	20	16 (80.0)	16 (80.0)	19	17 (89.5)	11 (57.9)	-9.5	+22.1	
Carbapenemase/ESBL	8 ^h	8 (100.0)	4 (50.0)	6 ⁱ	4 (66.7)	3 (50.0)	+33.3	0.0	
<i>P. aeruginosa</i>	13 ^j	11 (84.6)	11 (84.6)	5	5 (100.0)	3 (60.0)	-15.4	+24.6	

^aPercentages for clinical cure rates are calculated as m/n , where m is defined by the number of patients with a favorable clinical response at the test-of-cure visit and n is represented by the combined number of patients with clinical cure, clinical failure, and an indeterminate outcome.
^bPercentages for microbiological cure rates are calculated as m/n , where m is defined by the number of patients with a favorable microbiological response at the test-of-cure visit and n is represented by the combined number of patients with microbiological cure, failure, and an indeterminate outcome.
^cThe difference of the favorable clinical cure rate is calculated as CAZ-AVI treatment group minus BAT treatment group.
^dDifference of favorable microbiological responses is calculated as CAZ-AVI treatment group minus BAT treatment group.
^eTwo and three patients had polymicrobial infections in the CAZ-AVI and BAT groups, respectively.
^fIncludes patients with isolates carrying CTX-M-encoding genes with or without non-ESBL and/or ESBL genes (OXA-1, OXA-10, OXA-9, SHV-11, SHV-12, SHV-5, SHV-1, and TEM-1) (Table 1).
^gIncludes patients with isolates with documented overexpression of the intrinsic chromosomal AmpC gene, carrying plasmid AmpC, and/or with multiple non-ESBL and/or ESBL genes (Table 1).
^hIncludes KPC (3 isolates), VIM (1), NDM (2), and OXA-48 (2)-encoding genes.
ⁱIncludes KPC (3 isolates), VIM (1), NDM (1), and OXA-48 (1)-encoding genes.
^jOne patient had a polymicrobial cUTI (*P. aeruginosa* and *P. stuartii*) at the baseline visit.

TABLE 3 Patients with baseline isolates who presented clinical failure or had an indeterminate status at TOC^a

Arm and pathogen	Country	Infection type	CAZ-AVI MIC ($\mu\text{g/ml}$)	Molecular characterization
CAZ-AVI				
<i>K. pneumoniae</i>	Bulgaria	cUTI	0.5	CTX-M-15; SHV-1
<i>P. aeruginosa</i>	Bulgaria	cUTI	16	OXA-2; OXA-74; PER-1
<i>E. coli</i>	Croatia	cUTI	0.12	CTX-M-15
<i>P. mirabilis</i>	Croatia	cUTI	0.06	CMY-16
<i>E. coli</i>	Israel	cUTI	0.5	CTX-M-15; OXA-1
<i>E. coli</i>	Romania	cUTI	0.12	CTX-M-15; TEM-1
<i>P. mirabilis</i>	Romania	cUTI	0.5	ACC-4; TEM-1
<i>E. cloacae/K. pneumoniae</i>	Russia	cIAI	1/0.25	cAmpC/CTX-M-15; OXA-1; SHV-1; TEM-1
<i>E. coli</i>	Russia	cIAI	0.12	CTX-M-15; TEM-1
<i>E. coli</i>	Russia	cUTI	0.25	CTX-M-15; OXA-1
<i>E. coli</i>	South Africa	cUTI	0.12	CTX-M-15; OXA-1
<i>E. cloacae</i>	Ukraine	cUTI	0.25	CTX-M-15; cAmpC
<i>E. coli</i>	Ukraine	cUTI	0.06	CTX-M-15; TEM-1
<i>P. aeruginosa</i>	Ukraine	cUTI	64	OXA-10; VEB-1
BAT				
<i>K. pneumoniae</i>	Argentina	cUTI	NA	KPC-2; SHV-11; TEM-1
<i>E. coli</i>	Bulgaria	cIAI	NA	CTX-M-15; OXA-1
<i>K. pneumoniae</i>	Bulgaria	cUTI	NA	SHV-11; TEM-1; VIM-4
<i>E. cloacae</i>	Croatia	cUTI	NA	CTX-M-15; cAmpC; OXA-1; TEM-1
<i>E. cloacae</i>	Israel	cIAI	NA	CTX-M-15; OXA-1
<i>E. coli</i>	Israel	cIAI	NA	CTX-M-15; OXA-1
<i>E. coli</i>	Israel	cUTI	NA	CTX-M-15; OXA-1
<i>K. pneumoniae</i>	Romania	cUTI	NA	CTX-M-15; OXA-1; SHV-1
<i>E. coli</i>	Russia	cIAI	NA	CTX-M-15; OXA-1
<i>K. pneumoniae</i>	Russia	cIAI	NA	OXA-2; SHV-18
<i>E. coli</i>	Spain	cUTI	NA	CTX-M-27
<i>E. coli</i>	Turkey	cUTI	NA	CTX-M-15; TEM-1
<i>K. pneumoniae</i>	Ukraine	cUTI	NA	CMY-4; CTX-M-15; SHV-1

^aTOC, test of cure; CAZ-AVI, ceftazidime-avibactam; BAT, best available therapy; cUTI, complicated urinary tract infection; cIAI, complicated intra-abdominal infection; NA, not available. cAmpC represents overexpression of the intrinsic chromosomal *ampC* gene according to qRT-PCR experiments.

observed among patients enrolled in the BAT arm (63.0% and 91.1%, respectively) (Table 2). The favorable microbiological responses observed among patients enrolled in the CAZ-AVI arm (80.0 to 85.0%) were consistently higher than those noted for BAT (57.9 to 64.3%), except among patients infected with carbapenemase-producing *Enterobacteriaceae*, where favorable microbiological responses of 50.0% were documented in both arms (Table 2).

DISCUSSION

These study results confirm the high occurrence of *bla*_{CTX-M} among clinical trial *Enterobacteriaceae* isolates causing cIAI or cUTI. The majority of *bla*_{CTX-M} variants were *bla*_{CTX-M-15}, which corroborates other clinical trials (16–18) and surveillance studies (8, 19, 20) demonstrating the dominance of *bla*_{CTX-M-15} among *Enterobacteriaceae* (21, 22). This ESBL gene has disseminated globally; however, group 9 variants (especially CTX-M-14) appear dominant in China, Southeast Asia, South Korea, Japan, and Spain (21). Carbapenemase genes remained infrequently isolated (4.6%) among ceftazidime-resistant *Enterobacteriaceae* causing cIAI or cUTI. The occurrence of such isolates was lower than that observed in a surveillance study among cUTI and cIAI *Enterobacteriaceae* isolates in Europe (11.0%) (23). The lower carbapenemase rate in ceftazidime-resistant isolates in this clinical trial may be attributed to the respective sites included in the study or to the inclusion/exclusion criteria used to enroll each patient.

This study demonstrated similar clinical cure rates for CAZ-AVI and BAT (90.8 to 91.2%) in cIAI or cUTI caused by ceftazidime-resistant *Enterobacteriaceae*. These efficacy results also extended to the subset of pathogens carrying *bla*_{CTX-M} (92.5 to 92.9%). The analysis generated here showed clinical cure rates for CAZ-AVI lower than those for BAT on two occasions, when *Enterobacteriaceae* isolates overexpressing intrinsic AmpC or carrying pAmpC-encoding genes (80.0 versus 89.5% for BAT) were present or *P.*

TABLE 4 Patients with baseline isolates who presented an unfavorable microbiological response or had an indeterminate status at TOC^a

Arm and pathogen	Country	Infection	CAZ-AVI MIC ($\mu\text{g/ml}$)	Molecular characterization
CAZ-AVI				
<i>C. freundii</i>	Bulgaria	cUTI	0.5	CTX-M-15; OXA-1; TEM-1
<i>E. cloacae</i>	Bulgaria	cUTI	2	cAmpC; OXA-1; SHV-12; TEM-1
<i>E. cloacae</i>	Romania	cUTI	>256	CTX-M-15; NDM-1; OXA-1; TEM-1
<i>E. cloacae/K. pneumoniae</i>	Russia	cIAI	1/0.25	cAmpC/CTX-M-15; OXA-1; SHV-1; TEM-1
<i>E. coli</i>	Croatia	cUTI	0.12	CTX-M-15
<i>E. coli</i>	Israel	cUTI	0.5	CTX-M-15; OXA-1
<i>E. coli</i>	Romania	cUTI	0.06	CTX-M-15; OXA-1
<i>E. coli</i>	Russia	cUTI	0.25	CTX-M-15; OXA-1
<i>E. coli</i>	Romania	cUTI	0.12	CTX-M-15; TEM-1
<i>E. coli</i>	Russia	cIAI	0.12	CTX-M-15; TEM-1
<i>E. coli</i>	Russia	cUTI	≤ 0.008	CTX-M-15; TEM-1
<i>E. coli</i>	Ukraine	cUTI	0.06	CTX-M-15; TEM-1
<i>K. pneumoniae</i>	Bulgaria	cUTI	1	CTX-M-15; OXA-1; SHV-1; TEM-1
<i>K. pneumoniae</i>	Turkey	cUTI	0.25	CTX-M-15; OXA-1; SHV-1; TEM-1
<i>K. pneumoniae</i>	Turkey	cUTI	0.5	CTX-M-15; OXA-1; SHV-1; TEM-1
<i>K. pneumoniae</i>	Russia	cUTI	0.5	CTX-M-15; OXA-1; SHV-11; TEM-1
<i>K. pneumoniae</i>	Russia	cUTI	0.5	CTX-M-15; OXA-1; SHV-11; TEM-1
<i>K. pneumoniae</i>	Bulgaria	cUTI	0.12	CTX-M-15; OXA-1; SHV-38; TEM-1
<i>K. pneumoniae</i>	Bulgaria	cUTI	0.5	CTX-M-15; SHV-1
<i>K. pneumoniae</i>	Israel	cUTI	4	KPC-3; SHV-11
<i>K. pneumoniae</i>	Bulgaria	cUTI	>256	NDM-1; SHV-11
<i>P. aeruginosa</i>	Russia	cUTI	4	cAmpC
<i>P. aeruginosa</i>	Ukraine	cUTI	64	OXA-10; VEB-1
<i>P. mirabilis</i>	Romania	cUTI	0.5	ACC-4; TEM-1
<i>P. mirabilis</i>	Croatia	cUTI	0.06	CMY-16
<i>P. mirabilis</i>	Bulgaria	cUTI	0.5	CMY-16; SHV-12; TEM-1; VIM-1
BAT				
<i>C. freundii</i>	France	cUTI	NA	cAmpC; TEM-1
<i>C. freundii</i>	Bulgaria	cUTI	NA	CTX-M-15; CTX-M-3; cAmpC; OXA-1
<i>E. cloacae</i>	Bulgaria	cUTI	NA	CTX-M-15; CTX-M-3; cAmpC; OXA-1; TEM-1
<i>E. cloacae</i>	Israel	cIAI	NA	CTX-M-15; OXA-1
<i>E. coli</i>	Israel	cUTI	NA	CMY-2; TEM-1
<i>E. coli</i>	United States	cUTI	NA	CMY-2; TEM-1
<i>E. coli</i>	Russia	cUTI	NA	CMY-42; CTX-M-15; OXA-1
<i>E. coli</i>	Croatia	cUTI	NA	CTX-M-15
<i>E. coli</i>	Peru	cUTI	NA	CTX-M-15
<i>E. coli</i>	Bulgaria	cIAI	NA	CTX-M-15; OXA-1
<i>E. coli</i>	Bulgaria	cUTI	NA	CTX-M-15; OXA-1
<i>E. coli</i>	Israel	cIAI	NA	CTX-M-15; OXA-1
<i>E. coli</i>	Israel	cUTI	NA	CTX-M-15; OXA-1
<i>E. coli</i>	Mexico	cUTI	NA	CTX-M-15; OXA-1
<i>E. coli</i>	Romania	cUTI	NA	CTX-M-15; OXA-1
<i>E. coli</i>	Russia	cIAI	NA	CTX-M-15; OXA-1
<i>E. coli</i>	Turkey	cUTI	NA	CTX-M-15; OXA-1
<i>E. coli</i>	Bulgaria	cUTI	NA	CTX-M-15; OXA-1; TEM-1
<i>E. coli</i>	Romania	cUTI	NA	CTX-M-15; OXA-1; TEM-1
<i>E. coli</i>	Bulgaria	cUTI	NA	CTX-M-15; TEM-1
<i>E. coli</i>	Bulgaria	cUTI	NA	CTX-M-15; TEM-1
<i>E. coli</i>	Turkey	cUTI	NA	CTX-M-15; TEM-1
<i>E. coli</i>	Turkey	cUTI	NA	CTX-M-15; TEM-1
<i>E. coli</i>	Argentina	cUTI	NA	CTX-M-2
<i>E. coli/E. coli</i>	Turkey	cUTI	NA	CTX-M-15; CTX-M-3; TEM-1/CTX-M-15; TEM-1
<i>K. oxytoca</i>	Romania	cUTI	NA	CTX-M-15; OXA-1; TEM-1
<i>K. oxytoca</i>	Bulgaria	cUTI	NA	CTX-M-15; OXA-9; TEM-15
<i>K. pneumoniae</i>	Bulgaria	cUTI	NA	CMY-4; CTX-M-15; OXA-1; SHV-1
<i>K. pneumoniae</i>	Ukraine	cUTI	NA	CMY-4; CTX-M-15; SHV-1
<i>K. pneumoniae</i>	Bulgaria	cUTI	NA	CTX-M-15; CTX-M-3; SHV-1; TEM-1
<i>K. pneumoniae</i>	Romania	cUTI	NA	CTX-M-15; OXA-1; SHV-1
<i>K. pneumoniae</i>	Russia	cUTI	NA	CTX-M-15; OXA-1; SHV-1
<i>K. pneumoniae</i>	Bulgaria	cUTI	NA	CTX-M-15; OXA-1; SHV-1; SHV-11; TEM-1
<i>K. pneumoniae</i>	Bulgaria	cUTI	NA	CTX-M-15; OXA-1; SHV-1; TEM-1
<i>K. pneumoniae</i>	Peru	cUTI	NA	CTX-M-15; OXA-1; SHV-1; TEM-1
<i>K. pneumoniae</i>	Romania	cUTI	NA	CTX-M-15; OXA-1; SHV-1; TEM-1
<i>K. pneumoniae</i>	United States	cUTI	NA	CTX-M-15; OXA-1; SHV-1; TEM-1

(Continued on next page)

TABLE 4 (Continued)

Arm and pathogen	Country	Infection	CAZ-AVI MIC ($\mu\text{g/ml}$)	Molecular characterization
<i>K. pneumoniae</i>	Bulgaria	cUTI	NA	CTX-M-15; OXA-1; SHV-11
<i>K. pneumoniae</i>	Bulgaria	cUTI	NA	CTX-M-15; OXA-1; SHV-11; TEM-1
<i>K. pneumoniae</i>	Bulgaria	cUTI	NA	CTX-M-15; OXA-1; SHV-11; TEM-1
<i>K. pneumoniae</i>	Bulgaria	cUTI	NA	CTX-M-15; SHV-1
<i>K. pneumoniae</i>	Romania	cUTI	NA	CTX-M-15; SHV-1
<i>K. pneumoniae</i>	Bulgaria	cUTI	NA	CTX-M-15; SHV-1; TEM-1
<i>K. pneumoniae</i>	Romania	cUTI	NA	CTX-M-15; SHV-11; SHV-12; TEM-1
<i>K. pneumoniae</i>	Bulgaria	cUTI	NA	CTX-M-15; SHV-38; TEM-1
<i>K. pneumoniae</i>	Bulgaria	cUTI	NA	CTX-M-3; SHV-11
<i>K. pneumoniae</i>	Argentina	cUTI	NA	KPC-2; SHV-11; TEM-1
<i>K. pneumoniae</i>	Russia	cIAI	NA	OXA-2; SHV-18
<i>K. pneumoniae</i>	Bulgaria	cUTI	NA	SHV-11; TEM-1; VIM-4
<i>P. aeruginosa</i>	Turkey	cUTI	NA	cAmpC
<i>P. aeruginosa</i>	Turkey	cUTI	NA	OXA-2; PER-1
<i>P. mirabilis</i>	Argentina	cUTI	NA	CTX-M-3; OXA-9; SHV-5; TEM-1
<i>P. rettgeri</i>	Russia	cUTI	NA	CTX-M-3; NDM-1; TEM-1
<i>S. marcescens</i>	Bulgaria	cUTI	NA	CTX-M-15; CTX-M-3; OXA-1; TEM-1

^aTOC, test of cure; CAZ-AVI, ceftazidime-avibactam; BAT, best available therapy; cUTI, complicated urinary tract infection; cIAI, complicated intra-abdominal infection; NA, not available. cAmpC represents overexpression of the intrinsic chromosomal *ampC* gene according to qRT-PCR experiments.

aeruginosa (84.6 versus 100.0% for BAT) was present. The reasons for these results are unclear, and the number of such isolates remained low, compromising data analysis. Clinical cure rates presented here against AmpC-producing isolates were similar to those (75.0 to 86.7%) obtained in a IAI phase 3 trial for CAZ-AVI and comparator agents (17). Also in agreement were the small number of patients infected with such isolates in this and the previous IAI phase 3 trial (17).

REPRISE (ClinicalTrials.gov identifier NCT01644643) was the first pathogen-directed study, and only 14 patients infected with carbapenemase-producing *Enterobacteriaceae* pathogens were enrolled. This number of patients was likely due to the high occurrence of ESBL-producing isolates, even when including participating medical centers with a history of elevated rates of isolates carrying a variety of carbapenemase genes (e.g., Europe and adjacent regions) (24). All patients in the CAZ-AVI arm infected by carbapenemase-producing pathogens (including a patient with *P. aeruginosa*) had favorable clinical cure rates, including those infected with MBL-producing isolates (2 NDM-1 [CAZ-AVI MIC, >256 $\mu\text{g/ml}$], 1 VIM-2 [CAZ-AVI MIC, 32 $\mu\text{g/ml}$], and 1 VIM-1 [CAZ-AVI MIC, 1 $\mu\text{g/ml}$]). However, patients infected with the NDM-1- and VIM-1-producing isolates in the CAZ-AVI arm had unfavorable microbiological responses at the TOC. Although clinical improvement was seen in some patients infected with MBL-producing pathogens, persistence of the causative pathogen may occur.

CAZ-AVI (80.0 to 85.0%) demonstrated favorable microbiological responses consistently higher than those with BAT (57.9 to 64.3%) among patients with non-carbapenemase-producing *Enterobacteriaceae*. Although the vast majority of patients included here had cUTI, similar microbiological responses were obtained at TOC for CAZ-AVI (62.7%) and doripenem (60.7%) against ceftazidime-resistant isolates during the CAZ-AVI phase 3 trials (RECAPTURE 1 and 2) for cUTI (13). Differences in trial designs limit direct comparisons, and the reasons for these discrepancies are unknown. Nonetheless, imipenem was the agent most utilized as a BAT in the REPRISE trial, and most unfavorable microbiological responses were associated with this drug (15). In addition, the overall clinical cure rates observed here for both arms (90.6 to 91.1%) were higher than the favorable microbiological responses (63.0 to 82.6%). Similar findings were observed in the RECAPTURE phase 3 trials for cUTI, where the clinical cure rates (90.8 to 90.5%) were higher than the microbiological responses (64.0 to 60.0%) at TOC when ceftazidime-resistant isolates were present (13).

The small number of patients infected with carbapenemase-producing isolates limits the interpretation of the clinical and microbiological outcome findings in this study. However, the study results presented here add valuable information related to

the treatment of serious cUTI and cIAI caused by ceftazidime-resistant *Enterobacteriaceae*. Clinical cure rates for CAZ-AVI were similar to those obtained for the BAT arm against the overall ceftazidime-resistant *Enterobacteriaceae* population, whereas favorable microbiological responses for CAZ-AVI were consistently higher than those for BAT. These results expand on those reported previously (16, 17) by providing additional patients infected with ceftazidime-resistant pathogens, especially *bla*_{CTX-M}-carrying isolates. Moreover, the data analysis presented here indicates that CAZ-AVI may be used as an alternative agent to treat cUTI and cIAI.

MATERIALS AND METHODS

Patients, clinical isolates, study treatment, and endpoints. REPRIS (ClinicalTrials.gov identifier NCT01644643) was a prospective, international, randomized, open-label, phase 3 trial. Eligible patients were randomized in a 1:1 ratio to receive 5 to 21 days of treatment with either CAZ-AVI (2,000 mg CAZ–500 mg AVI), administered together as a 2-h intravenous (i.v.) infusion every 8 h, or BAT. Patients were stratified by entry diagnosis (cUTI and cIAI) and by region: (i) North America and Western Europe, (ii) Eastern Europe, and (iii) other regions. The investigator determined the BAT, which was based on the standard of care and local label recommendations, and BAT was documented before randomization. Patients with cUTI had 2 follow-up visits, at 21 to 25 days (FU1) and at 28 to 32 days (FU2) from randomization. Patients with cIAI had only 1 follow-up visit at 28 to 35 days from randomization (FU1). Carmeli et al. (15) provide additional information related to the clinical trial.

A total of 284 baseline *Enterobacteriaceae* and 18 *P. aeruginosa* isolates with elevated ceftazidime MIC results (≥ 8 $\mu\text{g/ml}$; referred to here as ceftazidime resistant) causing cIAI/cUTI in the microbiological modified intention-to-treat (mMITT) population were included. These baseline isolates were recovered during the first patient visit from 295 patients (aged 18 to 90 years) hospitalized in the Americas (Argentina, 8 subjects; Mexico, 7; Peru, 5; United States, 5), Europe (Bulgaria, 86; Croatia, 12; Czech Republic, 6; France, 3; Spain, 4; Ukraine, 28), Russia (66), Turkey (22), Israel (15), South Africa (2), and South Korea (2). When multiple isolates of the same species and demonstrating the same pulsed-field gel electrophoresis (PFGE) or multilocus sequence typing (MLST) profile were obtained from a patient, only 1 isolate was included. Six patients (3 from each arm) had 2 isolates of different species recovered at the baseline visit, and both isolates from each patient were included in the analysis. The primary endpoint was assessment of clinical response (cure, failure, or indeterminate) at the TOC visit 7 to 10 days after the last infusion of study therapy in the mMITT population, whereas a favorable microbiological response was defined by the eradication or presumed eradication of the causative pathogen (15).

Susceptibility testing, selection criteria, and screening of β -lactamases. All baseline clinical isolates were centrally tested for susceptibility by broth microdilution (Clinical and Laboratory Standards Institute [CLSI] document M07-A10, 2015) (25). *Enterobacteriaceae* were selected according to preestablished MIC criteria and subjected to screening for non-ESBL-, ESBL-, pAmpC-, and carbapenemase-encoding genes, and enzymes were assigned based on amino acid identity, as previously described (16, 17). The transcription levels of chromosomally encoded AmpC were determined in *Enterobacter* spp., *Citrobacter* spp., and *P. aeruginosa* by quantifying the target gene mRNA level using a normalized expression analysis method and relative comparison to susceptible control strains (16, 17). A given isolate was determined to overexpress the *ampC* gene when at least a 10-fold greater difference of *ampC* transcripts was detected than with a species-specific wild-type reference control strain.

Statistical analyses. Descriptive summaries are provided for clinical and microbiological responses at the TOC visit between groups. Analyses were performed between arms stratified by group of organisms/species, phenotype, and β -lactamase resistance mechanisms of baseline isolates.

Data availability. Upon request, and subject to certain criteria, conditions, and exceptions (see <https://www.pfizer.com/science/clinical-trials/trial-data-and-results> for more information), Pfizer will provide access to individual deidentified participant data from Pfizer-sponsored global interventional clinical studies conducted for medicines, vaccines, and medical devices (i) for indications that have been approved in the United States and/or European Union or (ii) in programs that have been terminated (i.e., development for all indications has been discontinued). Pfizer will also consider requests for the protocol, data dictionary, and statistical analysis plan. Data may be requested from Pfizer trials 24 months after study completion. The deidentified participant data will be made available to researchers whose proposals meet the research criteria and other conditions, and for which an exception does not apply, via a secure portal. To gain access, data requestors must enter into a data access agreement with Pfizer.

ACKNOWLEDGMENTS

We express appreciation to the following persons for significant contributions to the manuscript: L. Deshpande and T. Doyle.

The REPRIS study was originally sponsored by AstraZeneca and is now sponsored by Pfizer. AstraZeneca's rights to ceftazidime-avibactam were acquired by Pfizer in December 2016. This analysis was performed by JMI Laboratories and supported by AstraZeneca, which included funding for services related to preparing the manuscript. JMI Laboratories, Inc., has received research and educational grants from Achaogen,

Actelion, Allegra, Allergan, Ampliphi, API, Astellas, AstraZeneca, Basilea, Bayer, BD, Biomodels, Cardeas, CEM-102 Pharma, Cempra, Cidara, Cormedix, CSA Biotech, Cubist, Debiopharm, Dipexium, Duke, Durata, Entasis, Fortress, Fox Chase Chemical, GSK, Medpace, Melinta, Merck, Micurx, Motif, N8 Medical, Nabriva, Nexcida, Novartis, Paratek, Pfizer, Polyphor, Rempex, Scynexis, Shionogi, Spero Therapeutics, Symbal Therapeutics, Synolgoic, TGV Therapeutics, The Medicines Company, Theravance, Thermo Fisher, Venatorx, Wockhardt, and Zavante.

Some JMI employees are advisors/consultants for Allergan, Astellas, Cubist, Pfizer, Cempra, and Theravance. Regarding speakers' bureaus and stock options, we have none to declare. R.E.M., M.C., L.N.W., and R.K.F. are employees of JMI Laboratories, which received financial support from AstraZeneca in connection with the development of the manuscript. P.A.B. and G.G.S. were employees and shareholders of AstraZeneca at the time when the phase 3 clinical trials and the present analysis were undertaken. G.G.S. is currently an employee of Pfizer.

REFERENCES

- Laxminarayan R, Matsoso P, Pant S, Brower C, Rottingen JA, Klugman K, Davies S. 2016. Access to effective antimicrobials: a worldwide challenge. *Lancet* 387:168–175. [https://doi.org/10.1016/S0140-6736\(15\)00474-2](https://doi.org/10.1016/S0140-6736(15)00474-2).
- Friedman ND, Temkin E, Carmeli Y. 2016. The negative impact of antibiotic resistance. *Clin Microbiol Infect* 22:416–422. <https://doi.org/10.1016/j.cmi.2015.12.002>.
- Doi Y, Bonomo RA, Hooper DC, Kaye KS, Johnson JR, Clancy CJ, Thaden JT, Stryjewski ME, van Duin D, Gram-Negative Committee of the Antibacterial Resistance Leadership Group a. 2017. Gram-negative bacterial infections: research priorities, accomplishments, and future directions of the antibacterial resistance leadership group. *Clin Infect Dis* 64:S30–S35. <https://doi.org/10.1093/cid/ciw829>.
- Johnson JR, Johnston B, Clabots C, Kuskowski MA, Castanheira M. 2010. *Escherichia coli* sequence type ST131 as the major cause of serious multidrug-resistant *E. coli* infections in the United States. *Clin Infect Dis* 51:286–294. <https://doi.org/10.1086/653932>.
- Castanheira M, Mendes RE, Rhomberg PR, Jones RN. 2008. Rapid emergence of *bla*_{CTX-M} among *Enterobacteriaceae* in U.S. medical centers: molecular evaluation from the MYSTIC program (2007). *Microb Drug Resist* 14:211–216. <https://doi.org/10.1089/mdr.2008.0827>.
- Chang HJ, Hsu PC, Yang CC, Kuo AJ, Chia JH, Wu TL, Lee MH. 2011. Risk factors and outcomes of carbapenem-nonsusceptible *Escherichia coli* bacteremia: a matched case-control study. *J Microbiol Immunol Infect* 44:125–130. <https://doi.org/10.1016/j.jmii.2010.06.001>.
- Castanheira M, Farrell SE, Deshpande LM, Mendes RE, Jones RN. 2013. Prevalence of beta-lactamase-encoding genes among *Enterobacteriaceae* bacteremia isolates collected in 26 U.S. hospitals: report from the SENTRY antimicrobial surveillance program (2010). *Antimicrob Agents Chemother* 57:3012–3020. <https://doi.org/10.1128/AAC.02252-12>.
- Castanheira M, Farrell SE, Krause KM, Jones RN, Sader HS. 2014. Contemporary diversity of beta-lactamases among *Enterobacteriaceae* in the nine U.S. census regions and ceftazidime-avibactam activity tested against isolates producing the most prevalent beta-lactamase groups. *Antimicrob Agents Chemother* 58:833–838. <https://doi.org/10.1128/AAC.01896-13>.
- Falagas ME, Tansarli GS, Karageorgopoulos DE, Vardakas KZ. 2014. Deaths attributable to carbapenem-resistant *Enterobacteriaceae* infections. *Emerg Infect Dis* 20:1170–1175. <https://doi.org/10.3201/eid2007.121004>.
- Gupta N, Limbago BM, Patel JB, Kallen AJ. 2011. Carbapenem-resistant *Enterobacteriaceae*: epidemiology and prevention. *Clin Infect Dis* 53:60–67. <https://doi.org/10.1093/cid/cir202>.
- AstraZeneca AB. 2016. Zavicefta package insert. AstraZeneca AB, Waltham, MA. http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/004027/WC500210234.pdf. Accessed September 26, 2016.
- Allergan USA, Inc. 2017. Avycaz (ceftazidime-avibactam). Allergan USA, Inc, Madison, NJ. https://www.allergan.com/assets/pdf/avycaz_pi. Accessed February 2, 2017.
- Wagenlehner FM, Sobel JD, Newell P, Armstrong J, Huang X, Stone GG, Yates K, Gasink LB. 2016. Ceftazidime-avibactam versus doripenem for the treatment of complicated urinary tract infections, including acute pyelonephritis: RECAPTURE, a phase 3 randomized trial program. *Clin Infect Dis* 63:754–762. <https://doi.org/10.1093/cid/ciw378>.
- Mazuski JE, Gasink LB, Armstrong J, Broadhurst H, Stone GG, Rank D, Llorens L, Newell P, Pacht J. 2016. Efficacy and safety of ceftazidime-avibactam plus metronidazole versus meropenem in the treatment of complicated intra-abdominal infection—results from a randomized, controlled, double-blind, phase 3 program. *Clin Infect Dis* 62:1380–1389. <https://doi.org/10.1093/cid/ciw133>.
- Carmeli Y, Armstrong J, Laud PJ, Newell P, Stone G, Wardman A, Gasink LB. 2016. Ceftazidime-avibactam or best available therapy in patients with ceftazidime-resistant *Enterobacteriaceae* and *Pseudomonas aeruginosa* complicated urinary tract infections or complicated intra-abdominal infections (REPRISE): a randomised, pathogen-directed, phase 3 study. *Lancet Infect Dis* 16:661–673. [https://doi.org/10.1016/S1473-3099\(16\)30004-4](https://doi.org/10.1016/S1473-3099(16)30004-4).
- Mendes RE, Castanheira M, Gasink L, Stone GG, Nichols WW, Flamm RK, Jones RN. 2015. β -Lactamase characterization of Gram-negative pathogens recovered from patients enrolled in the phase 2 trials for ceftazidime-avibactam: clinical efficacies analyzed against subsets of molecularly characterized isolates. *Antimicrob Agents Chemother* 60:1328–1335. <https://doi.org/10.1128/AAC.01173-15>.
- Mendes RE, Castanheira M, Woosley LN, Stone GG, Bradford PA, Flamm RK. 2017. Molecular β -lactamase characterization of aerobic Gram-negative pathogens recovered from patients enrolled in the ceftazidime-avibactam phase 3 trials for complicated intra-abdominal infections, with efficacies analyzed against susceptible and resistant subsets. *Antimicrob Agents Chemother* 61:e02447-16. <https://doi.org/10.1128/AAC.02447-16>.
- Jones CH, Tuckman M, Keeney D, Ruzin A, Bradford PA. 2009. Characterization and sequence analysis of extended-spectrum- β -lactamase-encoding genes from *Escherichia coli*, *Klebsiella pneumoniae*, and *Proteus mirabilis* isolates collected during tigecycline phase 3 clinical trials. *Antimicrob Agents Chemother* 53:465–475. <https://doi.org/10.1128/AAC.00883-08>.
- Castanheira M, Mills JC, Costello SE, Jones RN, Sader HS. 2015. Ceftazidime-avibactam activity tested against *Enterobacteriaceae* isolates from U.S. hospitals (2011 to 2013) and characterization of β -lactamase-producing strains. *Antimicrob Agents Chemother* 59:3509–3517. <https://doi.org/10.1128/AAC.00163-15>.
- Jean SS, Hsueh PR, SMART Asia-Pacific Group. 2017. Distribution of ESBLs, AmpC beta-lactamases and carbapenemases among *Enterobacteriaceae* isolates causing intra-abdominal and urinary tract infections in the Asia-Pacific region during 2008–14: results from the Study for Monitoring Antimicrobial Resistance Trends (SMART). *J Antimicrob Chemother* 72:166–171. <https://doi.org/10.1093/jac/dkw398>.
- Bevan ER, Jones AM, Hawkey PM. 2017. Global epidemiology of CTX-M beta-lactamases: temporal and geographical shifts in genotype. *J Antimicrob Chemother* 72:2145–2155. <https://doi.org/10.1093/jac/dkx146>.
- Doi Y, Iovleva A, Bonomo RA. 2017. The ecology of extended-spectrum beta-lactamases (ESBLs) in the developed world. *J Travel Med* 24:S44–S51. <https://doi.org/10.1093/jtm/taw102>.

23. Pfaller MA, Bassetti M, Duncan LR, Castanheira M. 2017. Ceftolozane/tazobactam activity against drug-resistant *Enterobacteriaceae* and *Pseudomonas aeruginosa* causing urinary tract and intraabdominal infections in Europe: report from an antimicrobial surveillance programme (2012–15). *J Antimicrob Chemother* 72:1386–1395. <https://doi.org/10.1093/jac/dkx009>.
24. Albiger B, Glasner C, Struelens MJ, Grundmann H, Monnet DL, European Survey of Carbapenemase-Producing *Enterobacteriaceae* Working Group. 2015. Carbapenemase-producing *Enterobacteriaceae* in Europe: assessment by national experts from 38 countries, May 2015. *Euro Surveill* 20(45):pii=30062. <https://doi.org/10.2807/1560-7917.ES.2015.20.45.30062>.
25. CLSI. 2015. M07-A10. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard, 10th ed. CLSI, Wayne, PA.
26. Bush K, Jacoby GA. 2010. Updated functional classification of beta-lactamases. *Antimicrob Agents Chemother* 54:969–976. <https://doi.org/10.1128/AAC.01009-09>.